

Monroig O, Navarro JC, Amat F & Hontoria F (2007) Enrichment of *Artemia nauplii* in vitamin A, vitamin C and methionine using liposomes, *Aquaculture*, 269 (1-4), pp. 504-513

This is the peer reviewed version of this article

NOTICE: this is the author's version of a work that was accepted for publication in Aquaculture. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Aquaculture, [VOL 269, ISS 1-4 (2007)] DOI: <http://dx.doi.org/10.1016/j.aquaculture.2007.02.056>

Elsevier Editorial System(tm) for Aquaculture

Manuscript Draft

Manuscript Number:

Title: Enrichment of Artemia nauplii in vitamin A, vitamin C and methionine using liposomes

Article Type: Research Paper

Keywords: Artemia enrichment; liposomes; methionine; vitamin A; vitamin C

Corresponding Author: Dr. Francisco Hontoria,

Corresponding Author's Institution: Instituto de Acuicultura de Torre de la Sal (CSIC)

First Author: Óscar Monroig, Dr.

Order of Authors: Óscar Monroig, Dr.; Juan Carlos Navarro, Dr.; Francisco Amat, Dr.; Francisco Hontoria

Abstract: Several types of liposomes were used to enrich Artemia nauplii in vitamin A, vitamin C and free methionine. In a first experiment, unilamellar liposomes formulated with krill phospholipid extract and retinyl palmitate demonstrated their capability to enhance the retinol content of Artemia nauplii. Furthermore, the increase in retinol was related to the amount of retinyl palmitate included in the liposomes as vitamin A source. These findings yield the possibility of using such vesicles to bioencapsulate simultaneously both vitamin A and essential fatty acids present in the krill phospholipid extract. A second enrichment was carried out with unilamellar liposomes composed of soybean phosphatidylcholine and loaded with sodium ascorbate as vitamin C source. Our results did not show that vitamin C content in the nauplii could be increased using unilamellar liposomes. This was most likely due to the degradation of the vitamin C during enrichment as well as the ascorbate leakage. Finally, a third experiment assessed enrichment in free methionine using liposomes of different lamellarity (unilamellar or multilamellar) and composed of either soybean phosphatidylcholine or dipalmitoyl phosphatidylcholine, both combined with cholesterol as a membrane stabilizer. Results indicated that multilamellar liposomes represent a useful tool to deliver methionine to Artemia nauplii. Enhanced protection given by their multiple bilayers in comparison to unilamellar liposomes could account for the higher ability displayed by multilamellar vesicles for free methionine bioencapsulation.

1 Title:

2 Enrichment of *Artemia* nauplii in vitamin A, vitamin C and methionine using liposomes

3

4 Authors:

5 Óscar Monroig, Juan Carlos Navarro, Francisco Amat, Francisco Hontoria

6

7 Affiliation:

8 Instituto de Acuicultura de Torre de la Sal (CSIC). 12595 Torre de la Sal.

9 Castellón, Spain

10

11 Correspondence address:

12 Francisco Hontoria. Instituto de Acuicultura de Torre de la Sal (CSIC). 12595 Torre de

13 la Sal. Castellón. Spain. Telephone +34 964 319500. Fax +34 964 319509.

14 E-mail: hontoria@iats.csic.es

15

16 **Abstract**

17 Several types of liposomes were used to enrich *Artemia* nauplii in vitamin A,
18 vitamin C and free methionine. In a first experiment, unilamellar liposomes formulated
19 with krill phospholipid extract and retinyl palmitate demonstrated their capability to
20 enhance the retinol content of *Artemia* nauplii. Furthermore, the increase in retinol was
21 related to the amount of retinyl palmitate included in the liposomes as vitamin A source.
22 These findings yield the possibility of using such vesicles to bioencapsulate
23 simultaneously both vitamin A and essential fatty acids present in the krill phospholipid
24 extract. A second enrichment was carried out with unilamellar liposomes composed of
25 soybean phosphatidylcholine and loaded with sodium ascorbate as vitamin C source.

26 Our results did not show that vitamin C content in the nauplii could be increased using
27 unilamellar liposomes. This was most likely due to the degradation of the vitamin C
28 during enrichment as well as the ascorbate leakage. Finally, a third experiment assessed
29 enrichment in free methionine using liposomes of different lamellarity (unilamellar or
30 multilamellar) and composed of either soybean phosphatidylcholine or dipalmitoyl
31 phosphatidylcholine, both combined with cholesterol as a membrane stabilizer. Results
32 indicated that multilamellar liposomes represent a useful tool to deliver methionine to
33 *Artemia* nauplii. Enhanced protection given by their multiple bilayers in comparison to
34 unilamellar liposomes could account for the higher ability displayed by multilamellar
35 vesicles for free methionine bioencapsulation.

36

37 Keywords: *Artemia* enrichment; liposomes; methionine; vitamin A; vitamin C

38 **1. Introduction**

39 Most of the studies on nutrition of marine fish larvae focus their attention on the
40 importance of essential fatty acids (Izquierdo, 1996; Sargent et al., 1997). However,
41 vitamins and free amino acids are also considered as essential nutrients, and therefore
42 they have to be included in the diet for satisfying the requirements to maintain the
43 different biological functions working correctly, and guarantee the normal growth
44 (Rønnestad et al., 1999; Halver, 2002).

45 Vitamins are a heterogeneous group of compounds with a wide range of
46 biological activities. Vitamins are classified according to their solubility, as either lipid
47 or water soluble compounds (Gouillou-Coustans and Guillaume, 2001). Among
48 liposoluble vitamins, vitamin A represents an essential nutrient for fish since they
49 cannot synthesize this compound *de novo*. Vitamin A has been widely studied in fish
50 because it is involved in vision and most of the fish reared species are visual feeders

51 (Hunter, 1981). Consequently, dietary deficiencies in vitamin A can alter normal
52 function of vision in larvae, thereby reducing their predator ability. Moreover, dietary
53 vitamin A deficiencies have been related to the occurrence of pseudoalbinism in flatfish
54 such as halibut (*Hippoglossus hippoglossus*), turbot (*Psetta maxima*) and Japanese
55 flounder (*Paralichthys olivaceus*). This syndrome is characterized by the lack of
56 pigmentation in the body skin of the individual (Seikai et al., 1987; Estévez and
57 Kanazawa, 1995). Furthermore, low vitamin A content in the live preys as *Artemia* can
58 lead to an incomplete migration of the eye during flat fish metamorphosis (Saele et al.,
59 2003). On the contrary, hypervitaminosis of vitamin A has been related to the backbone
60 disease in Japanese flounder (Dedi et al., 1995, 1997), together with the jaw
61 deformation (Haga et al., 2003).

62 In regard to hydrosoluble vitamins, vitamin C or ascorbic acid is a dietary
63 essential nutrient for fish since these animals lack the enzyme gulonolactone oxidase
64 required in its synthesis (Dabrowski, 1990). In addition to its essentiality, vitamin C
65 also offers antioxidant (Hwang and Lin, 2002) and immunostimulant (Cuesta et al.,
66 2002) properties, and protection against stress (Henrique et al., 2002). For these reasons,
67 most diets used in aquaculture contain megadoses of vitamin C as a strategy to prevent
68 diseases in cultured fish.

69 Amino acids are believed to be important dietary nutrients for early larval stages
70 because it has been observed to be present in high amounts in fish eggs and in marine
71 zooplankton, the food of the larvae (Rønnestad, 1992, Fyhn et al., 1993; Rønnestad et
72 al., 1999). The main properties of free amino acids include their contribution to the
73 formation of new tissues during body growth, their role as energy source supply
74 (Rønnestad et al., 1999), and as effectors of chemical attraction (Kolkovski et al., 1997;
75 Hara, 2006). Among essential amino acids, methionine can be considered to be a

76 limiting amino acid since its content in *Artemia* nauplii is only at trace or low levels
77 (Helland et al., 2000).

78 Despite the fact that *Artemia* nauplii contain vitamin A, vitamin C and free
79 amino acids (Merchie et al., 1997; Takeuchi et al., 1998; Helland et al., 2000),
80 enrichment products normally include these compounds to enhance natural content of
81 nauplii. However, inclusion of these compounds in the enrichment diet is somewhat
82 difficult because of their different polarity. In the case of fish oil emulsions, one of the
83 most habitual products employed in *Artemia* enrichment worldwide, lipid micelles
84 formed when the product is dispersed in water possess a lipophilic core which impedes
85 the incorporation of hydrosoluble molecules. In the case of vitamin C, this handicap has
86 been overcome using lipophilic derivatives such as ascorbyl palmitate which is stable in
87 the emulsion micelles (Merchie et al., 1995a, b). Finally, amino acids are water soluble
88 compounds and, consequently, their inclusion in lipid emulsions seems to be difficult.

89 Liposomes represent a potential alternative for the enrichment of live preys in
90 both hydrophilic and lipophilic nutrients (Hontoria et al., 1994). Simply stated,
91 liposomes are lipid vesicles enclosing an aqueous space (New, 1990). According to this
92 structure, liposomes can be formulated with water-soluble compounds dissolved in the
93 aqueous space and lipophilic nutrients imbibed in the apolar moiety of phospholipid
94 fatty acid chains. Previous studies have assessed the capacity of liposomes to deliver
95 different compounds to live preys used in marine larviculture (Hontoria et al., 1994;
96 Ozkizilcik and Chu, 1994; Touraki et al., 1995; Tonheim et al., 2000; Monroig et al.,
97 2006).

98 The present study assesses several types of liposomes in the enrichment of
99 *Artemia* nauplii in vitamin A, vitamin C and methionine as models for lipophilic
100 vitamins, hydrophilic vitamins and amino acids, respectively.

101 **2. Materials and methods**

102 *2.1. Preparation of liposomes*

103 Liposomes prepared by means of three different methodologies were employed to
104 deliver vitamin A, vitamin C and methionine. Firstly, multilamellar vesicles (MLV)
105 were prepared by the method proposed by Bangham et al. (1965), but using filtered
106 seawater as the aqueous phase. Secondly, large unilamellar vesicles (LUV) were
107 obtained by extrusion (LUVext) of MLV suspensions using an extruder LiposoFast
108 equipped with 100 nm polycarbonate membranes (Avestin Inc., Ottawa, ON, Canada).
109 Finally, a second type of unilamellar liposomes was prepared by a detergent
110 solubilization methodology (LUVdet). In this latter typology, the aqueous phase
111 consisted of a saline solution (0.9% NaCl). More details about these methodologies are
112 found in Monroig et al. (2006).

113 Membranes of all three liposome typologies (MLV, LUVext and LUVdet) were
114 formulated with three different phospholipid sources purchased from Avanti Polar
115 Lipids (Alabaster, AL, USA): krill phospholipid extract (KPE), soybean
116 phosphatidylcholine (SPC), and dipalmitoyl phosphatidylcholine (DPPC). Lipid class
117 and fatty acid composition of phospholipids are presented in Table 1. Selection of the
118 membrane components and liposome typology was adopted according to the lipophilic
119 (retinyl palmitate) or hydrophilic (ascorbate and methionine) nature of the delivered
120 nutrient. Hence, liposomes chosen to enrich *Artemia* in vitamin A were composed of
121 KPE since this combination, although potentially leaky, would permit the simultaneous
122 enrichment in both essential fatty acids and vitamin A. Contrarily, vitamin C and
123 methionine were encapsulated in SPC and DPPC vesicles, which had demonstrated to
124 efficiently retain water-soluble substances when submitted to *Artemia* enrichment
125 conditions (Monroig et al., 2003). Occasionally, cholesterol (CHO) from Sigma

126 (Alcobendas, Spain) was included as a membrane stabilizer (New, 1990, p. 22).

127 Relevant features of the liposome suspensions used in this study are gathered in Table 2.

128 2.2. Enrichment of *Artemia nauplii* in vitamin A

129 Two types of unilamellar liposomes were used to enhance vitamin A of *Artemia*
130 nauplii, together with the commercial emulsion Super Selco (Inve, Ghent, Belgium) as a
131 reference product containing retinyl palmitate (Table 2). Both liposome suspensions
132 were formulated with KPE and retinyl palmitate in a ratio of 100:1 (w/w) in LUVext
133 (LUVext A), and 100:2 (w/w) in LUVdet (LUVdet A).

134 Commercial cysts (EG grade Inve, Ghent, Belgium) were incubated in aerated
135 seawater at a temperature of $28 \pm 1^\circ\text{C}$ with an illumination of 1500-2000 lux. The cyst
136 incubation lasted for 23 h. During the following hour, the nauplii were separated from
137 the cyst shells and stocked at approximate densities of 300 nauplii ml^{-1} in 1 l cylinder-
138 conical enrichment tubes. At this point the nauplii were ready to begin the enrichment
139 process (t_0). Afterwards, the enrichment products were dispensed in a single dose
140 yielding retinyl palmitate concentrations of 1.3, 5.0 and 10.0 mg l^{-1} corresponding to
141 Emulsion, LUVext A and LUVdet A, respectively. Similarly to the cyst incubation, the
142 tubes were placed in a thermostated bath at a temperature of $28 \pm 1^\circ\text{C}$ and illuminated
143 between 1500-2000 lux. Aeration was applied in each enrichment tube with 0.5
144 diameter section tubes, thereby yielding an approximate air flow of 1.0 litre per minute.

145 Samples of enriched nauplii were collected in a 100 μm plankton mesh and
146 gently washed with tap water, after 18 or 21 h of incubation in the enrichment medium.
147 Part of the nauplii were kept in clean seawater at the same conditions of nauplii
148 enrichment incubation stated above for 3 h more (24 h since the start of the incubation),
149 and then sampled. This sample point reflects the nutritional value of nauplii after some

150 hours in the culture tank, and emulates food quality in normal hatchery procedures.
151 Enriched nauplii samples were immediately frozen until nutrient analyses performed.

152 2.3. Enrichment of *Artemia nauplii* in vitamin C

153 Newly obtained *Artemia* nauplii were incubated with four enrichment diets
154 (Table 2). In addition to the commercial emulsion Super Selco (Emulsion) that contains
155 ascorbyl palmitate as vitamin C source, *Artemia* nauplii were incubated with LUVdet
156 liposomes composed of SPC as a unique membrane component and prepared with
157 sodium ascorbate (Sigma, Alcobendas, Spain) dissolved in the aqueous phase at a
158 concentration of 100 mg ml⁻¹. Liposomes were administered in a single dose at the
159 beginning of the incubation (t₀) (LUVdet C 1) or in two separate doses at t₀ and t₀+7 h
160 (LUVdet C 2). Finally, a fourth treatment consisted of sodium ascorbate directly
161 dissolved in the enrichment medium (Solution C). Both liposomes and solution
162 treatments yielded a sodium ascorbate concentration of 0.5 g l⁻¹, whereas the ascorbyl
163 palmitate content of the commercial emulsion was not specified by the manufacturer.
164 Sampling was carried out as described in the vitamin A enrichment.

165 2.4. Enrichment of *Artemia nauplii* in methionine

166 Four liposome-based treatments were used to enhance the free methionine
167 content of *Artemia* nauplii (Table 2). LUVext formulated with SPC and CHO (4:1, w/w)
168 as membrane components and methionine dissolved in the aqueous phase (40 mg ml⁻¹)
169 were administered in a single dose at t₀ (LUVext M 1) or in two separate doses at t₀ and
170 t₀+8 h (LUVext M 2). A third enrichment product consisted of LUVext composed of
171 DPPC and CHO (4:1, w/w) and methionine dissolved in the aqueous phase (40 mg ml⁻¹)
172 (LUVext M 3) administered in a single dose at the start of the nauplii incubation (t₀).
173 Liposome treatments were completed with multilamellar vesicles (MLV M) with an
174 equal composition than LUVext M 1 and LUV M 2, and administered in a single dose at

175 the beginning of incubation. As performed in the vitamin C experiment, liposomes were
176 compared with the efficiency displayed by methionine directly dissolved in the
177 enrichment medium and dispensed at t_0 (Solution M). In all five treatments,
178 concentration of methionine in the enrichment medium was 0.2 g l^{-1} after administration
179 of the total amount of product.

180 *2.5. Determination of vitamin A*

181 Analysis of retinol was carried out following the method proposed by Takeuchi
182 et al. (1998) with the following modifications. Total lipids were extracted from
183 biological samples (around 10 mg DW) according to the method described by Folch et
184 al. (1957), and then transferred to amber vials to avoid photodegradation. Lipid extracts
185 were dissolved in a methanol:acetone solution (1:1, v/v) containing $3.0 \text{ } \mu\text{g ml}^{-1}$ of
186 retinyl acetate (Sigma, Alcobendas, Spain) used as an internal standard (Aust et al.,
187 2001).

188 Lipid extracts (20 μl) were injected in a high performance liquid
189 chromatography (HPLC) system (Thermo Liquid Chromatography, San Jose, CA,
190 USA) equipped with a quaternary pump P4000 employed at isocratic conditions.
191 Detection was performed using a UV-VIS UV6000LP detector set at a wavelength of
192 325 nm. An oven Croco-Cil (Cluzeau, Sainte-Foy-la-Grande, France) was used to keep
193 a C-18 Lichrospher (Merck, Darmstadt, Germany) at 30°C during the elution. The
194 mobile phase consisted of 98% methanol containing 0.5% ammonium acetate (w/v) and
195 chloroform 85:15 (v/v) which was pumped at a flow rate of 1.5 ml min^{-1}

196 Peak identification was performed by comparing with commercial standards
197 obtained from Sigma (Alcobendas, Spain) and quantification of each peak was carried
198 out by comparing its relative area with that of the internal standard retinyl acetate.

199 ChromQuest Software (Thermo Quest, San Jose, CA, USA) was used in peak
200 integration.

201 *2.6. Determination of vitamin C*

202 Analyses of ascorbate (vitamin C) were performed following the method
203 described by Shiau and Hsu (1994). A weighed portion of enriched nauplii (around 10
204 mg DW) was suspended in 2 ml of ice cold 5% (w/v) metaphosphoric acid (Sigma,
205 Alcobendas, Spain) and homogenized with a tissue disrupter (Ika Labortechnik, Staufen,
206 Germany) for 30 s in an ice bath. Homogenates were centrifuged at 3000 g for 5 min,
207 and supernatants were filtered through a 0.45 µm-pore-size glass fiber-nylon syringe
208 filter (Whatman, Clifton, NJ, USA).

209 Ascorbate extracts (20 µl) were injected in the same HPLC system as described
210 in the vitamin A analysis. In this case, mobile phase consisting of an aqueous solution
211 of 0.05 M KH₂PO₄ (pH 2.8) was pumped at a flow rate of 0.9 ml min⁻¹. Elution was
212 carried out at room temperature and the relative peak area was compared with ascorbic
213 acid (Sigma, Alcobendas, Spain) as an external standard using a UV–VIS detector set at
214 254 nm. ChromQuest Software (Thermo Quest, San Jose, CA, USA) was used in peak
215 integration.

216 *2.7. Determination of free methionine*

217 A modification of the method described by Koop et al. (1982) was used in the
218 free amino acid determination. This method is based on a pre-column derivatization
219 with phenylisothiocyanate (PITC, Sigma, Alcobendas, Spain). Samples of enriched
220 nauplii (around 10 mg DW) were extracted in 1 ml of 6% (w/v) trichloroacetic acid
221 (Panreac, Castellar del Vallés, Spain) using a tissue disrupter (Ika Labortechnik, Staufen,
222 Germany). After maintaining homogenates for 24 h at 4°C, they were centrifuged

223 (10000 g) for 20 min, and supernatants were filtered through a 0.45 μm -pore-size glass
224 fiber-nylon syringe filter (Whatman, Clifton, NJ, USA).

225 Derivatization of free amino acids was accomplished after mixing 125 μl of the
226 filtrate and 25 μl of 2 mM of norleucine used as an internal standard (Sigma,
227 Alcobendas, Spain) with the following procedure. After desiccation of the mixture in a
228 speed vacuum, 20 μl of methanol:distilled water:trimethylamine (2:2:1, v/v/v) were
229 added to facilitate complete desiccation, and again introduced in the speed vacuum.
230 Finally, 15 μl of freshly prepared derivatizing reagent composed of methanol:distilled
231 water:trimethylamine:PITC (7:1:1:1, v/v/v/v) were added and incubated for 20 min at
232 room temperature. After this period, derivatized samples were completely desiccated in
233 the speed vacuum. At this point, samples were preserved at -20°C until HPLC injection.

234 Sample injection (20 μl) was carried out after dissolving derivatized samples
235 with 200 μl of 5 mM Na_2HPO_4 (pH 7.4) containing 5% (v/v) acetonitrile. A C-18
236 Lichrospher (Merck, Darmstadt, Germany) column maintained at 52°C during the
237 elution was employed. Injected samples were eluted through a gradient mixture of two
238 mobile phases: mobile phase A consisted of an aqueous solution of 70 mM sodium
239 acetate (pH 6.55) containing 2.5% (v/v) acetonitrile; and mobile phase B was composed
240 of acetonitrile:distilled water:methanol (45:40:15, v/v/v). All solvents were HPLC
241 grade (Merck, Darmstadt, Germany). The flow rate of the mobile phase was 1 ml min^{-1} ,
242 and followed the gradient program detailed in Table 3. The same UV-VIS detector as
243 described above was employed at 254 nm. Peaks were identified by comparison with
244 the amino acid commercial standard AA-S-18 (Sigma, Alcobendas, Spain), whereas
245 quantification was obtained by comparing its relative area with that of the internal
246 standard norleucine. ChromQuest Software (Thermo Quest, San Jose, CA, USA) was
247 used in peak integration.

248 2.8. Statistical analyses

249 Analytical data are expressed as means \pm standard deviations. Homogeneity of
250 variances was checked by Barlett's test. In cases of homoscedasticity, differences in the
251 nutrient content among treatments in each sampling point were analyzed by one-way
252 analysis of variance (ANOVA) followed, where appropriate, by Tukey's multiple
253 comparison test (Sokal and Rohlf, 1981). If heterogeneity of variances existed, robust
254 tests were applied. Welch's test was used to check differences between treatments and
255 Game-Howell's test to establish differences among groups. Comparisons of the means
256 with P values equal or less than 0.05 were considered significantly different. All the
257 statistical analyses were carried out using the SPSS statistical package (SPSS Inc.,
258 Chicago, IL, USA).

259 3. Results

260 3.1. Enrichment of *Artemia nauplii* in vitamin A

261 Fig. 1 shows the retinol content of nauplii enriched with two different liposome
262 preparations and the commercial emulsion. A relationship between the amount of
263 retinyl palmitate present in the enrichment diet and the retinol measured in the treated
264 nauplii can be observed. Thus, nauplii enriched with LUVdet A displayed higher retinol
265 levels than the other two treatments in all three sampling points ($P \leq 0.05$). At the same
266 time, LUVext A nauplii show higher retinol concentration than nauplii enriched with
267 the commercial emulsion after 18 and 21 h of incubation, although no differences can
268 be seen at the 24 h sampling point ($P \leq 0.05$). It is noteworthy to mention that nauplii
269 enriched with the emulsion, that includes retinyl palmitate in its composition, exhibit a
270 retinol contents below the detection threshold. In regards to retinol incorporation
271 throughout enrichment, both liposome-based treatments did not exhibit important

272 changes from 18 to 21 h. Nonetheless, the subsequent starvation period (24 h sampling
273 point) produced a drop in the retinol content of liposome enriched nauplii to the extent
274 that retinol from LUVext A nauplii was below detection threshold at the 24 h sampling
275 point (Fig. 1).

276 3.2. *Enrichment of Artemia nauplii in vitamin C*

277 Ascorbate analyses from nauplii enriched with the different products are shown
278 in Fig. 2. In general, nauplii enriched with the commercial emulsion contained lower
279 ascorbate levels in comparison to the other treatments. This difference became
280 significant ($P \leq 0.05$) at 18 h of incubation. Another remarkable result is the lack of
281 significant differences at any sampling point ($P \leq 0.05$) between the ascorbate content of
282 nauplii enriched with liposomes administered in a single dose (LUVdet C 1) and two
283 doses (LUVdet C 2). Nauplii incubated with dissolved ascorbate (Solution C) showed
284 not only statistically equivalent levels than the other treatments, but also their ascorbate
285 content was even higher ($P \leq 0.05$) than the other's after the starvation period.
286 Concerning the temporal incorporation, ascorbate levels from nauplii did not undergo
287 conclusive patterns, except for the increase observed during the starvation period in
288 nauplii treated with dissolved ascorbate (Fig. 2).

289 3.3. *Enrichment of Artemia nauplii in free methionine*

290 Levels of free methionine measured in *Artemia* nauplii incubated with different
291 enrichment products are represented in Fig. 3. After 18 h of incubation, nauplii enriched
292 with MLV liposomes (MLV M) registered significantly higher contents of free
293 methionine in comparison to the other treatments ($P \leq 0.05$). Such a difference was
294 maintained for up to 21 h, although LUVext M 2 (2 doses) did not differ from MLV
295 nauplii ($P \leq 0.05$) at that sampling point. Although this tendency is also observed at 24 h,

296 free methionine of nauplii enriched with MLV liposomes only showed significant
297 higher contents if compared to LUVext M 1 and Solution M nauplii. As seen in the
298 retinol and ascorbate experiments, temporal evolution of the free methionine levels of
299 nauplii showed minor changes from 18 to 21 h, except for LUVext M 1 nauplii, which
300 seemed to exhibit an increase during this period. On the contrary, the starvation period
301 was accompanied by a general decrease in the free methionine visible in all five
302 treatments.

303 **4. Discussion**

304 The ability of liposomes to deliver both hydrophilic and lipophilic substances
305 has been employed to improve the nutritional value of live preys used for feeding
306 marine fish larvae (Ozkizilcik and Chu, 1994; McEvoy et al., 1996; Tonheim et al.,
307 2000). With the exception of the pioneer study undertaken by Hontoria et al. (1997) on
308 vitamin encapsulation with liposomes, the present study represents the first report of
309 vitamin A bioencapsulation in live preys by means of these lipid vesicles. Some
310 nutritional studies on flat fish larvae have assessed the efficiency of several compounds
311 with vitamin A activity (i.e., retinol, retinyl esters, etc.) included in oil emulsions and
312 bioencapsulated in *Artemia* nauplii (Dedi et al., 1995; Estévez and Kanazawa, 1995;
313 Takeuchi et al., 1998; Haga et al., 2004). Results obtained in the present study illustrate
314 the efficiency of liposomes formulated with krill phospholipids and retinyl palmitate to
315 enhance the retinol content in *Artemia* nauplii. Additionally, the amount of retinol in the
316 nauplii can be directly related to the retinyl ester included in the liposome membrane,
317 thereby pointing at a metabolic transformation by naupliial enzymatic system as
318 previously described by Takeuchi et al. (1998). Retinyl palmitate incorporated in
319 liposomes would be hydrolyzed to free retinol before its absorption in the intestinal cells
320 (Furr and McGrane, 2003). Metabolic activity of nauplii may also explain the decrease

321 in the retinol levels observed throughout the starvation period in liposome enriched
322 nauplii, probably due to a reesterification of retinol (Furr and McGrane, 2003) within
323 digestive tract cells. A shocking result is the low retinol level obtained in nauplii
324 incubated with the commercial emulsion despite containing retinyl palmitate. This poor
325 efficiency of the commercial emulsion may, in part, be due to a partial degradation
326 during the incubation of the enrichment product under aggressive conditions of light,
327 oxygen and temperature (Woollard and Indyk, 2003). Indeed, Moren et al. (2005) have
328 recently proposed this degradation mechanism to explain why no vitamin A was
329 detected when another enrichment product (DC-DHA Selco) was submitted to
330 simulated enrichment conditions during 12 h. Despite the fact that liposomes could
331 undergo the same process, the higher amount of retinyl palmitate present in liposomes
332 in comparison to the commercial emulsion would preserve enough quantity to be
333 bioencapsulated in the *Artemia* nauplii before degradation.

334 Regarding the vitamin C enrichment, our results revealed that enhancement in
335 the ascorbate content of *Artemia* nauplii was more efficient when ascorbate was
336 encapsulated in liposomes or simply dissolved in the water in comparison to the oil
337 emulsion formulated with ascorbyl palmitate. The importance of such a result is
338 difficult to evaluate since the amount of vitamin C available in all treatments was not
339 the same. Whereas nauplii treated with liposomes and the aqueous solution were
340 incubated at the same initial ascorbate concentration in the medium (0.5 g l^{-1}), the
341 amount of vitamin C available for nauplii enriched with the commercial emulsion
342 depends on the concentration of ascorbyl palmitate present in the product (not specified
343 by the manufacturer), along with the efficiency of the nauplii to hydrolyze the ester
344 derivative to free ascorbate (Merchie et al., 1995a). However, results do not prove the
345 presumable advantage of using liposomes instead of the aqueous solution, since

346 differences between liposomes and the aqueous solution did not exist. Firstly, the low
347 efficiency of liposomes to bioencapsulate ascorbate in *Artemia* can be partly explained
348 through the susceptibility of ascorbate to degrade (Halver, 2002) as occurred in vitamin
349 A. Preservation of vitamin C by means of other antioxidant compounds such as vitamin
350 E included in the membrane (Urano et al., 1987; Fukuzawa et al., 1993a, b) or the
351 utilization of more resistant derivatives (Schüep et al., 1989; Shiao and Hsu, 1994;
352 Yoshitomi, 2004) would improve the efficiency of liposomes as an ascorbate carrier to
353 *Artemia* nauplii. Secondly, the turbulent regime of enrichment would produce the
354 leakage of water-soluble compounds encapsulated in liposomes. This phenomenon,
355 which is expected to affect other hydrosoluble substances such as methionine, is
356 discussed later on.

357 Results from the methionine enrichment indicate the possibility of using
358 liposomes to enhance the free amino acid pool in *Artemia* nauplii. Nauplii enriched with
359 multilamellar liposomes (MLV) presented higher levels of free methionine than
360 unilamellar liposomes (LUV), despite the fact that LUV are considered to have the most
361 efficient typology to maximize encapsulation of water-soluble compounds because of
362 the high internal volume:membrane lipid ratio (New, 1990, p. 28). Nevertheless, under
363 high turbulence conditions of enrichment procedures, MLV formulations seem to be
364 more suitable than LUV possibly due to their protective multiple bilayer structure
365 (between 5 and 20 layers). Therefore, methionine dissolved in the more internal
366 intermembrane spaces of MLV would remain encapsulated, whereas the turbulence
367 during incubation would cause the leakage of water-soluble compounds from the
368 aqueous compartments located between the more outer membranes, as would also occur
369 in the unique aqueous space of LUV. This low performance of LUV could neither be
370 improved through the dosage of liposomes in two separate doses, despite the fact that

371 this strategy would increase the time during which vesicles are still loaded with the
372 water-soluble nutrient (Monroig et al., 2003). Consequently, bioencapsulation of
373 methionine loaded particles (MLV liposomes) would be favoured in comparison to
374 methionine simply dissolved in the enrichment medium (case of original methionine
375 solution and empty unilamellar liposomes) since filtering appendages of nauplii are
376 adapted to retain discrete particles of a certain size present in the water column.

377 The direct incorporation of methionine through ingestion of enrichment medium
378 by nauplii should also be considered, as *Artemia* nauplii drink medium for
379 osmoregulatory balance (Navarro et al., 1993). Considering the content of free
380 methionine from newly hatched nauplii ($88.4 \mu\text{g g DW}^{-1}$) as the basal level, free
381 methionine measured in nauplii enriched with the methionine solution (Solution M)
382 (between 300 and $400 \mu\text{g g DW}^{-1}$) possibly indicate a net incorporation of the amino
383 acid through ingestion of the medium. The existence of this direct pathway was
384 demonstrated in *Artemia* enrichments with radioactive methionine (Tonheim et al.,
385 2000). In regards to temporal incorporation, analyses did not indicate an important
386 increase in the free methionine content during the period between 18 and 21 h. Despite
387 the supply of methionine during this period, the free methionine already
388 bioencapsulated in the nauplii could not be detected since nauplii can assimilate it into
389 the protein structure, transform it into other amino acids (cysteine or taurine), and
390 oxidize it to sulphate products (Tonheim et al., 2000). Moreover, these mechanisms
391 could still occur during the starvation period (from 21 to 24 h), thus explaining the drop
392 of free methionine levels in all treatments. Indeed, total absorbed methionine was
393 estimated to be two fold compared to that present in a free state (Tonheim et al., 2000).

394 In light of these results, unilamellar vesicles cannot be considered the best
395 alternative to encapsulate hydrophilic nutrients in liposomes employed in *Artemia*

396 enrichments, contrarily to the results obtained from another study (Monroig et al., 2003)
397 where LUV suspensions retained a fluorescent marker during a longer period than MLV
398 liposomes. This apparent discrepancy can be related to several reasons. Firstly, MLV
399 suspensions tested by Monroig et al. (2003) were formulated with krill phospholipids
400 which would form leaky vesicles due to the unsaturation degree of their fatty acid
401 chains (Alberts et al., 1996, p.512-513). On the contrary, formulation of multilamellar
402 liposomes with purified soybean phosphatidylcholine in the present study would
403 improve the retention of water-soluble substances included in these vesicles. Secondly,
404 differences in the hydrodynamic conditions set in both studies could also explain the
405 different behaviour of multilamellar liposomes. Incubation of nauplii in the present
406 study was carried out in 1 l tubes with an aeration flow around 1.0 lpm, whereas we
407 previously used smaller tubes (200 ml) where aeration was reduced to simulate normal
408 turbulence applied in standard 1 l enrichment tubes (Monroig et al., 2003). While
409 attempting to standardize turbulence in the 200 ml tubes, the degree of agitation in the
410 enrichment medium in this study could have resulted to be higher than the previous
411 study by Monroig et al. (2003). Finally, differences between the solubility of the
412 fluorescent marker carboxyfluorescein used in our preliminary experiment (Monroig et
413 al., 2003) and the water-soluble substances used here, such as methionine, could affect
414 their leakage rate and, consequently, make results from both studies not fully
415 comparable.

416 In summary, liposomes can be presented as an efficient system for the
417 enrichment of *Artemia* nauplii in retinol. Furthermore, retinol incorporation was related
418 to the amount of retinyl palmitate included in the liposome formulation. These findings
419 demonstrate the potential of these vesicles to bioencapsulate simultaneously both
420 vitamin A and essential fatty acids present in the krill phospholipid extract. Regarding

421 water-soluble nutrients, the present study could not demonstrate the efficiency of
422 unilamellar liposomes to increase the vitamin C content of nauplii probably due to
423 degradation during the enrichment process. The use of more stable derivatives, together
424 with the combination with other antioxidant compounds could improve liposome
425 capacity for vitamin C enrichment. In addition, leakage of vitamin C could be another
426 factor limiting the liposome efficiency. As shown in the methionine enrichment,
427 multilamellar liposomes could be a good candidate to deliver hydrosoluble compounds
428 to nauplii. Enhanced protection given by their multiple bilayers could account for a high
429 ability of water-soluble retention inside the liposomes. In these circumstances, nauplii
430 would incorporate particulate liposomes more efficiently than dissolved molecules by
431 means of their adapted filtering-natatory appendages.

432 **Acknowledgements**

433 This study was funded by the Ministry of Education and Culture of the Spanish
434 Government (project AGL2000-0135-P4-03). The authors would like to thank
435 Transtechnics, S.L. for liposome supply, and Mazdak Bagherie-Lachidan for his
436 corrections in the English manuscript.

437 **References**

- 438 Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., Watson, J.D., 1996. *Biología*
439 *Molecular de la Célula* (3rd edition). Ediciones Omega, S.A., Barcelona, 1387
440 pp.
- 441 Aust, O., Sies, H., Stahl, W., Polidori, M.C., 2001. Analysis of lipophilic antioxidants in
442 human serum and tissues: tocopherols and carotenoids. *J. Chromatogr.* 936A,
443 83-93.
- 444 Bangham, A.D., Standish, M.M., Watkins, J.C., 1965. Diffusion of univalent ions
445 across the lamellae of swollen phospholipids. *J. Mol. Biol.* 13, 238-252.

- 446 Cuesta, A., Esteban, M.A., Meseguer, J., 2002. Natural cytotoxic activity in seabream
447 (*Sparus aurata* L.) and its modulation by vitamin C. *Fish Shellfish Immunol.* 13,
448 97-109.
- 449 Dabrowski, K., 1990. Gulonolactone oxidase is missing in teleost fish – a direct
450 spectrophotometric assay. *Biol. Chem. Hoppe-Seyler* 371, 207-214.
- 451 Dedi, J., Takeuchi, T., Seikai, T., Watanabe, T., 1995. Hypervitaminosis and safe levels
452 of vitamin A for larval flounder (*Paralichthys olivaceus*) fed *Artemia* nauplii.
453 *Aquaculture* 133, 135-146.
- 454 Dedi, J., Takeuchi, T., Seikai, T., Watanabe, T., Hosoya, K., 1997. Hypervitaminosis A
455 during vertebral morphogenesis in larval Japanese flounder. *Fish. Sci.* 63, 466-
456 473.
- 457 Estévez, A., Kanazawa, A., 1995. Effect of (n-3) PUFA and vitamin A *Artemia*
458 enrichment on pigmentation success of turbot, *Scophthalmus maximus* (L).
459 *Aquacult. Nutr.* 1, 159-168.
- 460 Folch, J., Lees, N., Sloane-Stanley, G.H., 1957. A simple method for the isolation and
461 purification of total lipids from animal tissues. *J. Biol. Chem.* 226, 497-509.
- 462 Fukuzawa, K., Ikebata, W., Shibata, A., Sakanaka, T., Urano, S., 1993a. Location of α -
463 tocopherol in phospholipid vesicles and its dynamics in inhibiting lipid
464 peroxidation. In: Mino, M., Nakamura, H., Diplock, A.T., Kayden, H.J. *Vitamin*
465 *E - Its Usefulness in Health and in Curing Diseases.* Japan Scientific Society
466 Press, Tokyo (Japan). pp, 40 pp.
- 467 Fukuzawa, K., Ikebata, W., Sohmi, K., 1993b. Location, antioxidant and recycling
468 dynamics of α -tocopherol in liposome membranes. *J.Nutr.Sci.Vitaminol.* 39, S9-
469 S22.
- 470 Furr, H.C., McGrane, M.M., 2003. Retinol: physiology. In: Caballero, B., Trujo, L.,
471 Finglas, P. *Encyclopaedia of Food Sciences and Nutrition.* Academic Press,
472 4967 pp.

- 473 Fyhn, H.J., Finn, R.N., Helland, S., Rønnestad, I., Lømsland, E., 1993. Nutritional value
474 of phyto- and zooplankton as live food for marine fish larvae. In: Reinertsen, H.,
475 Dahle, L.A., Jørgensen, L., Tvinnerheim, K. Fish Farming Technology.
476 Balkema, Rotterdam, 126 pp.
- 477 Gouillou-Coustans, M.-F., Guillaume, J., 2001. Vitamin nutrition. In: Guillaume, J.,
478 Kaushik, S., Bergot, P., Métailler, R. (Eds.), Nutrition and Feeding of Fish and
479 Crustacean. Praxis Publishing Ltd., Chichester, pp. 145-167.
- 480 Haga, Y., Suzuki, T., Kagechika, H., Takeuchi, T., 2003. A retinoic acid receptor-
481 selective agonist causes jaw deformity in the Japanese flounder, *Paralichthys*
482 *olivaceus*. Aquaculture 221, 381-392.
- 483 Haga, Y., Takeuchi, T., Seikai, T., 2004. Changes of retinoid contents in larval Japanese
484 flounder *Paralichthys olivaceus* and *Artemia* nauplii enriched with a large dose
485 of all-trans retinoic acid. Fish. Sci. 70, 436-444.
- 486 Halver, J.E., 2002. The vitamins. In: Halver, J.E., Hardy, R.W. Fish Nutrition.
487 Academic Press, San Diego, 141 pp.
- 488 Hara, T.J., 2006. Feeding behaviour in some teleosts is triggered by single amino acids
489 primarily through olfaction. J. Fish Biol. 68, 810-825.
- 490 Helland, S., Triantaphyllidis, G.V., Fyhn, H.J., Evjen, M.S., Lavens, P., Sorgeloos, P.,
491 2000. Modulation of the free amino acid pool and protein content in populations
492 of the brine shrimp *Artemia* spp. Mar. Biol. 137, 1005-1016.
- 493 Henrique, M.M., Gouillou-Coustans, M.F., Gomes, E., 2002. Effect of dietary ascorbic
494 acid supplementation and chronic hypoxia on sea bream growth and vitamin C
495 status. J. Fish Biol. 60, 442-452.
- 496 Hontoria, F., Crowe, J.H., Crowe, L.M., Amat, F., 1994. Potential use of liposomes in
497 larviculture as a delivery system through *Artemia* nauplii. Aquaculture 127, 255-
498 264.
- 499 Hontoria, F., Navarro, J.C., Crowe, J.H., Crowe, L.M., Amat, F., 1997. Encapsulación
500 de ácidos grasos poliinsaturados esenciales y vitaminas en liposomas para su uso

- 501 como enriquecedores de nauplios de *Artemia*. Actas del VI Congreso Nacional
502 de Acuicultura, Cartagena (Spain), p. 127.
- 503 Hunter, J., 1981. Feeding ecology and predation. In: Lasker, R. (Ed.). Marine Fish
504 Larvae, Morphology, Ecology, and Relation to Fisheries. Wash. Sea Grant
505 Program, Univ. Wash. Press, Seattle, pp. 33-77.
- 506 Hwang, D.F., Lin, T.-K., 2002. Effect of temperature on dietary vitamin C requirement
507 and lipid in common carp. *Comp. Biochem. Physiol.* 131B, 1-7.
- 508 Izquierdo, M.S., 1996. Essential fatty acid requirements of cultured marine fish larvae.
509 *Aquacult. Nutr.* 2, 183-191.
- 510 Kolkovski, S., Koven, W., Tandler, A., 1997. The mode of action of *Artemia* in
511 enhancing utilization of microdiet by gilthead seabream *Sparus aurata* larvae.
512 *Aquaculture* 155, 193-205.
- 513 Koop, D.R., Morgan, E.T., Tarr, G.E., Coon, M.J., 1982. Purification and
514 Characterization of A Unique Isoenzyme of Cytochrome-P-450 from Liver-
515 Microsomes of Ethanol-Treated Rabbits. *J. Biol. Chem.* 257, 8472-8480.
- 516 Merchie, G., Lavens, P., Dhert, P., Dehasque, M., Nelis, H.J., de Leenheer, A.P.,
517 Sorgeloos, P., 1995a. Variation of ascorbic acid content in different live food
518 organisms. *Aquaculture* 134, 325-337.
- 519 Merchie, G., Lavens, P., Dhert, P., Pector, R., Mai Soni, A.F., Nelis, H.J., Ollevier, F.,
520 de Leenheer, A.P., Sorgeloos, P., 1995b. Live food mediated vitamin C transfer
521 to *Dicentrarchus labrax* and *Clarias gariiepinus*. *J. Appl. Ichthyol.* 11, 336-341.
- 522 Merchie, G., Lavens, P., Sorgeloos, P., 1997. Optimization of dietary vitamin C in fish
523 and crustacean larvae: a review. *Aquaculture* 155, 165-181.
- 524 Monroig, Ó., Navarro, J.C., Amat, I., González, P., Amat, F., Hontoria, F., 2003.
525 Enrichment of *Artemia* nauplii in PUFA, phospholipids, and water-soluble
526 nutrients using liposomes. *Aquacult. Int.* 11, 151-161.

- 527 Monroig, Ó., Navarro, J.C., Amat, F., González, P., Bermejo, A., Hontoria, F., 2006.
528 Enrichment of *Artemia* in essential fatty acids with different types of liposomes
529 and their use in the rearing of gilthead sea bream (*Sparus aurata*) larvae.
530 Aquaculture 251, 491-508.
- 531 Navarro, J.C., Ireland, J., Tytler, P., 1993. Effect of temperature on permeability and
532 drinking rates of the metanauplii of the brine shrimp *Artemia* sp. Mar. Biol. 116,
533 247-250.
- 534 New, R.R.C., 1990. Liposomes. A Practical Approach. IRL Press, Oxford, 301 pp.
- 535 Ozkizilcik, S., Chu, F.-L.-E., 1994. Uptake and metabolism of liposomes by *Artemia*
536 nauplii. Aquaculture 128, 131-141.
- 537 Rønnestad, I., 1992. Utilization of free amino acids in marine fish eggs and larvae. PhD.
538 Thesis. University of Bergen, Bergen, Norway.
- 539 Rønnestad, I., Thorsen, A., Finn, R.N., 1999. Fish larval nutrition: a review of recent
540 advances in the roles of amino acids. Aquaculture 177, 201-216.
- 541 Saele, Ø., Solbakken, J.S., Watanabe, K., Hamre, K., Pittman, K., 2003. The effect of
542 diet on ossification and eye migration in Atlantic halibut larvae (*Hippoglossus*
543 *hippoglossus* L.). Aquaculture 220, 683-696.
- 544 Sargent, J.R., McEvoy, L.A., Bell, J.G., 1997. Requirements, presentation and sources
545 of polyunsaturated fatty acids in marine fish larval feeds. Aquaculture 155, 117-
546 127.
- 547 Schüep, W., Marmet, J., Studer, W., 1989. Stability of ascorbate-2-sulfate in trout feed
548 measured by HPLC. Aquaculture 79, 249-258.
- 549 Seikai, T., Matsumoto, J., Shimozaki, M., Oikawa, A., Akiyama, T., 1987. An association
550 of melanophores appearing at metamorphosis as vehicles of asymmetric skin
551 color formation with pigment anomalies developed under hatchery conditions in
552 the Japanese flounder, *Paralichthys olivaceus*. Pigment Cell Res. 1, 143-151.

- 553 Shiau S.-Y., Hsu T.-S., 1994. Vitamin C requirement of grass shrimp, *Penaeus*
554 *monodon*, as determined with L-ascorbyl-2monophosphate. *Aquaculture* 122,
555 347-357.
- 556 Sokal, R.R., Rohlf, F.J., 1981. *Biometry*. Freeman, New York, 859 pp.
- 557 Takeuchi, T., Dedi, J., Haga, Y., Watanabe, T., 1998. Effect of vitamin A compounds
558 on bone deformity in larval Japanese flounder (*Paralichthys olivaceus*).
559 *Aquaculture* 169, 155-165.
- 560 Tonheim, S.K., Koven, W., Rønnestad, I., 2000. Enrichment of *Artemia* with free
561 methionine. *Aquaculture* 190, 223-235.
- 562 Touraki, M., Rigas, P., Kastritsis, C., 1995. Liposome mediated delivery of water
563 soluble antibiotics to the larvae of aquatic animals. *Aquaculture* 136, 1-10.
- 564 Urano, S., Iida, M., Otani, I., Matsuo, M., 1987. Membrane stabilization of vitamin E;
565 interactions of α -tocopherol with phospholipids in bilayer liposomes. *Biochem.*
566 *Biophys. Res. Comm.* 146, 1413-1418.
- 567 Woollard, D.C., Indyk, H.E., 2003. Retinol: properties and determination. In: Caballero,
568 B., Trujo, L., Finglas, P. (Eds.), *Encyclopaedia of Food Sciences and Nutrition*
569 (2nd edition), Academic Press, London, pp. 4952-4957.
- 570 Yoshitomi, B., 2004. Depletion of ascorbic acid derivatives in fish feed by the
571 production process. *Fish. Sci.* 70, 1153-1156.
- 572
- 573
- 574
- 575
- 576
- 577

578 Tables

579 Table 1. Selected fatty acid (percent of total fatty acids) and lipid class composition
 580 (percent of total lipids) of phospholipid sources used in the liposome formulation.

	KPE	SPC	DPPC
<i>Fatty acid</i>			
16:0	25.5	15.1	100.0
18:0	1.0	3.0	0.0
18:1n-9	5.4	13.6	0.0
18:2n-6	1.8	60.6	0.0
18:3n-3	1.4	3.6	0.0
20:4n-6	0.7	0.0	0.0
20:5n-3	29.7	0.0	0.0
22:5n-3	0.8	0.0	0.0
22:6n-3	18.4	0.0	0.0
Saturates	28.6	18.7	100.0
Monounsaturates	13.2	15.8	0.0
Polyunsaturates	55.8	64.5	0.0
Total n-3	52.5	3.8	0.0
Total n-6	3.2	60.7	0.0
HUFA n-3	49.3	0.2	0.0
HUFA n-6	1.4	0.1	0.0
DHA/EPA	0.6	0.0	0.0
<i>Phospholipid class</i>			
PC	67.0	95.0	100.0
PE	9.0	0.0	0.0
Other	24.0	5.0	0.0

581 HUFA n-3: $\geq 20:3n-3$; HUFA n-6: $\geq 20:2n-6$; DHA/EPA: docosahexaenoic to

582 eicosapentaenoic fatty acid ratio; PC: phosphatidylcholine; PE:

583 phosphatidylethanolamine.

584

585

586

587 Table 2. Enrichment products assessed in the *Artemia* enrichment in vitamin A, vitamin
588 C and methionine.

<i>Delivered nutrient</i>	<i>Enrichment products</i>	<i>Lamellarity (liposomes)</i>	<i>Preparation method (liposomes)</i>	<i>Dosage</i>	<i>Membrane composition (w/w)</i>
<i>Vitamin A</i>	Emulsion	-	-	1	-
	LUVext A	Unilamellar	Extrusion	1	KPE:vitA (100:1)
	LUVdet A	Unilamellar	Detergent solubilization	1	KPE:vitA (100:2)
<i>Vitamin C</i>	Emulsion	-	-	1	-
	LUVdet C 1	Unilamellar	Detergent solubilization	1	SPC (100%)
	LUVdet C 2	Unilamellar	Detergent solubilization	2	SPC (100%)
	Solution C	-	-	1	-
<i>Methionine</i>	LUVext M 1	Unilamellar	Extrusion	1	SPC:CHO (4:1)
	LUVext M 2	Unilamellar	Extrusion	2	SPC:CHO (4:1)
	LUVext M 3	Unilamellar	Extrusion	1	DPPC:CHO (4:1)
	MLV M	Multilamellar	Simple lipid hydration	1	SPC:CHO (4:1)
	Solution M	-	-	1	-

589 LUVdet: large unilamellar vesicles prepared by the detergent solubilization

590 methodology; LUVext: large unilamellar vesicles prepared by extrusion through

591 polycarbonate membranes; MLV: multilamellar vesicles; KPE: krill phospholipid

592 extract; SPC: soybean phosphatidylcholine; DPPC: dipalmitoyl phosphatidylcholine;

593 CHO: cholesterol; vitA: retinyl palmitate.

594

595 Table 3. Gradient elution program followed in the analysis of free amino acids.

<u>Time (min)</u>	<u>% phase A</u>	<u>% phase B</u>
0.0	100.0	0.0
13.5	97.0	3.0
16.5	97.0	3.0
19.0	96.5	3.5
21.0	95.5	4.5
24.0	94.0	6.0
25.0	93.0	7.0
26.5	92.0	8.0
28.5	91.0	9.0
30.0	91.0	9.0
50.0	66.0	34.0
65.0	30.0	70.0

596 Mobile phase A, aqueous solution of 70 mM sodium acetate (pH 6.55) containing 2.5%
597 (v/v) acetonitrile; Mobile phase B, solution of acetonitrile:distilled water:methanol
598 (45:40:15, v/v/v).

599

600


601 Figure legends

602 Figure 1. Contents of retinol (μg per gram of dry weight) of *Artemia* nauplii enriched
603 with different products. Samples of 24 h belong to nauplii kept during 3 h in clean
604 seawater after 21 h of incubation with the enrichment product. Data are means of three
605 replicates. Error bars represent standard deviations. Treatments sharing the same
606 superscript letter in a sampling point are not significantly different from each other
607 ($P \leq 0.05$).

608 Figure 2. Contents of ascorbate (mg per gram of dry weight) of *Artemia* nauplii
609 enriched with different products. Samples of 24 h belong to nauplii kept during 3 h in
610 clean seawater after 21 h of incubation with the enrichment product. Data are means of
611 three replicates. Error bars represent standard deviations. Treatments sharing the same
612 superscript letter in a sampling point are not significantly different from each other
613 ($P \leq 0.05$).

614 Figure 3. Contents of free methionine (μg per gram of dry weight) of *Artemia* nauplii
615 enriched with different products. Samples of 24 h belong to nauplii kept during 3 h in
616 clean seawater after 21 h of incubation with the enrichment product. Data are means of
617 three replicates. Error bars represent standard deviations. Treatments sharing the same
618 superscript letter in a sampling point are not significantly different from each other
619 ($P \leq 0.05$).

620






Figure 3

