

Stroke Ready Intervention: Community Engagement to Decrease Prehospital Delay

Lesli E. Skolarus, MD, MS; Marc A. Zimmerman, PhD; Sarah Bailey, MA; Mackenzie Dome, MA; Jillian B. Murphy, MPH; Christina Kobrossi, MSW, MPH; Stephan U. Dombrowski, PhD; James F. Burke, MD, MS; Lewis B. Morgenstern, MD

Background—Time-limited acute stroke treatments are underused, primarily due to prehospital delay. One approach to decreasing prehospital delay is to increase stroke preparedness, the ability to recognize stroke, and the intention to immediately call emergency medical services, through community engagement with high-risk communities.

Methods and Results—Our community–academic partnership developed and tested “Stroke Ready,” a peer-led, workshop-based, health behavior intervention to increase stroke preparedness among African American youth and adults in Flint, Michigan. Outcomes were measured with a series of 9 stroke and nonstroke 1-minute video vignettes; after each video, participants selected their intended response (primary outcome) and symptom recognition (secondary outcome), receiving 1 point for each appropriate stroke response and recognition. We assessed differences between baseline and posttest appropriate stroke response, which was defined as intent to call 911 for stroke vignettes and not calling 911 for nonstroke, nonemergent vignettes and recognition of stroke. Outcomes assessments were performed before workshop 1 (baseline), at the conclusion of workshop 2 (immediate post-test), and 1 month later (delayed post-test). A total of 101 participants completed the baseline assessment (73 adults and 28 youths), 64 completed the immediate post-test, and 68 the delayed post-test. All participants were African American. The median age of adults was 56 (interquartile range 35–65) and of youth was 14 (interquartile range 11–16), 65% of adults were women, and 50% of youths were women. Compared to baseline, appropriate stroke response was improved in the immediate post-test (4.4 versus 5.2, $P<0.01$) and was sustained in the delayed post-test (4.4 versus 5.2, $P<0.01$). Stroke recognition did not change in the immediate post-test (5.9 versus 6.0, $P=0.34$), but increased in the delayed post-test (5.9 versus 6.2, $P=0.04$).

Conclusions—Stroke Ready increased stroke preparedness, a necessary step toward increasing acute stroke treatment rates.

Clinical Trial Registration—URL: <https://www.clinicaltrials.gov/>. Unique identifier: NCT01499173. (*J Am Heart Assoc.* 2016;5:e003331 doi: 10.1161/JAHA.116.003331)

Key Words: community • outcomes research • stroke

Acute stroke treatments reduce post-stroke disability and are cost effective.^{1–3} Yet, they are dramatically underutilized—administered to less than 5% of stroke patients in the United States.⁴ The primary reason that patients do not receive time-dependent, acute stroke treatments is their delay

in hospital presentation.^{5,6} Stroke preparedness, the ability to recognize stroke and the intention to respond immediately by calling emergency medical services (EMS), is a crucial step to increasing the number of stroke patients who are eligible for acute stroke treatments.⁷ With the recent success of acute endovascular stroke treatment,³ stroke preparedness interventions are needed now more than ever.

The community is vital to any stroke preparedness intervention strategy. For example, stroke patients rarely activate EMS themselves⁸; aphasia, dysarthria, and hemiparesis make getting to the phone and communicating difficult. Furthermore, witnesses may be better able to recognize stroke symptoms in someone else.⁹ Thus, bystanders are most likely to call for EMS.⁸ Community Based Participatory Research (CBPR), a form of community engagement, is a collaborative approach to research where community and academic partners share responsibility in conceiving, designing, testing, and disseminating interventions to improve the

From the Stroke Program, University of Michigan Medical School, Ann Arbor, MI (L.E.S., M.D., J.B.M., C.K., J.F.B., L.B.M.); Department of Health Behavior of Health Education, University of Michigan School of Public Health, Ann Arbor, MI (M.A.Z.); Bridges Into the Future, Flint, MI (S.B.); Division of Psychology, School of Natural Sciences, University of Stirling, UK (S.U.D.).

Correspondence to: Lesli E. Skolarus, MD, MS, 1500 East Medical Center Dr SPC#5855, Ann Arbor, MI 48109. E-mail: lerusche@umich.edu

Received February 19, 2016; accepted April 20, 2016.

© 2016 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

health of the community.¹⁰ Novel approaches such as this may be needed to reach the ultimate goal of increased acute stroke treatments.

Stroke preparedness interventions may prove particularly valuable for African Americans, who experience greater stroke incidence, prehospital delay, and are less likely to receive acute stroke treatments than non-Hispanic whites.^{4,11–14} Overall stroke preparedness interventions have been shown to decrease prehospital delay and increase acute stroke treatment rates.¹⁵ Yet, many of these studies have not included African Americans¹⁶ or have had less success in African Americans compared to non-Hispanic whites.¹⁷ Thus, for the development and pilot testing of stroke preparedness interventions, engaging African American communities is important. Within the context of a CBPR approach, we present the development and results of pilot testing of a theory-based, peer-led intervention to increase stroke preparedness among African American youth and adults in an urban, underserved, predominately African American community.

Methods

Setting and Partnership

Flint, Michigan, the birthplace of General Motors, was once a thriving industrial city. Today, the majority of the population is African American (60%), and over 40% of the population live below the poverty level. Genesee County, where Flint is the largest city, has one of the highest age-adjusted stroke hospitalization rates in Michigan.¹⁸ Furthermore, Flint has the lowest acute stroke treatment rate of any community of its size in the United States,⁴ suggesting a substantial community need for increasing stroke preparedness.

Our community–academic partnership was established in 2009 to improve the cardiovascular health of Flint. Our partnership is composed of academic partners from the University of Michigan, including stroke neurologists and experts in health behavior and health education, and community partners from Bridges into the Future, a faith-based organization dedicated to improving the health of Flint. By means of a CBPR approach, we developed and tested a theory-based, peer-led intervention, “Stroke Ready,” between September 2010 and August 2015. Our purpose was to increase stroke preparedness among adults and youth in 3 African American churches.

Stroke Ready Health Behavior Theory

While educational levels are associated with acute stroke treatment rates, previous stroke and stroke knowledge are not associated with stroke preparedness, suggesting that

knowledge is likely necessary but not sufficient to decrease prehospital delay.^{19,20} Thus, interventions based on behavioral theory are needed. Constructs from the Theory of Planned Behavior were used to guide Stroke Ready development and assessments.²¹ According to the Theory of Planned Behavior, the most immediate predictor of behavior is intention, which in turn is best predicted by an individual’s attitude toward the behavior, perceived subjective norms around the behavior, and perceived self-efficacy with regard to the behavior.²² For example, individuals with greater intent to call EMS for stroke will be more likely to actually do so when an acute stroke situation arises.²³ An individual’s intention can be enhanced by increasing self-efficacy (one’s perceived confidence in her ability to recognize and react to stroke symptoms), improving attitudes (one’s negative or positive feelings toward the behavior), and reshaping subjective norms (one’s perception about how others will judge the behavior). Stroke Ready materials and workshops focused on augmenting self-efficacy, attitude, and subjective norms around calling 911 as well as providing education in stroke recognition and response.

Stroke Ready Development and Testing

Phase 1: development of culturally sensitive Stroke Ready materials

Stroke Ready materials consisted of 5 main components: (1) Participant workbook, (2) Peer leader manual, (3) Workshop PowerPoint slides, (4) Stroke video vignettes, and (5) Stroke Ready music video. The 41-page Stroke Ready participant workbook was conceptualized and drafted by the community–academic partners, reviewed by the community advisory boards, and designed by a local Flint graphic designer. It emphasized community members’ description of stroke symptoms and community-relevant stroke statistics (Figure 1). The 97-page peer leader manual was constructed with 1 page for the PowerPoint slide and the opposing page with peer leader instructions for the delivery of the slide. The Stroke Ready PowerPoint presentation topics are outlined in Table 1. Stroke video vignettes were embedded into the PowerPoint to show stroke symptoms. Previous interventions have had success incorporating music, particularly hip-hop, into stroke education.²⁴ While community partners were enthusiastic about including music, they felt that hip-hop would not be appropriate for older adult church members and instead suggested Gospel music. Thus, we collaborated with a Flint church music director to create an original score and lyrics incorporating the National Stroke Association’s FAST stroke symptoms mnemonic (eg, F-facial droop, A-arm weakness, S-slurred speech, T-time to call 911). This original Gospel music score and video (<https://www.youtube.com/watch?v=VIPZ4h-e2nQ>) was performed by local youth and adults




Figure 1. Stroke Ready workbook.

and featured dancing by the academic and community partners. The music video included a strong focus on self-efficacy by asking viewers to participate in demonstrating stroke signs. All materials were reviewed by community

focus groups²⁵ and our adult and youth community advisory board to ensure cultural sensitivity.

Phase 2: peer leader training

To increase community capacity and promote sustainability of the intervention,¹⁰ Stroke Ready was delivered by peer leaders. We recruited and trained youth (ages 10–18) and adult peer-leaders from three Flint churches. Peer leaders were selected based on their previous participation or leadership roles within the church. We exceeded our goal of 12 peer leaders by recruiting and training 27 peer leaders, including 11 youth.

Each peer leader attended a 3-hour training focused on 3 core areas: stroke preparedness, skills-based knowledge (communication, technology, leading activities), and research implementation (basic understanding of program planning including monitoring and evaluation).²⁶ We utilized a teach-back model for training peer leaders, whereby workshops were first taught by stroke neurologists and community partners and then later presented back to the research team by the peer leaders.²⁷ Peer leaders received t-shirts to wear during the workshops, monetary compensation for their time, and certificates of training completion.

Phase 3: intervention recruitment and delivery

Community partners and peer-leaders recruited workshop participants through word-of-mouth, church bulletins, flyers, text messaging, and church service announcements; each church recruited 20 adults and 20 youths ages 10 and above. The workshops were delivered 1 week apart and were led by at least 2 peer leaders using the peer leader manual and

Table 1. Overview of the Stroke Ready Intervention

Method of Peer Leader Delivery	Workshop 1	Workshop 2
Introduction	Recognize that stroke is an emergency and treatable	Recognize that stroke is an emergency and treatable
Read	Stroke is common in Flint For tPA, the faster you call 911 the better Review: F.A.S.T.	Review: F.A.S.T. Overcoming barriers to calling 911 Coping with stress Review: calling 911
Audio	What is a stroke Stroke is common among African Americans African Americans have greater post-stroke disability Stroke is treatable: tPA is a clot-buster medicine Call 911 to get help	What to expect when you call 911 Waiting for help to arrive Navigating the Emergency Department Stroke risk factors Stroke prevention: hypertension
Interactive activities	Think F.A.S.T. tPA Activity Workshop review: discussion	What to expect when you call 911: Discussion Stroke role play Workshop review: discussion
Video media	Stroke Clips: F.A.S.T. Signs of Stroke Music Video	Signs of Stroke Music Video

tPA indicates tissue plasminogen activator.

PowerPoint presentations. The multimedia workshops included PowerPoint slides that were read aloud, videos showing stroke symptoms, facilitated discussions, and audio recording for more medically based information (Table 1).

The first workshop included sections on (1) stroke recognition and the importance of calling 911; (2) stroke burden in the Flint African American community; (3) improving outcome expectations (ie, stroke is treatable); and (4) an interactive demonstration using straws for arteries with play dough for clots and pipe cleaners for acute stroke treatment. This demonstration showed the value of the acute stroke treatment and encouraged positive attitudes toward securing this outcome.²⁸ The second session focused on subjective norms and self-efficacy to call 911 through stroke role-play scenarios. Scripts were provided for characters including stroke victim, family member, and 911 operators to allow participants to practice the desired behavior (ie, self-efficacy). Workshop 2 also contained a discussion of barriers to calling EMS and how to overcome these barriers that were identified during community focus groups (ie, subjective norms).²⁵ Participants received certificates, bags, pens, and a small monetary compensation for their involvement.

Outcome Assessment

We created an outcome measure to assess stroke recognition and response using video vignettes. While previous stroke preparedness assessments included written vignettes, health video vignettes were created to provide a more realistic experience for the viewer.²⁹ The stroke preparedness outcome measure included 9, 45-second video vignettes (5 acute stroke, 2 nonacute stroke (strokes that occurred weeks or months ago), and 2 nonstroke (peripheral neuropathy, myocardial infarction); participants selected the diagnosis

and their intended response after each video. Each video had a unique set of answer choices that were determined from cognitive interviews with community members. Stroke recognition answer choices included 4 unique diagnoses and “don’t know”; stroke response options included 5 unique choices and “don’t know.” For acute stroke scenarios, 1 point was awarded for the appropriate responses of recognizing stroke and calling 911. For nonacute stroke and nonstroke scenarios, 1 point was awarded for each appropriate response, which was defined as any response other than stroke and calling 911. By including nonacute stroke and nonstroke scenarios, the instrument selected against participants calling 911 for all scenarios. Stroke recognition (score 0–9) and response (score 0–8) were scored independently. One of the vignettes was an acute myocardial infarction, which did not fit the response scoring, leaving 8 response vignettes. Assessments were completed before the first workshop (baseline), at the conclusion of workshop 2 (immediate post-test), and 1 month later (delayed post-test). Our primary outcome was the change in appropriate response to call 911 between the baseline and the immediate post-test and the secondary outcome was the change in appropriate stroke recognition between the baseline and the immediate post-test. We also measured stroke attitude, self-efficacy, and subjective norms using a 5-point Likert scale (Table 2). Process measures assessed Stroke Ready quality, comprehension, and satisfaction using a Likert scale.

Analysis

Sociodemographics were described using descriptive statistics. Multilevel linear regression models with a fixed church-level intercept and a random participant-level intercept to account for the times series analysis were used to explore the

Table 2. Theory of Planned Behavior Psychological Constructs

Psychological Construct	Immediate Post-Test Odds Ratio (95% CI)	P Value	Delayed Post-Test Odds Ratio (95% CI)	P Value
Stroke self-efficacy*				
I would be able to tell if someone is having a stroke	1.2 (0.5, 2.0)	<0.01	0.8 (0.07, 1.5)	0.03
I know what to do if I saw someone having a stroke	0.8 (0.03, 1.6)	0.04	0.3 (–0.5, 1.0)	0.51
Stroke attitude				
If I were to see signs of a stroke, calling 911 would be [†]	0.05 (–0.8, 0.9)	0.92	–0.07 (–0.9, 0.8)	0.87
If a person has signs of a stroke, calling 911 right away could be [‡]	0.3 (–0.8, 1.5)	0.57	–0.5 (–1.6, 0.7)	0.43
Subjective norms*				
Most people would call 911 if they were to see a stroke	–0.5 (–1.1, 0.09)	0.09	–0.9 (–1.5, –0.2)	<0.01
My family would want me to call 911 if I were to see a stroke	0.2 (–0.7, 1.0)	0.72	0.1 (–0.8, 1.1)	0.75

*Response stems range from “strongly agree” to “strongly disagree.”

[†]Response stems range from “extremely pleasant” to “very unpleasant.”

[‡]Response stems range from “very helpful” to “very harmful.”

Table 3. Sociodemographics and Medical Conditions of Stroke Ready Participants (N=10)

	Adult No. Responding (n=73)	Adult % of Respondents	Youth No. Responding (n=28)	Youth % of Respondents
Age y, median (IQR)	65	56 (35–65)	28	14 (11–16)
Women	68	65	28	50
Married	68	37		
Live alone	67	22		
Education	67			
High School graduate or less		38		
Some college		29		
College graduate		15		
Advanced degree		9		
Medical condition				
Stroke (including TIA)	55	4		
High blood pressure	61	49		
Heart attack	51	6		
Diabetes mellitus	56	34		
Atrial fibrillation	48	2		
Insurance status	64			
Private		39		
Medicare		30		
Medicaid		20		
Genesee County Health plan		3		
Uninsured		8		

IQR indicates interquartile range; TIA, transient ischemic attack.

change in stroke response and recognition. Average marginal effects were used to report the change in appropriate stroke response and recognition after completing the immediate post-test and delayed post-test. We determined the intraclass coefficient to estimate the impact of clustering at the church level on intervention efficacy. The majority of the missing data were for 1 video and rather than drop these participants from the primary analysis, we used the conservative approach of including them in the analysis with missing data recoded as “don’t know.” We explored differences in the effect of the intervention based on whether the participant was a youth or an adult by including interaction terms of outcome assessment and age group (youth versus adult) into the multilevel linear regression models assessing stroke response and recognition. Multilevel mixed-effects ordered logistic regression models with a fixed church-level intercept and a random participant-level intercept were used to explore change between baseline and immediate post-test and baseline and delayed post-test in stroke attitude, self-efficacy, and subjective norms after accounting for the participants’ church.

We conducted a sensitivity analysis to evaluate the robustness of the findings based on 3 different approaches to missing

data. For participants with missing data from 1 video, data were recoded as the correct answer and in a second analysis recoded as the incorrect answer. Additionally, we excluded participants with any missing data. All analyses were performed using STATA 11 and STATA 14. The research was approved by the University of Michigan Institutional Review Board and all participants signed written informed consent. The trial was registered on clinicaltrials.gov NCT01499173.

Results

A total of 104 participants attended workshop 1 and 77 attended workshop 2. A total of 101 participants (73 adults and 28 youth) completed the baseline assessment, 64 completed the immediate post-test (47 adults and 17 youth), and 68 completed the delayed post-test (53 adults and 15 youth). All participants were African American (Table 3). The median age of the adult participants was 56 (interquartile range 35–65), 65% were women and 38% had a high school education or less. Forty-nine percent of adult respondents had hypertension and 4% have had a stroke. The median age of the youth participants was 14 (interquartile range 11–16) and

50% were women. Of the 9 participants in the 16- to 20-year-old age group, 8 were men. We found no difference in age, sex, pretest stroke recognition, or appropriate response between those who completed the immediate and delayed post-test and those lost to follow-up.

Stroke Preparedness

Compared to baseline scores, the appropriate stroke response increased in the immediate post-test (4.4 versus 5.2, $P<0.01$) and delayed post-test (4.4 versus 5.2, $P<0.01$) (Figure 2). We found no significant increase in stroke recognition when comparing the baseline and immediate post-test (5.9 versus 6.0, $P=0.34$), but stroke recognition increased in the delayed post-test (5.9 versus 6.2, $P=0.04$). No interaction effects between age group and stroke response ($P=0.72$) or stroke recognition ($P=0.23$) were found. The intraclass correlation coefficient in an empty model was 0.062, suggesting that church explains about 6% of all variance in stroke action. In the sensitivity analysis, we found no change in the study results under different scenarios of missing data.

Stroke Attitude, Self-Efficacy, and Subjective Norms

We found no difference between baseline and immediate post-test or delayed post-test measures of stroke attitude or family norms (Table 2). Compared to the baseline, self-efficacy for stroke recognition increased in the immediate post-test (odds ratio [OR]=1.2, 95% CI: 0.5, 2.0, $P<0.01$), which was sustained in the delayed post-test (OR=0.8, 95% CI: 0.07, 1.5, $P=0.03$). When comparing the baseline to the immediate post-test, self-efficacy for stroke response increased (OR=0.8, 95% CI: 0.03, 1.6, $P=0.04$) and became nonsignificant in the delayed post-test (OR=0.3, 95% CI: -0.5,

1.0, $P=0.51$). No change in the community subjective norms between baseline and immediate post-test (OR=-0.5, 95% CI: -1.1, 0.09, $P=0.09$) was found, but community subjective norms decreased in the delayed post-test (OR=-0.9, 95% CI: -1.5, -0.2, $P<0.01$).

Acceptability Measures

A total of 95% (95% CI: 87–99) percent of participants felt Stroke Ready was good or excellent quality and 84% (95% CI: 74–92) reported they understood most or all of the workshop. A total of 80% (95% CI: 70–89) of participants were very or extremely satisfied with Stroke Ready and 97% (95% CI: 91–99) would recommend it to a friend.

Discussion

Stroke Ready is the first stroke preparedness intervention developed and tested through a CBPR approach in collaboration with African American churches. Stroke Ready increased appropriate stroke response immediately after completion of the intervention, and this increased response was sustained after 1 month. Stroke recognition increased 1 month after the intervention. The improvement in stroke self-efficacy, which formed the basis of the behavioral theory for the Stroke Ready intervention, suggests a possible pathway by which appropriate stroke response was increased. Moreover, participants reported high levels of quality, understanding, and satisfaction with Stroke Ready. With the promising results of Stroke Ready on the intermediate outcomes of stroke response and recognition, further studies are needed to test clinical outcomes such as decreased prehospital delay and increased acute stroke treatments.

To measure improvement in acute treatment rates, large-scale interventions such as the acute myocardial infarction study, REACT, or the TLL Temple Foundation Stroke Project are needed.^{30,31} During both trials communities that received a multilevel hospital and community intervention were compared to usual care. No differences in prehospital delay were found in REACT, while acute stroke treatment rates increased in the TLL Temple Foundation Stroke Project. Single-level interventions have also shown an increase in acute stroke treatments. Community stroke preparedness interventions such as mailings and mass media have decreased prehospital delay and increased acute stroke treatments.^{32,33} Similarly randomized, hospital implementation trials of acute stroke practices have resulted in small increases in the number of stroke patients receiving acute stroke treatments.^{34,35} Because both community and hospital-level interventions have shown an increase in acute stroke treatment rates, determining the optimal use of resources to increase acute stroke treatments will likely be community




Figure 2. Stroke response scores among Stroke Ready participants. $P<0.01$ for baseline vs. immediate post-test; $P<0.01$ for baseline vs. delayed post-test.

specific. Stroke Ready is designed for communities where the largest barrier to acute stroke treatment is prehospital delay and is specifically intended for communities with large African American populations. The Stroke Ready's religious content was minimal and found to be acceptable by participants of various religions and levels of religiosity. Thus, we think that Stroke Ready could be widely disseminated in relevant communities.

Stroke Ready highlights the benefits of community engagement. We attribute the excellent recruitment of peer leaders and adult participants to the CBPR approach. The churches were well aware that Stroke Ready was created for them by members of their church and community. We received strong support from church leadership, in part because stroke education is a recognized need in the community, and also because of the long-standing relationship between community partners and Flint churches. Stroke Ready enrolled a substantial number of young African American men, a population historically underserved in medical research.³⁶ This success may be because peer leaders, many of whom were young men, assisted in participant recruitment. The community embraced Stroke Ready and made it their own. A church youth group leader, for example, held additional training sessions for her youth peer leaders.

Stroke Ready exceeded its enrollment goals for peer leaders, met the goals for adult participants, but fell short of the youth enrollment goals. Alternate recruitment strategies for youth such as social media could be considered in the future. The intervention evaluation was limited by attrition. Attrition was greater than in some stroke preparedness interventions where testing was performed for all participants simultaneously during school hours.^{24,37} Retention was also higher in other behavioral interventions partnering with African American churches that incorporated strategies such as church incentives for outcome assessment and multiple time options for outcome assessments.³⁸ These strategies should be considered in the future.

This study has limitations. This was a pre–post design and thus we cannot conclude that Stroke Ready caused an increase in stroke preparedness and psychological variables. However, the short duration between baseline and outcomes and the lack of any known local or national stroke preparedness campaign during the time frame suggest an impact of the intervention. Second, Stroke Ready experienced substantial attrition resulting in missing data. The demographic characteristics of the participants and those lost to follow-up, however, were similar and our primary analyses arrived at similar conclusions using different approaches to missing data. However, we cannot confirm that those lost to follow-up were truly missing at random and thus it is possible that these missing data may contribute subtle biases. Stroke Ready

consisted of 2 sessions and thus it is unknown whether fewer workshops would have resulted in improved stroke preparedness. We did not collect data on occupations and thus cannot exclude that many participants may work in the healthcare sector. Finally, our church-based educational intervention may limit the generalizability of the study beyond this context. Nevertheless, it is an efficient way to engage African Americans who are more likely to identify themselves as religious and attend church than non-Hispanic whites.³⁹

In conclusion, Stroke Ready increased stroke preparedness and stroke self-efficacy, which are necessary steps toward increasing acute stroke treatment rates. Further study is needed to determine whether Stroke Ready can decrease prehospital delay.

Acknowledgments

We would like to thank Pastor Patrick Sanders and Bishop Otis Floyd of New Jerusalem Full Gospel Baptist Church, Pastors Seon Thompson and Matthew Jackson of Refuge Temple, Pastor Phillip Thompson of Bethlehem Temple Church, and Bishop Urundi Knox of Ebenezer Ministry for their support. We would also like to thank Barry Turner, Gloria Moses-Colon, and Donald Moses from the Community Outreach for Family and Youth (COFY) center. We thank Elder Jeffrey LaValley of Candied Jams, Jamal Bransford, Alliance recording, Mike Richardson of InnerCity Productions, and Joshua Worth and Tony Trischler of T3 promotions for their creative contributions. We especially thank E. Hill DeLoney of the Community Based Organization partners (CBOP) for her guidance and support.

Sources of Funding

We received funding from NIH/NINDS K23 NS073685 (Skolarus), NIH/NINDS K08NS082597 (Burke).

Disclosures

None.

References

1. Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med*. 1995;333:1581–1587.
2. Fagan SC, Morgenstern LB, Petitta A, Ward RE, Tilley BC, Marler JR, Levine SR, Broderick JP, Kwiatkowski TG, Frankel M, Brott TG, Walker MD. Cost-effectiveness of tissue plasminogen activator for acute ischemic stroke. NINDS rt-PA Stroke Study Group. *Neurology*. 1998;50:883–890.
3. Powers WJ, Derdeyn CP, Biller J, Coffey CS, Hoh BL, Jauch EC, Johnston KC, Johnston SC, Khalessi AA, Kidwell CS. 2015 AHA/ASA focused update of the 2013 guidelines for the early management of patients with acute ischemic stroke regarding endovascular treatment: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2015;46:3020–3035.
4. Skolarus LE, Meurer WJ, Shanmugasundaram K, Adelman EE, Scott PA, Burke JF. Marked regional variation in acute stroke treatment among Medicare beneficiaries. *Stroke*. 2015;46:1890–1896.
5. Moser DK, Kimble LP, Alberts MJ, Alonzo A, Croft JB, Dracup K, Evenson KR, Go AS, Hand MM, Kothari RU, Mensah GA, Morris DL, Pancioli AM, Riegel B,

- Zerwic JJ. Reducing delay in seeking treatment by patients with acute coronary syndrome and stroke: a scientific statement from the American Heart Association Council on Cardiovascular Nursing and Stroke Council. *Circulation*. 2006;114:168–182.
6. Kleindorfer D, Kissela B, Schneider A, Woo D, Khoury J, Miller R, Alwell K, Gebel J, Szafarski J, Pancioli A, Jauch E, Moomaw C, Shukla R, Broderick JP. Eligibility for recombinant tissue plasminogen activator in acute ischemic stroke: a population-based study. *Stroke*. 2004;35:e27–e29.
 7. Schroeder EB, Rosamond WD, Morris DL, Evenson KR, Hinn AR. Determinants of use of emergency medical services in a population with stroke symptoms: the Second Delay in Accessing Stroke Healthcare (DASH II) Study. *Stroke*. 2000;31:2591–2596.
 8. Wein TH, Staub L, Felberg R, Hickenbottom SL, Chan W, Grotta JC, Demchuk AM, Groff J, Bartholomew LK, Morgenstern LB. Activation of emergency medical services for acute stroke in a nonurban population: the T.L.L. Temple Foundation Stroke Project. *Stroke*. 2000;31:1925–1928.
 9. Dombrowski SU, Sniehotta FF, Mackintosh J, White M, Rodgers H, Thomson RG, Murtagh MJ, Ford GA, Eccles MP, Araujo-Soares V. Witness response at acute onset of stroke: a qualitative theory-guided study. *PLoS One*. 2012;7:e39852.
 10. Israel BA, Schulz AJ, Parker EA, Becker AB. Review of community-based research: assessing partnership approaches to improve public health. *Annu Rev Public Health*. 1998;19:173–202.
 11. Schwamm LH, Reeves MJ, Pan W, Smith EE, Frankel MR, Olson D, Zhao X, Peterson E, Fonarow GC. Race/ethnicity, quality of care, and outcomes in ischemic stroke. *Circulation*. 2010;121:1492–1501.
 12. Johnston SC, Fung LH, Gillum LA, Smith WS, Brass LM, Lichtman JH, Brown AN, Wang DZ. Utilization of intravenous tissue-type plasminogen activator for ischemic stroke at academic medical centers. *Stroke*. 2001;32:1061–1068.
 13. Hsia AW, Edwards DF, Morgenstern LB, Wing JJ, Brown NC, Coles R, Loftin S, Wein A, Koslosky SS, Fatima S, Sanchez BN, Fokar A, Gibbons MC, Shara N, Jayam-Trouth A, Kidwell CS. Racial disparities in tissue plasminogen activator treatment rate for stroke: a population-based study. *Stroke*. 2011;42:2217–2221.
 14. Howard VJ, Kleindorfer DO, Judd SE, McClure LA, Safford MM, Rhodes JD, Cushman M, Moy CS, Soliman EZ, Kissela BM. Disparities in stroke incidence contributing to disparities in stroke mortality. *Ann Neurol*. 2011;69:619–627.
 15. Mellon L. Stroke warning campaigns: delivering better patient outcomes? A systematic review. *Patient Relat Outcome Meas*. 2015;6:61.
 16. Gardois P, Booth A, Goyder E, Ryan T. Health promotion interventions for increasing stroke awareness in ethnic minorities: a systematic review of the literature. *BMC Public Health*. 2014;14:409.
 17. Boden-Albala B, Edwards DF, Wing JJ, Gibbons MC, Hsia A, Kidwell CS. Abstract T MP71: evaluation of a community-wide stroke preparedness intervention: results of the ASPIRE study. *Stroke*. 2014;45:ATMP71.
 18. Anderson B, Lyon-Callo S, Heiler P, Miller H, Theisen V. *Impact of Heart Disease and Stroke in Michigan: 2008 Report on Surveillance*. Lansing, MI: Michigan Department of Community Health, Bureau of Epidemiology, Chronic Disease Epidemiology Section; 2015.
 19. Williams LS, Bruno A, Rouch D, Marriott DJ, Mas DJ. Stroke patients' knowledge of stroke: influence on time to presentation. *Stroke*. 1997;28:912–915.
 20. Fussman C, Rafferty AP, Lyon-Callo S, Morgenstern LB, Reeves MJ. Lack of association between stroke symptom knowledge and intent to call 911. *Stroke*. 2010;41:1501–1507.
 21. Ajzen I. Theories of cognitive self-regulation: the theory of planned behavior. *Organ Behav Hum Decis Process*. 1991;50:179–211.
 22. Ajzen I. Perceived behavioral control, self-efficacy, locus of control, and the theory of planned behavior. *J Appl Soc Psychol*. 2002;32:665–683.
 23. Skolarus LE, Zimmerman MA, Murphy J, Brown DL, Kerber KA, Bailey S, Fowlkes S, Morgenstern LB. Community-based participatory research: a new approach to engaging community members to rapidly call 911 for stroke. *Stroke*. 2011;42:1862–1866.
 24. Williams O, Noble JM. 'Hip-Hop' stroke. *Stroke*. 2008;39:2809–2816.
 25. Skolarus LE, Murphy JB, Zimmerman MA, Bailey S, Fowlkes S, Brown DL, Lisabeth LD, Greenberg E, Morgenstern LB. Individual and community determinants of calling 911 for stroke among African Americans in an urban community. *Circ Cardiovasc Qual Outcomes*. 2013;6:278–283.
 26. O'Brien MJ, Squires AP, Bixby RA, Larson SC. Role development of community health workers: an examination of selection and training processes in the intervention literature. *Am J Prev Med*. 2009;37:S262–S269.
 27. The share approach—using the teach-back technique: a reference guide for health care providers. Rockville, MD: Agency for Healthcare Research and Quality; July 2014. Available at: <http://www.ahrq.gov/professionals/education/curriculum-tools/shareddecisionmaking/tools/tool-6/index.html>. Accessed September 9, 2015.
 28. Morgenstern LB, Gonzales NR, Maddox KE, Brown DL, Karim AP, Espinosa N, Moyé LA, Pary JK, Grotta JC, Lisabeth LD, Conley KM. A randomized, controlled trial to teach middle school children to recognize stroke and call 911. *Stroke*. 2007;38:2972–2978.
 29. Chan D, Schmitt N. Video-based versus paper-and-pencil method of assessment in situational judgment tests: subgroup differences in test performance and face validity perceptions. *J Appl Psychol*. 1997;82:143–159.
 30. Luepker RV, Raczynski JM, Osganian S, Goldberg RJ, Finnegan JR, Hedges JR, Goff DC, Eisenberg MS, Zapka JG, Feldman HA, Labarthe DR, McGovern PG, Cornell CE, Proschan MA, Simons-Morton DG. Effect of a community intervention on patient delay and emergency medical service use in acute coronary heart disease. *JAMA*. 2000;284:60–67.
 31. Morgenstern LB, Bartholomew LK, Grotta JC, Staub L, King M, Chan W. Sustained benefit of a community and professional intervention to increase acute stroke therapy. *Arch Intern Med*. 2003;163:2198–2202.
 32. Flynn D, Ford GA, Rodgers H, Price C, Steen N, Thomson RG. A time series evaluation of the FAST National Stroke Awareness Campaign in England. *PLoS one*. 2014;9:e104289.
 33. Mueller M, Purnell TS, Mensah GA, Cooper LA. Reducing racial and ethnic disparities in hypertension prevention and control: what will it take to translate research into practice and policy? *Am J Hypertens*. 2015;28:699–716.
 34. Scott PA, Meurer WJ, Frederiksen SM, Kalbfleisch JD, Xu Z, Haan MN, Silbergleit R, Morgenstern LB; Investigators I. A multilevel intervention to increase community hospital use of alteplase for acute stroke (INSTINCT): a cluster-randomised controlled trial. *Lancet Neurol*. 2013;12:139–148.
 35. Dirks M, Niessen LW, van Wijngaarden JD, Koudstaal PJ, Franke CL, van Oostenbrugge RJ, Huijsman R, Lingsma HF, Minkman MM, Dippel DW. Promoting thrombolysis in acute ischemic stroke. *Stroke*. 2011;42:1325–1330.
 36. Woods VD, Montgomery SB, Herring RP. Recruiting Black/African American men for research on prostate cancer prevention. *Cancer*. 2004;100:1017–1025.
 37. Williams O, Hecht MF, DeSorbo AL, Huq S, Noble JM. Effect of a novel video game on stroke knowledge of 9-to-10-year-old, low-income children. *Stroke*. 2014;45:889–892.
 38. Resnicow K, Kramish Campbell M, Carr C, McCarty F, Wang T, Periasamy S, Rahotep S, Doyle C, Williams A, Stables G. Body and soul: a dietary intervention conducted through African-American churches. *Am J Prev Med*. 2004;27:97–105.
 39. A religious portrait of African-Americans. 2015. Available at: <http://www.pewforum.org/2009/01/30/a-religious-portrait-of-african-americans/>. Accessed July 15, 2015.



Stroke Ready Intervention: Community Engagement to Decrease Prehospital Delay

Lesli E. Skolarus, Marc A. Zimmerman, Sarah Bailey, Mackenzie Dome, Jillian B. Murphy, Christina Kobrossi, Stephan U. Dombrowski, James F. Burke and Lewis B. Morgenstern

J Am Heart Assoc. 2016;5:e003331; originally published May 20, 2016;
doi: 10.1161/JAHA.116.003331

The *Journal of the American Heart Association* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Online ISSN: 2047-9980

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jaha.ahajournals.org/content/5/5/e003331>