

Norambuena F, Morais S, Estevez A, Bell JG, Tocher DR, Navarro JC, Cerda J & Duncan N (2013) Dietary modulation of arachidonic acid metabolism in senegalese sole (*Solea Sengalensis*) broodstock reared in captivity, *Aquaculture*, 372-375, pp. 80-88.

This is the peer reviewed version of this article

NOTICE: this is the author's version of a work that was accepted for publication in Aquaculture. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Aquaculture, [VOL 372-375 (2013)] DOI: <http://dx.doi.org/10.1016/j.aquaculture.2012.10.035>

1 **DIETARY MODULATION OF ARACHIDONIC ACID METABOLISM IN**
2 **SENEGALESE SOLE (*SOLEA SENEGALENSIS*) BROODSTOCK REARED IN**
3 **CAPTIVITY**

4 Fernando Norambuena^{1*}, Sofia Morais¹, Alicia Estévez¹, J Gordon Bell², Douglas R. Tocher²,
5 Juan C. Navarro³, Joan Cerdà⁴ and Neil Duncan¹

6 ¹IRTA-Sant Carles de la Ràpita, Ctra. Poble Nou Km 6, 43540-Sant Carles de la Ràpita,
7 Tarragona, Spain

8 ²Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9 4LA,
9 Scotland, UK

10 ³Institute of Aquaculture of Torre la Sal, Spanish Council for Scientific Research (CSIC), Torre
11 la Sal s/n, 12595-Cabanes, Castellón, Spain

12 ⁴IRTA-Institute of Marine Sciences (CSIC), 08003 Barcelona, Spain

13 *Corresponding author: Fernando Norambuena, Phone: +34- 977 745427, Fax +34-977744138
14 email: norambuena52@hotmail.com

15

16 ***Abstract***

17 Previous studies have shown higher levels of arachidonic acid (20:4n-6, ARA) in testis, liver,
18 and muscle of wild Senegalese sole (*Solea senegalensis*) compared to fish reared in captivity
19 (first generation, G1). The present study was conducted to establish the optimal level of dietary
20 ARA for G1 Senegalese sole broodstock, using as a reference the fatty acid profile of wild
21 broodstock (gonads, liver and muscle). A total of 120 Senegalese sole broodstock were randomly
22 distributed into 12 tanks and (1:1 male and female), fed in duplicate with six experimental diets
23 containing increasing amounts of ARA (0.7%, 1.6%, 2.3%, 3.2%, 5.0%, and 6.0 % of total fatty
24 acids) for nine months. The relative ARA levels in liver, muscle and male and female gonads at
25 the end of the feeding period increased in a dose dependent manner. Dietary ARA was mainly
26 incorporated and stored in testis or ovary, followed by liver and muscle. Fish fed 2.3% and 3.2%
27 ARA showed no differences in the ARA content of testis, ovary and liver when compared to

28 wild fish. In male fish, a significant increase in the levels of 22:4n-6 and 22:5n-6 fatty acids was
29 also observed, which was consistent with the up-regulation of fatty acyl elongase (*elovl5*) and
30 desaturase (*d4fad*) transcript levels in the liver of fish fed 0.7%, 2.3% and 6% ARA. These
31 results suggest that dietary inclusion of 3.2% ARA during periods shorter than nine months, or of
32 2.3% ARA for prolonged periods, can maintain optimal levels of tissue ARA in captive
33 Senegalese sole broodstock. In addition, the data indicate that male Senegalese sole is able to
34 elongate and desaturate ARA to 22:4n-6 and 22:5n-6, suggesting that these fatty acids may be
35 important for male reproduction.

36 **Keywords:** Arachidonic acid, broodstock fish, fish nutrition, fatty acids.

37 **1. Introduction**

38 One of the most important nutritional factors for successful fish reproduction, and among the
39 most studied, is the fatty acid arachidonic acid (20:4n-6, ARA) [1-6]. Arachidonic acid is the
40 main precursor for production of 2-series prostaglandins (PGs) [7,8], which stimulate ovarian
41 and testicular steroidogenesis, triggering oocyte maturation in females and milt production in
42 males, and are involved in female sexual behaviour [9-13]. Arachidonic acid itself and its
43 metabolites regulate cholesterol (CHOL) transfer from the outer to inner mitochondrial
44 membrane where the P450 enzyme resides to initiate steroid hormone synthesis [11,14].
45 Moreover, ARA had differential effects on steroid biosynthesis. Although it stimulates
46 testosterone production by elevating cAMP levels in a dose-dependent manner, ARA at high
47 doses can also inhibit steroidogenesis by affecting the availability of CHOL [11,15].

48 Senegalese sole (*Solea senegalensis*) is a promising species for aquaculture in Southern Europe.
49 However, an important problem encountered in this species is the fact that first generation (G1)
50 of reared fish often fail to spawn viable eggs, contrary to wild animals, which produce eggs of
51 sufficient quality and quantity after variable times of acclimation in captivity [16]. This is
52 hindering the expansion of Sole aquaculture and hence several studies have been recently
53 performed in an attempt to understand the underlying causes. Studies on wild sole broodstock
54 showed higher levels of ARA and ARA-derived fatty acids in different tissues compared to those
55 observed in G1 fish [17], similarly to that has been reported in other fish species [18-27]. High
56 accumulation of ARA in sperm has also been reported in rainbow trout (*Oncorhynchus mykiss*)

57 fed diets low in docosahexaenoic acid (22:6n-3, DHA) [28], and in wild European seabass
58 (*Dicentrarchus labrax*) [29]. In addition, previous studies on Senegalese sole showed that
59 differences in ARA tissue content resulted in differences in cyclooxygenase (COX-2) gene
60 expression, which was significantly up-regulated in the sperm-duct, oviduct and gills of males
61 from wild origin compared to G1 fish [30]. Thus, wild fish showed significantly higher levels of
62 2-series PGs compared to cultured fish, especially in testis, whereas G1 Senegalese sole, with a
63 lower ARA tissue content, exhibited significantly higher levels of 3-series PGs and lower levels
64 of CHOL [17], the precursor of steroid hormones in vertebrates [31]. On the other hand,
65 Senegalese sole fed artificial diets formulated with graded ARA levels showed an increase in
66 ARA levels in circulating blood, which in turn may induce an increase in CHOL and steroid
67 production, especially in males [32]. Higher levels of ARA in the tissues of wild fish and
68 increased levels in the blood of G1 fish previously fed ARA-enriched diets resulted in an
69 increase in ARA-derived fatty acids, 22:4n-6 and 22:5n-6 [32]. A similar increase in these n-6
70 long-chain polyunsaturated fatty acids (LC-PUFA) was observed in the sperm of wild European
71 seabass [29]. These fatty acids are present in the cells of reproductive (i.e., seminiferous tubules,
72 sperm) and nervous tissues in larger quantities (human, bull, boar and rabbit) [33-36] than those
73 reported in any other fish tissue and mammals [37,38]. On the other hand, although the
74 physiological function of these LC-PUFAs in sperm is not well known, in mammals they are
75 considered indicators of normal testicular development, spermatogenesis, germ cell populations
76 and fertility [36,39-42] as well as in sperm formation and transportation in the rat testicle
77 [36,38].

78 Biologically active essential fatty acids such as ARA, eicosapentaenoic acid (20:5n-3, EPA) and
79 DHA can be synthesized to some extent by some mammals and freshwater fish through
80 elongation and desaturation of dietary shorter chain precursors. Carnivores and marine fish have
81 only negligible biosynthetic capacity and hence require preformed LC-PUFA in the diet [6,43].
82 However, it was recently demonstrated that desaturation of 22:4n-6 to 22:5n-6 may be carried
83 out by a direct pathway involving a delta 4 desaturase in both a marine herbivorous fish [44], as
84 well as in Senegalese sole larvae [45]. This indicates that there is more than one possible
85 pathway for the synthesis of 22:5n-6 and DHA in vertebrates, i.e., not only the classical
86 'Sprecher pathway' [43].

87 Based on these latest observations, in the present study we conducted a nine-month feeding trial
88 on broodstock G1 Senegalese sole using a standard commercial feed formulation with six graded
89 levels of dietary ARA. The objectives were: (1) to determine the optimal dietary level of ARA
90 for G1 Senegalese sole, using as a reference the fatty acid profile in gonads, liver and muscle of
91 wild broodstock [17]; and (2) to investigate the regulation of fatty acyl desaturase (*d4fad*) and
92 elongase (*elovl5*) gene expression in the liver of G1 sole fed different amounts of ARA.

93 **2. Materials and Methods**

94 Research involving animal experimentation conformed to the principles for the use and care of
95 laboratory animals, in agreement with the Spanish and European regulations on animal welfare
96 (Federation of Laboratory Animal Science Associations, FELASA).

97 **2.1. Fish and Diets**

98 One hundred and twenty Senegalese sole (four year old and 524 ± 11 g average weight), reared
99 in captivity were PIT tagged (AVID, UK) and sexed using a heterologous vitellogenin ELISA
100 for European seabass (*Dicentrarchus labrax*) and validated for Senegalese sole [46]. The fish
101 were distributed among twelve experimental tanks (10 fish per tank, 5 males and 5 females) and
102 fed in duplicate standard commercial (extruded) diet with six graded ARA contents (Tables 1
103 and 2) for nine months (from September 2009 until May 2010). The fish were held in a
104 recirculation system with simulated natural photoperiod and temperature ($40^{\circ} 37'$ and $40^{\circ} 48'$ N
105 and between $0^{\circ} 21'$ and $0^{\circ} 40'$ E., Tarragona, Spain), with minimum temperature observed during
106 two weeks in January - February (13°C) and maximum temperature over twelve weeks during
107 June - September (21°C). The fish were fed six days per week at a daily ration of 0.15- 0.3%
108 body weight.

109 **2.2. Fish Sampling**

110 In May 2010, seventy two fish were sacrificed by pithing the spinal cord (12 fish per dietary
111 treatment, 6 males and 6 females) after anaesthesia with 0.3 ml L^{-1} Aqui-S[®] (Scan Aqua A.S,
112 Årnes, Norway) [47]. Gonads, liver and muscle were collected, and 2 g of liver frozen
113 immediately in liquid nitrogen and subsequently stored at -70°C until RNA extraction, whereas
114 the rest of the tissues for lipid and fatty acid profile were stored at -20°C . All the fish used for

115 these analyses were in advanced stages of sexual maturation, females with vitellogenic oocytes
116 and males containing spermatozoa in the seminiferous tubules.

117 **2.3. Lipid and Fatty Acid Analyses**

118 Samples of tissues and feeds were homogenized and total lipids extracted [48] and quantified
119 gravimetrically. Tissue samples of six males and six females for each diet treatment were
120 analyzed, and feeds were analyzed in triplicate every three months during the experiment. Fatty
121 acid methyl esters were prepared by acid-catalyzed transmethylation [49], and extracted and
122 purified following [50]. Methyl esters were separated and quantified by gas-liquid
123 chromatography (Thermo Trace GC, Thermo Finningan, Milan, Italy) using a 30 m x 0.25 mm
124 ID capillary column (BPX 70, SGE Europe Ltd., UK) with on-column injection and flame
125 ionization detection using helium as carrier gas (1.2 mL min⁻¹ constant flow rate). Individual
126 methyl esters were identified by comparison with known standards (Supelco Inc., Madrid) and a
127 well-characterized fish oil, and quantified in relation to the internal standard, 21:0. The results
128 are presented as percentage of the total fatty acids (TFA) as mean ± standard error of the mean
129 (SEM). Water content was calculated by drying samples at 105°C until a constant weigh was
130 obtained [51].

131 **2.4. Tissue RNA Extraction and Quantitative Real-Time (qRT-PCR)**

132 In order to study the expression of fatty acyl desaturase (*d4fad*) and elongase (*elovl5*) in liver,
133 which is the main metabolic organ where LC-PUFA biosynthesis occurs, total RNA was
134 extracted by organic solvent (Tri-reagent), according to the manufacturer's instructions (Ambion,
135 Applied Biosystems). RNA quality and quantity were assessed by gel electrophoresis and
136 spectrophotometry (NanoDrop ND-1000, Thermo Scientific, Wilmington, USA), respectively.
137 One microgram of total RNA per sample was reverse-transcribed into cDNA using a Verso™
138 cDNA kit (ABgene, Surrey, UK), following the manufacturer's instructions, using a mixture of
139 random hexamers and anchored oligo-dT (3:1, v/v). The cDNA was then diluted 50-fold with
140 water, after a similar amount of cDNA was pooled from all samples. The expression levels of
141 *d4fad* and *elovl5* transcripts were determined by real-time quantitative (qRT-PCR) and
142 normalized using ubiquitin (*UBQ*) and ribosomal protein S4 (*RPS4*) expression.using primers
143 described previously by Morais et al. [45] and normalized using ubiquitin (*UBQ*) and ribosomal

144 protein S4 (*RPS4*) [52]. Details of the primers used can be found in Table 3. The amplification
145 efficiency of the primer pairs was assessed using serial dilutions of cDNA pooled from the
146 samples. All amplifications were carried out in duplicate using an Eppendorf qPCR Cycler
147 (Stanford) in a final volume of 20 μ l containing 2 μ l (for reference genes) or 5 μ l (for *d4fad* and
148 *elovl5*) diluted cDNA (1/50), 0.5 μ M of each primer and 10 μ l of AbsoluteTM qPCR SYBR[®]
149 Green mix (ABgene). Every amplification experiment also included non-template controls
150 (NTC). The RT-qPCR profiles contained an initial activation step at 95°C for 15 min, followed
151 by 35 cycles: 15 s at 95°C, 15 s at the specific primer pair annealing T_m (Table 3), and 30 s at
152 72°C. After the amplification phase, a melt curve of 0.5°C increments from 75°C to 90°C was
153 performed, enabling confirmation of the amplification of a single product in each reaction. The
154 RT-qPCR product sizes were checked by agarose gel electrophoresis and their identity was
155 confirmed by sequencing. No primer–dimer formation occurred in the NTC.

156 **2.5. Statistical Analysis**

157 Statistical differences in lipid and fatty acid compositions among the six experimental groups
158 and wild fish measured previously [17] were analysed separately in males and females by one-
159 way ANOVA followed by the post-hoc multiple comparison test Tukey’s HSD, at a significance
160 level (P) of 0.05. Moreover, correlations among ARA and C22 fatty acids of the n-6 series were
161 calculated at $P = 0.05$. The compliance of data with normality and homogeneity of variance were
162 tested using the Kolmogorov–Smirnov and Bartlett (Chi-Sqr) tests and, when necessary, log-
163 transformation was carried out. Fatty acid content was expressed as mean % TFA \pm SEM.
164 Statistical analysis was performed using the Statistica[®] package for windows (version 6.0;
165 StatSoft Inc, Tulsa, USA).

166 The relative expression of *d4fad* and *elovl5* in fish from groups A (control), C and F was
167 normalized by the expression of *UBQ* and *RPS4* using the normalization factor calculated by
168 geNorm[®] Software version 3.5 [53] and then analyzed for statistical significance using the
169 relative expression software tool (REST-MCS[®], version 2, [http://www.gene-
170 quantification.info/](http://www.gene-quantification.info/)), which employs a pairwise fixed reallocation randomization test (10,000
171 randomizations) with efficiency correction [54].

172

173 **3. Results**

174 After 9 months of feeding the experimental diets the weight of females increased 1.8-fold (from
175 $533 \pm 13\text{g}$ to $950 \pm 25\text{g}$, $\text{SGR} = 0.20\% \text{ day}^{-1}$) whereas weights of males increased 1.5-fold, (from
176 $515 \pm 16\text{g}$ to $755 \pm 25\text{g}$, $\text{SGR} = 0.13\% \text{ day}^{-1}$), with a feed conversion ratio of 1.3 ± 0.04 . No
177 significant differences were noted among the six experimental groups in either growth or feed
178 conversion.

179 **3.1. Fatty acid composition of tissues**

180 The total fatty acid compositions of the different male and female tissues are shown in Tables 4-
181 9. All the tissues analyzed showed a significant accumulation of ARA in a dose dependent
182 manner. Thus, in testis the fish in groups C, D, E and F showed a significantly higher ARA
183 content compared to the control group A, particularly group F which had 13% ARA, 2.7-fold
184 higher than group A with 5% ARA. In the ovary, fish from group F showed a 3.2-fold higher
185 ARA content than group A, whereas groups B, C, D and E did not show significant differences
186 compared to A. The liver of the fish from groups fed diets C, D, E and F had higher ARA levels
187 than control diet A both in males and females. Additionally, males from group B also had
188 significantly higher liver ARA than those from group A, while in females there were no
189 significant differences. The muscle of male fish from groups C, D, E and F showed significantly
190 higher ARA levels than those from group A, whereas in females only those from groups E and F
191 had significantly higher ARA levels than group A.

192 In the case of 22:4n-6 and 22:5n-6 contents, fish fed the six experimental diets showed a dose
193 dependent and significant increase in testis (Table 4) and liver (Table 6) of male fish, whereas no
194 differences could be found in females, although a similar trend was observed but with much
195 smaller differences (Table 5 and 7). The relative level of 22:4n-6 in testis was significantly
196 higher in groups B (2.2-fold), C (4.2-fold), D (4.5-fold), E (6.1-fold) and F (8.1-fold) compared
197 with the lowest levels found in the control group A (0.2% TFA) (Table 4). The 22:5n-6 content
198 in testis was also higher in groups C (1.5-fold), D (1.8-fold), E (2.2-fold) and F (2.4-fold)
199 compared with the control group A (0.4% TFA). The 22:4n-6 levels in the liver of males were
200 also significantly higher in groups E (7.1-fold) and F (9.5-fold) compared to control group A

201 (0.1% TFA), while 22:5n-6 was 5-fold higher in group F compared with group A (0.2% TFA)
202 (Table 6).

203 As a consequence of the increasing ARA levels, the EPA/ARA ratio was significantly reduced in
204 similar dose-dependent manner in gonads (Tables 4 and 5), liver (Tables 6 and 7) and muscle
205 (Tables 8 and 9) of both males and females. The increase of ARA in gonad, liver and muscle of
206 males resulted in a concomitant significant increase in total n-6 PUFA whereas a significant
207 reduction in total n-3 PUFA was only observed in testis.

208 **3.2. Comparison of tissue ARA levels with wild fish**

209 The tissue ARA levels in the present study were compared those of wild broodstock reported
210 previously [17], and different results were obtained depending on the tissue (Fig. 1). In gonads
211 ARA was significantly higher in males compared with females however in liver and muscle no
212 differences in ARA content between males and females were observed. The levels of ARA in
213 testis of fish from groups E and F showed significantly higher accumulation of ARA compared
214 with those from wild fish. However, ARA levels in testis from groups A, B, C, and D were
215 similar and not significantly different to those from wild fish. In ovary no differences in ARA
216 between the experimental groups and wild fish were observed. ARA content in liver (males and
217 females) from groups A and B were significantly lower than the wild group. However, groups C,
218 D, E and F showed no differences compared to the wild group. In muscle of males the ARA
219 levels in all the fish groups were significantly lower compared to wild fish, and the same as in
220 females from groups A, B, C and D. However, females from groups E and F showed similar
221 ARA levels to that found in the wild.

222 **3.3. Elongase and desaturase gene expression (RT-qPCR)**

223 The expression of *d4fad* and *elovl5* increased in a dose dependent manner in fish fed diets C and
224 F in comparison to the control (A) treatment, but only in males (Fig. 2). Levels of *d4fad*
225 transcripts were significantly higher in the liver of male fish from groups C (3.2-fold) and F (6.8-
226 fold). On the other hand, expression of *elovl5* increased significantly in group F, being 5.3-fold
227 higher than in group A, but not in group C (1.8-fold increase).

228

229

230 4. Discussion

231 One of the main bottlenecks to the culture of Senegalese sole, a species with high market value
232 and interest for aquaculture diversification in Southern Europe, is the poor reproductive
233 performance of G1 fish. Previous studies have indicated that the problem is more likely
234 associated with male sperm production and quality than female performance [55-58]. When
235 comparing the fatty acid profile of cultured and wild fish major differences were found in the
236 contents of ARA which, due to its important multiple roles in reproduction, was deemed a
237 potential factor associated with reproductive failure of fish produced in captivity. Hence, the
238 objective of the present study was to test a gradient of dietary ARA to determine the levels that
239 would raise ARA tissue contents in G1 fish to those similar to wild fish. In addition, given
240 previous observations that increased ARA in the tissue and blood of fish raise the levels of its
241 elongated and desaturated products, 22:4n-6 and 22:5n-6, [17,32], the pathway of LC-PUFA
242 biosynthesis was also studied by assessing the expression levels of two genes coding for Elov15
243 and d4Fad, which are two key enzymes in this pathway.

244 The results obtained show that the relative content of ARA in tissues of G1 Senegalese sole
245 generally correlated with dietary ARA levels, increasing in a dose-dependent manner in both
246 sexes. Feeding fish with diets containing 2.3% and 3.2% ARA levels (diets C and D,
247 respectively) resulted in similar ARA contents in testis, ovary and liver compared to wild fish.
248 Muscle, on the other hand, did not accumulate ARA at the same rate as liver and gonads and
249 only the female fish fed diets containing 5% or higher ARA achieved similar ARA levels in
250 muscle as the wild fish. A previous study on dietary ARA preference of Senegalese sole
251 broodstock using a self-feeding system over 16 months revealed that fish regulate the ingestion
252 of ARA to around 3.0% TFA in diet and the resulting ARA content in the tissues was 8.9% TFA
253 in testis, 4% in ovary and 2.5% in liver [59]. These values were similar to those found previously
254 in wild Senegalese sole [17] and also to those obtained in the present study in the testis, ovary
255 and liver of G1 fish in groups C and D. Thus, the ARA accumulation in the muscle was lower
256 than in the self-feeding experiment, except for females of groups E and F. Considering that the
257 self-feeding experiment was conducted for 16 months and the present study lasted only 9
258 months, optimal dietary ARA level seems to be dependent on the duration of the feeding period.
259 If fish are fed for a period shorter than nine months it might be appropriate to use at least 3.2%

260 ARA in the diet. For longer feeding periods, 2.3% ARA in the diet might be sufficient for
261 Senegalese sole G1, such that no differences in ARA content could be found in the testis and
262 ovary of the G1 fish compared to wild fish, or fish in the self-feeding experiment. Alornd [4],
263 studied ARA requirements for G1 Atlantic halibut over 3 years and suggested 2.3% as the
264 optimal dietary content, as this level led to the longest milt production period and highest
265 fecundity. This result is likely due to ARA being rapidly incorporated into the reproductive
266 tissues [4].

267 Previous studies on ARA requirements for broodstock fish showed that the optimal dietary level
268 is species-specific, with higher or lower ARA levels producing detrimental effects in
269 reproductive physiology [2-4,11,12,15,30,60]. A significant increase in the production of steroids
270 was observed in Senegalese sole males fed 3.2% ARA [32], and the highest egg production was
271 obtained using diets with 3.6% ARA for Japanese flounder (*Paralichthys olivaceus*) [3]. On the
272 other hand, a study with Atlantic cod (*Gadus morhua*) found that a diet with 4% ARA, increased
273 the production of estradiol and extended the length of the spawning season [60]. However,
274 negative effects on steroid production, fecundity, egg and larval quality have been observed
275 when fish were fed high ARA levels. Japanese flounder fed 7.3% ARA exhibited a significant
276 reduction in egg and larval quality [3] whereas Atlantic halibut fed 3.2% ARA showed a delay in
277 the spawning season [4]. Other effects such as an earlier estradiol peak in Senegalese sole [32],
278 and an earlier peak in estradiol and vitellogenesis in Atlantic cod [60] were also observed. More
279 research is required to establish the effects of dietary ARA on sperm and oocyte quality in *S.*
280 *senegalensis* and the feeding time required to incorporate ARA into the reproductive organs, but
281 the present study suggests that dietary levels between 2.3% and 3.2%, depending on the duration
282 of the feeding, might improve the reproductive performance of Senegalese sole G1.

283 The results obtained in the present study also showed that dietary ARA was preferentially
284 transferred to and accumulated in the gonads (testis and ovary), followed by the liver and
285 muscle, similarly to what had been previously observed in Atlantic halibut, white seabream,
286 black seabream and silver pomfret [4,22,23,61,62].

287 A positive correlation between the level of ARA and the concentrations of 22:4n-6 and 22:5n-6
288 in testis, liver and muscle of wild fish had been shown previously [17], and it was suggested that
289 this probably occurred from metabolism (elongation and desaturation) of ARA. A similar pattern

290 of 22:4n-6 and 22:5n-6 concentration in the tissues after graded dietary ARA was observed in the
291 testis and liver of males in the present study. Accumulation of these fatty acids in the muscle of
292 wild black seabream [23] and Senegalese sole [17], in the gonads of seabass [63], seabream [21],
293 silver pomfret [62] and in the sperm of seabass [29] and rainbow trout [28,29] have been
294 reported previously, with the accumulation of 22:5n-6 being suggested as deriving from local
295 production and uptake from the circulatory system [64-66]. These fatty acids are found in storage
296 lipids in testis of mammals [38] and are considered indicators of normal testicular and sperm
297 condition [36-42]. Although the physiological function of these LC-PUFA in sperm is not well
298 understood, Lenzie et al. [36] suggested that they are involved in sperm formation and in
299 fertilization and, in mammals, there is an increase in the degree of fatty acid metabolism and
300 desaturation during spermatogenesis and sperm maturation [42]. However, the function of 22:4n-
301 6 and 22:5n-6 in fish and their effects on reproduction and spermatogenesis have not been
302 established. It has been suggested that 22:5n-6 is accumulated in the testis as an ARA reservoir,
303 being retro-converted into ARA by hydrogenation and subsequent oxidation [37,64,65,67].
304 Cultured fish fed commercial extruded diets show a significantly lower accumulation of 22:5n-6
305 and 22:4n-6 compared to wild fish [17]. Nonetheless, in the present study, when fish were fed
306 increasing ARA levels, a parallel increase in these ARA-derived fatty acids was observed,
307 especially in the liver and testis. Recently, the cloning and characterization of *elov15* and *d4fad*
308 transcripts in Senegalese sole revealed that this species is able to elongate ARA to 22:4n-6 and
309 then directly desaturate this substrate to 22:5n-6 [45]. Although the activity of this pathway
310 could not be assessed *in vivo*, results from this experiment suggest that it is physiologically
311 relevant given that ARA induced a marked dose-dependent up-regulation in the expression of
312 both *elov15* and *d4fad* in liver, with *elov15* transcripts increasing 5.3-fold in males fed 6.0% ARA
313 in relation to those fed 0.7%, and *d4fad* being up-regulated 3.2- and 6.8-fold in fish fed diets
314 containing 2.3% and 6.0% ARA, respectively. However, an interesting observation was that the
315 up-regulation of these two genes was only observed in males, which was also clearly reflected in
316 the fatty acid composition of the tissues, where significant changes in 22:4n-6 and 22:5n-6
317 contents between treatments were only observed in testis, liver and flesh of males whereas in
318 female tissues only ARA levels were significantly different between fish fed the experimental
319 diets. Gender differences in liver desaturase expression have also been observed in Wistar rats
320 fed an n-3 PUFA enriched diet [68]. In that case, desaturation activity was significantly increased

321 in the females compared to males. It was also shown that female rats have higher plasma DHA
322 concentrations than males [69]. It is conceivable that the LC-PUFA biosynthesis pathway in
323 female fish liver is also more directed to the provision of DHA for later incorporation into eggs
324 and, at least in post-larvae, the expression of *d4fad* was up-regulated at lower dietary levels of n-
325 3 LC-PUFA (mainly DHA and EPA) [45]. However, in the present experiment only the levels of
326 ARA varied significantly between diets and hence *elovl5* and *d4fad* expression was associated
327 with the conversion of ARA into 22:4n-6 and subsequently to 22:5n-6 in G1 male fish, both of
328 which are important fatty acids involved in testis and sperm composition in mammals [39-42].
329 Further studies are necessary to understand the importance of these two fatty acids and their
330 physiological function in male fish reproduction, but the present results suggest that they may be
331 as important in Senegalese sole as in higher vertebrates.

332 **5. Conclusion**

333 Based on previous data on ARA content in wild Senegalese sole [17], and considering the results
334 presented here, fish fed either 2.3% or 3.2% ARA enriched diets showed levels of ARA in testis,
335 ovary and liver compared to those of wild fish. Thus, diets with 3.2% ARA for feeding periods
336 up to nine months or 2.3% ARA diets for prolonged feeding periods are suggested. The ARA
337 was preferentially transferred and conserved in the gonads (testis and ovary), followed by the
338 liver and muscle. The increase in the expression of *elovl5* and *d4fad* transcripts in liver in
339 response to dietary ARA content and a parallel increase in tissue 22:4n-6 and 22:5n-6 levels
340 suggest the ability of Senegalese sole to elongate ARA to 22:4n-6 followed by desaturation to
341 22:5n-6, suggesting an important role of these ARA-derived fatty acids in male fish
342 reproduction. Further studies are required to establish the dietary ARA effect on reproductive
343 performance of Senegalese sole and the time required for effective incorporation of ARA into
344 reproductive organs.

345 **Acknowledgments**

346 Thanks to N. Gras, G. Macia and J.R. Dick for technical support and G. Rosenlund from
347 Skretting ARC (Norway) for formulating and providing the diets used in the study. This research
348 was funded by Ministerio de Agricultura, Pesca, Alimentación y Medio Ambiente -JACUMAR
349 Project Lenguado coordinated nationally by Pedro Cañavate and in IRTA by N.D. and (INIA)-

350 FEDER project RTA2005-00113-00-00 coordinated by N.D. F.N. thanks Agència de Gestió
351 d'Ajuts Universitaris i de Recerca (AGAUR) for the PhD grant he was awarded.

352 6. Reference

353

- 354 [1] Sargent J, Bell G, McEvoy L, Tocher D, Estevez A. Recent developments in the essential fatty acid
355 nutrition of fish. *Aquaculture* 1999; **177**: 191-199.
- 356 [2] Mazorra C, Bruce M, Bell JG, Davie A, Alorend E, Jordan N *et al.* Dietary lipid enhancement of
357 broodstock reproductive performance and egg and larval quality in Atlantic halibut
358 (*Hippoglossus hippoglossus*). *Aquaculture* 2003; **227**: 21-33.
- 359 [3] Furuita H, Yamamoto T, Shima T, Suzuki N, Takeuchi T. Effect of arachidonic acid levels in
360 broodstock diet on larval and egg quality of Japanese flounder *Paralichthys olivaceus*.
361 *Aquaculture* 2003; **220**: 725-735.
- 362 [4] Alorend E. The effect of dietary arachidonic acid concentration on Atlantic Halibut (*Hippoglossus*
363 *hippoglossus*) broodstock performance, assessment of egg, milt and larval quality. *PhD Thesis,*
364 *Institute of Aquaculture* University of Stirling, UK; 2004. p. 176.
- 365 [5] Meunpol O, Meejing P, Piyatiratitivorakul S. Maturation diet based on fatty acid content for
366 male *Penaeus monodon* (Fabricius) broodstock. *Aquac Res* 2005; **36**: 1216-1225.
- 367 [6] Tocher DR. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquac Res* 2010;
368 **41**: 717-732.
- 369 [7] Smith WL, Murphy RC, Dennis E, Vance JEV. The eicosanoids: cyclooxygenase, lipoxygenase, and
370 epoxygenase pathways. *New Compr Biochem* 2002; **36**: 341-371.
- 371 [8] Tocher DR. Metabolism and functions of lipids and fatty acids in teleost fish. *Rev Fish Sci* 2003;
372 **11**: 107-184.
- 373 [9] Van der Kraak G, Chang JP. Arachidonic acid stimulates steroidogenesis in goldfish preovulatory
374 ovarian follicles. *Gen Comp Endocr* 1990; **77**: 221-228.
- 375 [10] Wade MG, Van der Kraak G. Arachidonic acid and prostaglandin E2 stimulate testosterone
376 production by goldfish testis in vitro. *Gen Comp Endocr* 1993; **90**: 109-118.
- 377 [11] Mercure F, Van der Kraak G. Inhibition of gonadotropin-stimulated ovarian steroid production
378 by polyunsaturated fatty acids in teleost fish. *Lipids* 1995; **30**: 547-554.
- 379 [12] Sorbera LA, Asturiano JF, Carrillo M, Zanuy S. Effects of polyunsaturated fatty acids and
380 prostaglandins on oocyte maturation in a marine teleost, the European Sea bass (*Dicentrarchus*
381 *labrax*). *Biol Reprod* 2001; **64**: 382-389.
- 382 [13] Sorensen PW, Stacey NE. Brief review of fish pheromones and discussion of their possible uses
383 in the control of non-indigenous teleost fishes. *New Zeal J Mar Fresh* 2004; **38**: 399-417.
- 384 [14] Wang X, Stocco DM. The decline in testosterone biosynthesis during male aging: A consequence
385 of multiple alterations. *Mol Cell Endocrinol* 2005; **238**: 1-7.
- 386 [15] Mercure F, Van der Kraak G. Mechanisms of Action of Free Arachidonic Acid on Ovarian Steroid
387 Production in the Goldfish. *Gen Comp Endocr* 1996; **102**: 130-140.
- 388 [16] Carazo I, Martin I, Chereguini O, Mañanós E, Duncan N. The absence of reproductive behaviour
389 in cultured (G1 generation) Senegalese sole (*Solea senegalensis*) explains poor reproductive
390 performance. *Proceeding of 5th Workshop on the Cultivation of Soles* Centre of Marine
391 Sciences, University of the Algarve, Faro, Portugal, 5-7 April; 2011. p. 12
- 392 [17] Norambuena F, Estevez A, Bell G, Carazo I, Duncan N. Proximate and fatty acid composition in
393 muscle, liver and gonads of wild versus cultured broodstock of Senegalese sole (*Solea*
394 *senegalensis*). *Aquaculture* 2012; **356-357**: 176-185.

- 395 [18] Harrell RM, Woods LC. Comparative fatty acid composition of eggs from domesticated and wild
396 striped bass (*Morone saxatilis*). *Aquaculture* 1995; **133**: 225-233.
- 397 [19] Sheikh-Eldin M, De Silva SS, Anderson TA, Gooley G. Comparison of fatty acid composition of
398 muscle, liver, mature oocytes, and diets of wild and captive Macquarie perch, *Macquaria*
399 *australasica*, broodfish. *Aquaculture* 1996; **144**: 201-216.
- 400 [20] Silversand C, Norberg B, Haux C. Fatty-acid composition of ovulated eggs from wild and cultured
401 turbot (*Scophthalmus maximus*) in relation to yolk and oil globule lipids. *Mar Biol* 1996; **125**:
402 269-278.
- 403 [21] Cejas JR, Almansa E, Villamandos JE, Badía P, Bolaños A, Lorenzo A. Lipid and fatty acid
404 composition of ovaries from wild fish and ovaries and eggs from captive fish of white sea bream
405 (*Diplodus sargus*). *Aquaculture* 2003; **216**: 299-313.
- 406 [22] Cejas JR, Almansa E, Jerez S, Bolaños A, Samper M, Lorenzo A. Lipid and fatty acid composition
407 of muscle and liver from wild and captive mature female broodstocks of white seabream,
408 *Diplodus sargus*. *Comp Biochem Physiol B* 2004; **138**: 91-102.
- 409 [23] Rodríguez C, Acosta C, Badía P, Cejas JR, Santamaría FJ, Lorenzo A. Assessment of lipid and
410 essential fatty acids requirements of black seabream (*Spondyliosoma cantharus*) by comparison
411 of lipid composition in muscle and liver of wild and captive adult fish. *Comp Biochem Physiol B*
412 2004; **139**: 619-629.
- 413 [24] Salze G, Tocher DR, Roy WJ, Robertson DA. Egg quality determinants in cod (*Gadus morhua* L.):
414 egg performance and lipids in eggs from farmed and wild broodstock. *Aquac Res* 2005; **36**: 1488-
415 1499.
- 416 [25] Blanchet C, Lucas M, Julien P, Morin R, Gingras S, Dewailly E. Fatty acid composition of wild and
417 farmed Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*). *Lipids* 2005; **40**:
418 529-531.
- 419 [26] Aslan SS, Guven KC, Gezgin T, Alpaslan M, Tekinay A. Comparison of fatty acid contents of wild
420 and cultured rainbow trout *Oncorhynchus mykiss* in Turkey. *Fisheries Sci* 2007; **73**: 1195-1198.
- 421 [27] Lund I, Steenfeldt SJ, Suhr KI, Hansen BW. A comparison of fatty acid composition and quality
422 aspects of eggs and larvae from cultured and wild broodstock of common sole (*Solea solea* L.).
423 *Aquacult Nutr* 2008; **14**: 544-555.
- 424 [28] Vassallo-Agius R, Watanabe T, Yoshizaki G, Satoh S, Takeuchi Y. Quality of eggs and spermatozoa
425 of rainbow trout fed an n-3 essential fatty acid-deficient diet and its effects on the lipid and fatty
426 acid components of eggs, semen and livers. *Fisheries Sci* 2001; **67**: 818-827.
- 427 [29] Bell MV, Dick JR, Thrush M, Navarro JC. Decrease 20:4n-6/20:5n-3 ratio in sperm from cultured
428 sea bass, *Dicentrarchus labrax*, broodstock compared with wild fish *Aquaculture* 1996; **144**: 189-
429 199.
- 430 [30] Norambuena F, Mackenzie S, Bell JG, Callol A, Estevéz A, Duncan N. Prostaglandin (F and E, 2-
431 and 3-series) production and cyclooxygenase (COX-2) gene expression of wild and cultured
432 broodstock of Senegalese sole (*Solea senegalensis*). *Gen Comp Endocr* 2012; **177**: 256-262.
- 433 [31] Baron SF, Hylemon PB. Biotransformation of bile acids, cholesterol, and steroids hormones. In:
434 White RL MaBA (ed). *Gastrointestinal microbiology*. London, England: Chapman and Hall; 1997.
435 p. 663.
- 436 [32] Norambuena F. Senegalese sole (*Solea senegalensis*) broodstock nutrition: arachidonic acid
437 (20:4n-6, ARA) and reproductive physiology. *PhD Thesis, Institute of Animal Biology, Vegetal and*
438 *Ecology*: Autonomous University of Barcelona, Spain; 2012. p. 187.
- 439 [33] Ahaluwalia B, Holman RT. Fatty acid composition of lipid of bull, boar, rabbit and human semen.
440 *J Reprod Fertil* 1969; **18**: 431-437.
- 441 [34] Tinoco J. Dietary requirements and functions of α -linolenic acid in animals. *Prog Lipid Res* 1982;
442 **21**: 1-45.

- 443 [35] Picardo M, Gramatico P, De Luca C, Passi S, Nazzaro-Porro M. Differential effect of azelaic acid in
444 melanoma cell cultures. *Pigm Cell Res* 1990; **20**: 142-149.
- 445 [36] Lenzi A, Picardo M, Gandini L, Dondero F. Lipids of the sperm plasma membrane: from
446 polyunsaturated fatty acids considered as markers of sperm function to possible scavenger
447 therapy. *Hum Reprod Update* 1996; **2**: 246-256.
- 448 [37] Bridges RB, Coniglio JG. The metabolism of 4,7,10,13,16-[5-14c]docosapentaenoic acid in the
449 testis of the rat. *BBA-Lipid Lipid Met* 1970; **218**: 29-35.
- 450 [38] Ayala S, Gaspar G, Brenner RR, Peluffo RIO, Kunau W. Fate of linoleic, arachidonic, and docosa-
451 7,10,13,16-tetraenoic acids in rat testicles. *J Lipid Res* 1973; **14**: 296-305.
- 452 [39] Leat WMF, Northrop CA, Harrison FA, Cox RW. Effect of dietary linoleic and linolenic acids on
453 testicular development in the rat. *Exp Physiol* 1983; **68**: 221-231.
- 454 [40] MacDonald ML, Rogers QR, Morris JG, Cupps PT. Effects of linoleate and arachidonate
455 deficiencies on reproduction and spermatogenesis in the cat. *J Nutr* 1984; **114**: 719-726.
- 456 [41] Lenzi A, Gandini L, Maresca V, Rago R, Sgro P, Dondero F *et al.* Fatty acid composition of
457 spermatozoa and immature germ cells. *Mol Hum Reprod* 2000; **6**: 226-231.
- 458 [42] Furland NE, Maldonado EN, Aresti PA, Aveldaño MI. Changes in lipids containing long- and very
459 long-chain polyunsaturated fatty acids in cryptorchid rat testes. *Biol Reprod* 2007; **77**: 181-188.
- 460 [43] Sprecher H. Metabolism of highly unsaturated n-3 and n-6 fatty acids. *Biochim Biophys Acta*
461 2000; **1486**: 219-231.
- 462 [44] Li Y, Monroig O, Zhang L, Wang S, Zheng X, Dick JR *et al.* Vertebrate fatty acyl desaturase with $\Delta 4$
463 activity. *P Natl Acad Sci USA* 2010; **107**: 16840-16845.
- 464 [45] Morais S, Castanheira F, Martinez-Rubio L, Conceição LEC, Tocher DR. Long chain
465 polyunsaturated fatty acid synthesis in a marine vertebrate: Ontogenetic and nutritional
466 regulation of a fatty acyl desaturase with delta 4 activity. *BBA- Mol Cell Biol L* 2012; **1821**: 660-
467 671.
- 468 [46] Mañanós E, Núñez J, Zanuy S, Carrillo M, Le Menn F. Sea bass (*Dicentrarchus labrax* L.)
469 vitellogenin. II-Validation of an enzyme-linked immunosorbent assay (ELISA). *Comp Biochem*
470 *Physiol B* 1994; **107**: 217-223.
- 471 [47] Norambuena F, Carazo I, Duncan N, Estevez A, Roque A. Anesthesia of broodstock
472 mediterranean fish. *Humane Slaughter Association Centenary International Symposium: Recent*
473 *Advances in the Welfare of Livestock at Slaughter* Portsmouth Historic Dockyard, Portsmouth,
474 UK, 30 June - 1 July; 2011. p. 30.
- 475 [48] Folch JM, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total
476 lipids from animal tissues. *J Biol Chem* 1957; **226**: 497-509.
- 477 [49] Christie WW. *Lipid Analysis*. Pergamon, Oxford, UK; 1982. p. 207
- 478 [50] Tocher DR, Harvie DG. Fatty acid compositions of the major phosphoglycerides from fish neural
479 tissues; (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout (*Salmo gairdneri*) and cod
480 (*Gadus morhua*) brains and retinas. *Fish Physiol Biochem* 1988; **5**: 229-239.
- 481 [51] AOAC. Crude protein content by the Dumas procedure, by means of a Nitrogen/protein FP-528
482 determinator and moisture. *Official Methods of Analysis*. Washington, USA: Association of
483 Official Analytical Chemist; 2000.
- 484 [52] Infante C, Matsuoka M, Asensio E, Canavate J, Reith M, Manchado M. Selection of housekeeping
485 genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Mol Biol* 2008;
486 **9**: 28.
- 487 [53] Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A *et al.* Accurate
488 normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal
489 control genes. *Gen Biol* 2002; **3**: 34.31-34.11.

- 490 [54] Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST©) for group-wise
491 comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids*
492 *Res* 2004.
- 493 [55] Guzmán JM, Cal R, García-López A, Chereguini O, Kight K, Olmedo M *et al.* Effects of in vivo
494 treatment with the dopamine antagonist pimozide and gonadotropin-releasing hormone agonist
495 (GnRHa) on the reproductive axis of Senegalese sole (*Solea senegalensis*). *Com Biochem Phys A*
496 2011; **158**: 235-245.
- 497 [56] Agulleiro MJ, Anguis V, Cañavate JP, Martínez-Rodríguez G, Mylonas CC, Cerda J. Induction of
498 spawning of captive-reared Senegal sole (*Solea senegalensis*) using different administration
499 methods for gonadotropin-releasing hormone agonist. *Aquaculture* 2006; **257**: 511-524.
- 500 [57] Agulleiro MJ, Scott AP, Duncan N, Mylonas CC, Cerdà J. Treatment of GnRHa-implanted
501 Senegalese sole (*Solea senegalensis*) with 11-ketoandrostenedione stimulates spermatogenesis
502 and increases sperm motility. *Comp Biochem Phys A* 2007; **147**: 885-892.
- 503 [58] Forné I, Agulleiro MJ, Asensio E, Abián J, Cerdà J. 2-D DIGE analysis of Senegalese sole (*Solea*
504 *senegalensis*) testis proteome in wild-caught and hormone-treated F1 fish. *Proteomics* 2009; **9**:
505 2171-2181.
- 506 [59] Norambuena F, Sánchez-Vásquez J, Estevez A, Carazo I, Duncan N. Self-selection of diets with
507 different content of arachidonic acid by Senegalese sole (*Solea senegalensis*) broodstock.
508 *Aquaculture, submitted* 2012.
- 509 [60] Norberg B, Hamre K, Araujo P, van der Meeren T, Rosenlund G. Effect of graded levels of
510 arachidonic acid on reproductive physiology, spawning performance, and egg quality of atlantic
511 cod *5th Fish & shellfish larviculture symposium, European Aquaculture Society* Oostende,
512 Belgium, 7-11 September; 2009.
- 513 [61] Pérez MJ, Rodríguez C, Cejas JR, Martín MV, Jerez S, Lorenzo A. Lipid and fatty acid content in
514 wild white seabream (*Diplodus sargus*) broodstock at different stages of the reproductive cycle.
515 *Comp Biochem Physiol B* 2007; **146**: 187-196.
- 516 [62] Huang X, Yin Y, Shi Z, Li W, Zhou H, Lv W. Lipid content and fatty acid composition in wild-caught
517 silver pomfret (*Pampus argenteus*) broodstocks: Effects on gonad development. *Aquaculture*
518 2010; **310**: 192-199.
- 519 [63] Bell JG, Farndale BM, Bruce MP, Navas JM, Carillo M. Effects of broodstock dietary lipid on fatty
520 acid compositions of eggs from sea bass (*Dicentrarchus labrax*). *Aquaculture* 1997; **149**: 107-119.
- 521 [64] Rosenthal MD, Garcia MC, Jones MR, Sprecher H. Retroconversion and D4 desaturation of
522 docosatetraenoate (22:4(n-6)) and docosapentaenoate (22:5(n-3)) by human cells in culture.
523 *BBA-Lipid Lipid Met* 1991; **1083**: 29-36.
- 524 [65] Tam P, Umeda-Sawada R, Yaguchi T, Akimoto K, Kiso Y, Igarashi O. The metabolism and
525 distribution of docosapentaenoic acid (n-6) in rats and rat hepatocytes. *Lipids* 2000; **35**: 71-75.
- 526 [66] Tam PS, Sawada R, Cui Y, Matsumoto A, Fujiwara Y. The metabolism and distribution of
527 docosapentaenoic acid (n-6) in the liver and testis of growing rats. *Biosci Biotechnol Biochem*
528 2008; **72**: 2548-2554.
- 529 [67] Kunau W. Partial degradation of 4.7.10.13.16-docosapentaenoic acid in rat liver. *Febs Lett* 1971;
530 **16**: 54-56.
- 531 [68] Extier A, Langelier Bnd, Perruchot M-Hln, Guesnet P, Van Veldhoven PP, Lavielle M *et al.* Gender
532 affects liver desaturase expression in a rat model of n-3 fatty acid repletion. *J Nutr Biochem*
533 2010; **21**: 180-187.
- 534 [69] Childs CE, Romeu-Nadal M, Burdige GC, Calder PC. Gender differences in the n-3 fatty acid
535 content of tissues. *P Nut Soc* 2008; **67**: 19-27.
- 536
537

538 **Fig. 1** Arachidonic acid content (20:4n-6, ARA) in a) gonads, b) liver and c) muscle of wild fish
539 (Norambuena et al., 2012a) compared with cultured fish fed with different dietary ARA levels
540 (A= 0.7% ARA, B=1.6% ARA, C= 2.3% ARA,D= 3.2% ARA, E= 5.0% ARA and F= 6.0% ARA.
541 Different letters indicate significant differences (ANOVA, P<0.05, N=6) between fish groups.
542 Capital letters are used for males and small letters for females. (*) Indicate significant
543 differences (P<0.05) between males and females.

544

545 **Fig. 2** Relative expression (RT-qPCR) of fatty acyl (a) elongase (elovl5) and (b) desaturase
546 (d4fad) in the liver of female and male Senegalese sole in relation to group A (control),
547 normalized by the expression of UBQ and RPS4 (reference genes). (*) Denote significant
548 differences of groups C and F with respect to control (group A), calculated by REST (P<0.05,
549 N=6).

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576


577

578


579

580

581



583
 584
 585
 586
 587
 588
 589
 590
 591



593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619

620 **Table 1.** *Ingredients and proximate composition of the experimental diets (A, B, C, D, E and F)*

Ingredients (g/Kg)	A	B	C	D	E	F
Fish meal ¹	645.0	645.0	645.0	645.0	645.0	645.0
Wheat gluten ²	120.0	120.0	120.0	120.0	120.0	120.0
Wheat ³	125.8	125.8	125.8	125.8	125.8	125.8
Fish oil ⁴	80.0	76.0	71.8	67.6	63.2	59.0
Vevodar ⁵	0.0	4.0	8.2	12.4	16.8	21.0
Premixes ⁶	29.2	29.2	29.2	29.2	29.2	29.2
Analysed values						
Moisture, %	8.0	7.8	8.3	8.4	8.6	8.3
Crude protein, % DM ⁷	61,2	61,4	61,6	61,7	61,8	62,2
Crude fat, % DM	13.8	14.1	14.4	13.7	14.1	14.3

¹ LT fish meal, Skretting, Stavanger, Norway

² Cargill Nordic, Charlottenlund, Denmark

³ Skretting, Stavanger, Norway

⁴ Scandinavian fish oil, Skretting, Stavanger, Norway

⁵ Contains 35% arachidonic acid, DSM Food Specialities, Delft, The Netherlands

⁶ Include micronutrients, vitamin and mineral supplementation. Trouw Nutrition, Boxmeer, Netherlands, proprietary composition Skretting ARC

⁷ Dry matter

622
623
624

Table 2. Lipid, fatty acid content and fatty acid composition (% TFA \pm SEM) of the diets used (A, B, C, D, E and F) for feeding G1 Senegalese sole (*Solea senegalensis*). Columns assigned different letters were significantly different (ANOVA, $P < 0.05$, $N = 3$)

	A	B	C	D	E	F
TFA ($\mu\text{g mg}^{-1}$ L)	880 \pm 87	702 \pm 34	820 \pm 74	861 \pm 82	961 \pm 19	911 \pm 94
Fatty acid composition (%TFA)						
14:0	2.5 \pm 1.2	3.6 \pm 2.8	3.8 \pm 2.1	3.6 \pm 2.3	3.7 \pm 2.8	4.0 \pm 2.9
16:0	14.9 \pm 1.1	15.3 \pm 3.7	17.6 \pm 2.6	15.8 \pm 1.8	15.0 \pm 2.8	16.1 \pm 2.7
18:0	1.8 \pm 0.9	2.4 \pm 0.9	2.6 \pm 0.7	2.6 \pm 0.6	3.0 \pm 0.7	3.2 \pm 0.4
Total SFA	19.4 \pm 1.5	21.7 \pm 5.7	24.4 \pm 4.2	22.3 \pm 3.6	21.9 \pm 4.8	23.4 \pm 5.2
16:1n-7	4.9 \pm 0.8	4.8 \pm 2.0	5.0 \pm 0.8	4.9 \pm 0.8	4.3 \pm 1.8	4.4 \pm 1.2
18:1n-9	15.0 \pm 1.3	15.7 \pm 2.4	16.9 \pm 1.9	15.9 \pm 2.0	15.1 \pm 2.7	15.7 \pm 0.6
18:1n-7	1.2 \pm 2.0	0.9 \pm 1.6	1.0 \pm 1.7	0.8 \pm 1.4	0.8 \pm 1.4	1.0 \pm 1.7
20:1n-9	7.2 \pm 1.5	7.1 \pm 0.3	7.2 \pm 1.1	6.3 \pm 0.5	6.9 \pm 0.7	6.6 \pm 0.4
22:1n-9	3.5 \pm 6.0	3.9 \pm 6.7	2.9 \pm 5.0	3.0 \pm 5.2	3.9 \pm 6.7	2.9 \pm 5.1
Total MUFA	32.3 \pm 10.5	32.7 \pm 4.7	33.4 \pm 4.4	31.3 \pm 4.9	31.4 \pm 5.5	31.1 \pm 6.0
18:2n-6	5.9 \pm 0.7	6.4 \pm 0.7	6.0 \pm 0.9	6.6 \pm 0.4	5.9 \pm 0.1	7.2 \pm 0.7
20:4n-6, ARA	0.7 \pm 0.3 ^c	1.6 \pm 0.6 ^c	2.3 \pm 0.8 ^{bc}	3.2 \pm 0.7 ^b	5.0 \pm 0.6 ^a	6.0 \pm 0.1 ^a
22:4n-6	0.1 \pm 0.1	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0
22:5n-6	0.3 \pm 0.6	0.3 \pm 0.2	0.2 \pm 0.3	0.2 \pm 0.3	0.2 \pm 0.2	0.3 \pm 0.4
Total n-6 PUFA	9.3 \pm 3.4	8.4 \pm 1.3	8.9 \pm 1.5	10.6 \pm 0.8	12.4 \pm 1.0	14.0 \pm 1.6
18:3n-3	1.3 \pm 0.2	1.4 \pm 0.2	1.3 \pm 0.3	1.3 \pm 0.2	1.2 \pm 0.1	1.2 \pm 0.1
18:4n-3	2.2 \pm 0.3	2.0 \pm 0.1	1.8 \pm 0.3	2.0 \pm 0.1	1.8 \pm 0.2	1.7 \pm 0.2
20:4n-3	0.7 \pm 0.1	0.7 \pm 0.1	0.6 \pm 0.1	0.7 \pm 0.0	0.6 \pm 0.1	0.6 \pm 0.1
20:5n-3, EPA	13.0 \pm 8.4	16.8 \pm 5.3	15.9 \pm 5.4	16.4 \pm 5.4	14.8 \pm 6.0	14.7 \pm 4.9
22:5n-3, DPA	1.6 \pm 0.4	3.0 \pm 2.3	2.0 \pm 1.0	2.3 \pm 1.3	4.6 \pm 5.7	2.0 \pm 0.9
22:6n-3, DHA	14.4 \pm 2.0	13.0 \pm 2.1	11.3 \pm 2.5	13.0 \pm 1.6	11.1 \pm 0.6	11.3 \pm 2.3
Total n-3 PUFA	39.0 \pm 11.0	37.1 \pm 0.7	33.4 \pm 2.9	35.8 \pm 2.1	34.3 \pm 1.9	31.5 \pm 1.4
Total PUFA	48.3 \pm 11.2	45.5 \pm 1.4	42.3 \pm 2.6	46.4 \pm 1.4	46.6 \pm 2.0	45.5 \pm 1.7
EPA/ARA	23.6 \pm 18.2 ^a	12.4 \pm 6.6 ^a	7.8 \pm 4.2 ^{ab}	5.5 \pm 2.5 ^{ab}	3.0 \pm 1.3 ^b	2.4 \pm 0.8 ^b
EPA/DHA	0.9 \pm 0.5	1.3 \pm 0.6	1.5 \pm 0.8	1.3 \pm 0.5	1.3 \pm 0.6	1.4 \pm 0.7
DHA/ARA	23.5 \pm 9.0 ^a	8.8 \pm 1.6 ^b	5.1 \pm 1.1 ^{bc}	4.1 \pm 0.5 ^{bc}	2.2 \pm 0.3 ^{bc}	1.9 \pm 0.4 ^c
n-3/n-6	4.6 \pm 2.3	4.5 \pm 0.7	3.8 \pm 0.8	3.4 \pm 0.4	2.8 \pm 0.3	2.3 \pm 0.3

625
626
627
628
629
630

L: lipids, DW: dry weight, TFA: total fatty acids, ARA: arachidonic acid, DPA: docosapentaenoic acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid.

Table 3. Sequences of PCR primers utilized in this study [45,52]

Transcript	Primer name	Sequence	Amplicon	Tm ⁴	Accession no.
<i>d4fad</i>	Δ4fad -Solea-F8	AAGCCTCTGCTGATTGGAGA	131 bp ³	60	JN673546
	Δ4fad-Solea-R5	GGCTGAGCTTGAAACAGACC			
<i>Elov15</i>	Elov15-Solea-F3	TTTCATGTTTTTGCACACTGC	161 bp	60	JN793448
	Elov15-Solea-R3	GACACCTTTAGGCTCGGTTTT			
<i>UBQ</i> ¹	qUBQ-F	AGCTGGCCCAAAAATATAACTGCGACA	93 bp	70	AB291588
	qUBQ-R	ACTTCTTCTTGCGGCAGTTGACAGCAC			
<i>RPS4</i> ²	qRPS4-F	GTGAAGAAGCTCCTTGTCGGCACCA	83 bp	70	AB291557
	qRPS4-R	AGGGGGTCGGGGTAGCGGATG			

¹UBQ: Ubiquitin

²RPS4: 40S ribosomal protein S4

³bp: base pairs

⁴Tm: annealing temperature.

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655 **Table 4.** Lipid, fatty acid content and fatty acid composition (% TFA \pm SEM) of testis of
 656 *Senegalese sole* fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,
 657 $P < 0.05$, $N = 6$)

	A	B	C	D	E	F
TL (mg g ⁻¹ DW)	78 \pm 7	68 \pm 22	66 \pm 20	81 \pm 3	76 \pm 22	81 \pm 17
TFA (μ g mg ⁻¹ L)	503 \pm 46	533 \pm 52	546 \pm 39	561 \pm 56	538 \pm 34	520 \pm 13
Fatty acid composition (%TFA)						
14:0	0.7 \pm 0.1	0.6 \pm 0.1	0.9 \pm 0.1	0.7 \pm 0.1	0.6 \pm 0.1	0.6 \pm 0.1
16:0	17.8 \pm 0.9	17.8 \pm 0.6	18.9 \pm 0.6	19.8 \pm 0.4	17.7 \pm 1.1	19.0 \pm 0.9
18:0	7.0 \pm 0.5	6.4 \pm 0.4	6.3 \pm 0.8	6.7 \pm 0.7	6.8 \pm 0.6	6.4 \pm 0.3
Total SFA	26.3 \pm 0.8	26.3 \pm 1.7	26.9 \pm 1.2	28.0 \pm 0.8	25.4 \pm 1.7	26.5 \pm 0.9
16:1n-7	3.3 \pm 0.4	2.6 \pm 0.1	3.2 \pm 0.4	3.2 \pm 0.3	2.4 \pm 0.3	3.6 \pm 0.3
18:1n-9	15.8 \pm 1.4	14.2 \pm 1.6	15.1 \pm 1.8	14.9 \pm 0.6	13.0 \pm 1.0	17.3 \pm 1.2
18:1n-7	6.1 \pm 1.7	6.5 \pm 0.6	6.7 \pm 0.7	5.2 \pm 1.5	7.4 \pm 0.6	6.0 \pm 1.6
20:1n-9	1.9 \pm 0.3	1.7 \pm 0.4	1.8 \pm 0.2	1.3 \pm 0.2	1.8 \pm 0.2	1.5 \pm 0.1
22:1n-9	0.5 \pm 0.1	0.2 \pm 0.1	0.2 \pm 0.1	0.4 \pm 0.1	0.3 \pm 0.1	0.3 \pm 0.1
Total MUFA	27.8 \pm 1.8	24.9 \pm 2.5	26.1 \pm 1.1	24.2 \pm 1.3	24.6 \pm 0.9	28.9 \pm 1.1
18:2n-6	8.5 \pm 0.5	8.0 \pm 0.2	6.9 \pm 0.2	8.1 \pm 0.3	7.0 \pm 0.4	7.0 \pm 0.4
20:4n-6, ARA	5.0 \pm 0.5 ^d	7.1 \pm 0.8 ^{cd}	8.6 \pm 1.4 ^c	9.5 \pm 0.3 ^{bc}	12.5 \pm 1.0 ^{ab}	13.4 \pm 0.5 ^a
22:4n-6	0.2 \pm 0.1 ^d	0.4 \pm 0.1 ^c	0.8 \pm 0.1 ^{bc}	0.8 \pm 0.1 ^b	1.1 \pm 0.1 ^{ab}	1.5 \pm 0.1 ^a
22:5n-6, DPA	0.4 \pm 0.1 ^d	0.5 \pm 0.1 ^{cd}	0.6 \pm 0.0 ^{bcd}	0.7 \pm 0.1 ^{abc}	0.9 \pm 0.1 ^{ab}	1.0 \pm 0.1 ^a
Total n-6 PUFA	16.1 \pm 0.5 ^c	16.6 \pm 0.8 ^c	18.8 \pm 1.5 ^{bc}	19.6 \pm 0.5 ^{ab}	22.2 \pm 0.6 ^{ab}	22.3 \pm 0.2 ^a
18:3n-3	0.4 \pm 0.0	0.4 \pm 0.0	0.4 \pm 0.1	0.4 \pm 0.0	0.3 \pm 0.0	0.3 \pm 0.1
18:4n-3	0.2 \pm 0.1	0.1 \pm 0.0	0.2 \pm 0.1	0.2 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0
20:4n-3	1.5 \pm 0.7	0.8 \pm 0.3	0.4 \pm 0.1	0.1 \pm 0.1	0.4 \pm 0.0	0.2 \pm 0.0
20:5n-3, EPA	4.1 \pm 0.5	5.1 \pm 0.6	3.6 \pm 0.2	3.4 \pm 0.8	3.9 \pm 0.4	2.9 \pm 0.3
22:5n-3	3.6 \pm 0.3	4.4 \pm 0.5	3.7 \pm 0.4	3.9 \pm 0.4	3.5 \pm 0.3	2.9 \pm 0.2
22:6n-3, DHA	16.5 \pm 1.2	18.8 \pm 0.5	18.9 \pm 1.7	19.2 \pm 0.5	17.8 \pm 0.5	15.1 \pm 0.8
Total n-3 PUFA	27.4 \pm 1.7 ^a	30.7 \pm 1.6 ^a	27.3 \pm 2.0 ^a	27.4 \pm 0.4 ^a	26.1 \pm 0.6 ^a	20.7 \pm 0.8 ^b
Total PUFA	43.5 \pm 1.5 ^{ab}	47.3 \pm 2.3 ^{ab}	46.1 \pm 0.7 ^{ab}	46.9 \pm 0.6 ^{ab}	48.4 \pm 0.6 ^a	43.0 \pm 0.8 ^b
EPA/ARA	0.8 \pm 0.1 ^a	0.7 \pm 0.1 ^{ab}	0.5 \pm 0.1 ^{bc}	0.4 \pm 0.1 ^c	0.3 \pm 0.1 ^c	0.2 \pm 0.0 ^c
EPA/DHA	0.2 \pm 0.0	0.3 \pm 0.0	0.2 \pm 0.0	0.2 \pm 0.0	0.2 \pm 0.0	0.2 \pm 0.0
DHA/ARA	3.5 \pm 0.5	2.3 \pm 0.6	3.0 \pm 1.3	2.0 \pm 0.1	1.5 \pm 0.1	0.9 \pm 0.2
n-3/n-6	1.7 \pm 0.1 ^a	1.8 \pm 0.1 ^a	1.5 \pm 0.3 ^{ab}	1.4 \pm 0.0 ^{ab}	1.2 \pm 0.1 ^b	0.9 \pm 0.0 ^c

658 TL: Total lipids, L: lipids, DW: dry weight, TFA: total fatty acids, ARA: arachidonic acid, DPA: docosapentaenoic
 659 acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid. Diet A= 0.7, B= 1.6, C= 2.3, D= 3.2, E= 5.0 and F=
 660 6.0% TFA. Data within a row assigned different letters were significantly different (ANOVA, $P < 0.05$, $N = 6$).
 661

662 **Table 5.** Lipid, fatty acid content and fatty acid composition (% TFA \pm SEM) of ovary of
 663 *Senegalese sole* fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,
 664 $P < 0.05$, $N = 6$)

	A	B	C	D	E	F
TL (mg g ⁻¹ DW)	108 \pm 20	100 \pm 22	125 \pm 10	127 \pm 6	131 \pm 9	130 \pm 8
TFA (μ g mg ⁻¹ L)	503 \pm 46	533 \pm 52	546 \pm 39	561 \pm 56	538 \pm 34	520 \pm 13
Fatty acid composition (%TFA)						
14:0	2.0 \pm 0.2	1.8 \pm 0.3	1.7 \pm 0.4	2.0 \pm 0.2	2.0 \pm 0.1	1.9 \pm 0.4
16:0	19.1 \pm 1.0	19.8 \pm 1.5	18.2 \pm 0.9	19.0 \pm 1.0	18.7 \pm 0.7	19.6 \pm 0.5
18:0	4.2 \pm 0.4	4.3 \pm 1.4	4.4 \pm 0.8	3.2 \pm 0.3	3.5 \pm 0.1	4.8 \pm 0.6
Total SFA	25.5 \pm 0.9	28.1 \pm 2.9	25.3 \pm 1.1	24.5 \pm 1.4	24.9 \pm 1.0	27.4 \pm 0.2
16:1n-7	5.3 \pm 0.5	4.8 \pm 0.7	4.6 \pm 0.6	6.2 \pm 0.4	5.0 \pm 0.2	4.6 \pm 0.7
18:1n-9	16.5 \pm 1.1	14.9 \pm 1.0	15.7 \pm 1.3	13.3 \pm 2.8	14.9 \pm 0.8	16.2 \pm 0.8
18:1n-7	3.8 \pm 0.4	3.8 \pm 0.3	4.4 \pm 0.7	3.1 \pm 0.1	3.8 \pm 0.4	3.3 \pm 1.1
20:1n-9	2.9 \pm 0.3	2.5 \pm 0.5	2.6 \pm 0.5	2.9 \pm 0.2	3.1 \pm 0.3	2.6 \pm 0.4
22:1n-9	1.6 \pm 0.3	1.8 \pm 0.2	1.8 \pm 0.6	1.6 \pm 0.2	1.4 \pm 0.5	1.4 \pm 0.4
Total MUFA	30.7 \pm 1.4	26.6 \pm 2.1	28.9 \pm 1.8	25.1 \pm 2.7	28.4 \pm 1.1	27.4 \pm 1.2
18:2n-6	7.8 \pm 0.4	8.4 \pm 0.4	7.1 \pm 0.7	7.9 \pm 0.5	7.1 \pm 0.3	7.5 \pm 0.5
20:4n-6, ARA	2.0 \pm 0.9 ^b	2.9 \pm 1.0 ^b	3.8 \pm 1.3 ^{ab}	4.1 \pm 0.1 ^{ab}	5.1 \pm 0.2 ^{ab}	6.4 \pm 1.6 ^a
22:4n-6	0.3 \pm 0.1	0.4 \pm 0.0	0.5 \pm 0.1	0.5 \pm 0.0	0.5 \pm 0.2	0.7 \pm 0.2
22:5n-6, DPA	0.3 \pm 0.1	0.4 \pm 0.1	0.4 \pm 0.1	0.5 \pm 0.0	0.6 \pm 0.0	0.6 \pm 0.1
Total n-6 PUFA	11.3 \pm 1.3	12.9 \pm 1.1	14.1 \pm 2.0	13.2 \pm 0.8	14.1 \pm 0.5	15.9 \pm 1.8
18:3n-3	0.6 \pm 0.2	0.6 \pm 0.2	0.6 \pm 0.2	1.0 \pm 0.1	0.9 \pm 0.0	0.8 \pm 0.1
18:4n-3	0.6 \pm 0.1	0.9 \pm 0.4	0.5 \pm 0.1	0.8 \pm 0.1	0.8 \pm 0.0	0.6 \pm 0.1
20:4n-3	1.4 \pm 0.7	1.1 \pm 0.5	0.4 \pm 0.1	0.5 \pm 0.1	0.6 \pm 0.0	0.6 \pm 0.1
20:5n-3, EPA	2.8 \pm 0.5	2.0 \pm 0.8	2.9 \pm 0.7	2.8 \pm 0.2	2.3 \pm 0.0	2.7 \pm 0.4
22:5n-3	4.1 \pm 0.4	4.3 \pm 0.6	3.9 \pm 0.3	4.3 \pm 0.4	4.3 \pm 0.1	4.0 \pm 0.3
22:6n-3, DHA	20.5 \pm 1.5	21.2 \pm 3.4	22.8 \pm 2.9	23.3 \pm 1.3	22.7 \pm 0.8	20.0 \pm 0.8
Total n-3 PUFA	31.0 \pm 1.5	30.8 \pm 2.0	31.1 \pm 2.0	35.8 \pm 1.4	31.6 \pm 0.8	28.0 \pm 1.2
Total PUFA	42.4 \pm 1.6	43.6 \pm 1.7	45.2 \pm 1.9	49.0 \pm 1.4	45.7 \pm 0.9	43.9 \pm 1.3
EPA/ARA	1.9 \pm 0.0 ^a	0.9 \pm 0.2 ^b	0.8 \pm 0.1 ^b	0.7 \pm 0.0 ^b	0.4 \pm 0.0 ^b	0.5 \pm 0.0 ^b
EPA/DHA	0.1 \pm 0.1	0.1 \pm 0.1	0.1 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0
DHA/ARA	17.3 \pm 0.2	12.3 \pm 5.1	12.0 \pm 5.8	6.0 \pm 0.2	4.5 \pm 0.3	3.4 \pm 0.9
n-3/n-6	2.9 \pm 0.2	2.5 \pm 0.3	2.5 \pm 0.4	2.8 \pm 0.2	2.3 \pm 0.1	1.9 \pm 0.3

Abbreviations as in Table 4.

665
666

667

668 **Table 6.** Lipid, fatty acid content and fatty acid composition (% TFA \pm SEM) of liver of male
 669 Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,
 670 $P < 0.05$, $N = 6$)

	A	B	C	D	E	F
TL (mg g ⁻¹ DW)	485 \pm 39	472 \pm 41	477 \pm 29	455 \pm 43	457 \pm 36	450 \pm 22
TFA (μ g mg ⁻¹ L)	562 \pm 60	538 \pm 109	555 \pm 59	566 \pm 66	504 \pm 69	539 \pm 50
Fatty acid composition (%TFA)						
14:0	3.9 \pm 0.7	2.8 \pm 0.8	3.8 \pm 0.5	2.2 \pm 1.1	2.9 \pm 0.7	3.4 \pm 0.7
16:0	19.9 \pm 0.4	17.8 \pm 0.8	18.6 \pm 0.5	19.4 \pm 1.2	16.6 \pm 0.9	16.7 \pm 1.5
18:0	3.7 \pm 0.7	3.5 \pm 0.5	2.7 \pm 0.2	2.5 \pm 0.4	2.5 \pm 0.3	3.0 \pm 0.3
Total SFA	28.8 \pm 1.1	24.8 \pm 1.0	25.7 \pm 1.1	24.4 \pm 1.7	22.5 \pm 1.1	23.5 \pm 2.3
16:1n-7	7.3 \pm 0.6	6.5 \pm 1.0	8.8 \pm 0.5	7.4 \pm 0.4	6.2 \pm 0.7	6.1 \pm 1.7
18:1n-9	20.9 \pm 1.3	17.1 \pm 2.2	21.9 \pm 1.3	21.1 \pm 1.4	18.3 \pm 1.0	20.4 \pm 1.2
18:1n-7	3.7 \pm 0.0	3.3 \pm 0.3	3.6 \pm 0.3	3.8 \pm 0.1	3.3 \pm 0.2	3.5 \pm 0.9
20:1n-9	3.5 \pm 0.6	3.2 \pm 0.7	3.7 \pm 0.4	3.8 \pm 0.6	3.8 \pm 0.4	2.7 \pm 0.3
22:1n-9	3.0 \pm 0.4	2.9 \pm 0.4	2.9 \pm 0.5	3.1 \pm 0.8	3.4 \pm 0.6	2.8 \pm 0.4
Total MUFA	29.9 \pm 5.8	30.7 \pm 3.8	40.5 \pm 2.6	45.8 \pm 4.5	33.7 \pm 2.1	33.7 \pm 2.0
18:2n-6	6.7 \pm 0.5	6.7 \pm 1.0	6.4 \pm 0.5	6.8 \pm 0.7	7.9 \pm 0.4	7.5 \pm 0.4
20:4n-6, ARA	0.5 \pm 0.1 ^d	1.2 \pm 0.2 ^c	1.5 \pm 0.1 ^c	1.8 \pm 0.1 ^{bc}	2.4 \pm 0.2 ^{ab}	2.8 \pm 0.4 ^a
22:4n-6	0.1 \pm 0.0 ^c	0.1 \pm 0.1 ^c	0.5 \pm 0.0 ^{bc}	0.5 \pm 0.2 ^{bc}	0.9 \pm 0.1 ^{ab}	1.2 \pm 0.1 ^a
22:5n-6, DPA	0.2 \pm 0.1 ^b	0.3 \pm 0.1 ^b	0.6 \pm 0.1 ^{ab}	0.6 \pm 0.1 ^{ab}	0.9 \pm 0.1 ^{ab}	1.1 \pm 0.1 ^a
Total n-6 PUFA	7.8 \pm 0.6	8.4 \pm 1.1	9.0 \pm 0.6	9.0 \pm 1.1	12.6 \pm 0.6	12.8 \pm 0.8
18:3n-3	0.8 \pm 0.1	0.8 \pm 0.1	0.8 \pm 0.0	0.6 \pm 0.1	0.9 \pm 0.1	0.8 \pm 0.1
18:4n-3	0.4 \pm 0.0	0.5 \pm 0.1	0.3 \pm 0.1	0.3 \pm 0.1	0.5 \pm 0.0	0.3 \pm 0.1
20:4n-3	0.6 \pm 0.1	0.4 \pm 0.1	0.6 \pm 0.0	0.4 \pm 0.1	0.6 \pm 0.1	0.6 \pm 0.1
20:5n-3, EPA	2.7 \pm 0.7	2.2 \pm 0.8	2.7 \pm 0.8	1.7 \pm 0.8	2.3 \pm 0.6	3.2 \pm 0.7
22:5n-3	4.3 \pm 0.4	4.8 \pm 0.8	4.2 \pm 0.7	4.0 \pm 0.8	5.2 \pm 0.2	4.7 \pm 0.4
22:6n-3, DHA	20.2 \pm 3.7	23.8 \pm 4.1	18.8 \pm 1.9	17.3 \pm 2.5	22.0 \pm 2.0	23.0 \pm 4.8
Total n-3 PUFA	29.3 \pm 3.3	33.5 \pm 3.3	24.2 \pm 2.2	24.4 \pm 4.3	30.7 \pm 2.1	29.6 \pm 3.8
Total PUFA	39.4 \pm 3.0	41.7 \pm 3.6	34.8 \pm 2.4	29.9 \pm 2.6	42.5 \pm 1.8	42.5 \pm 3.9
EPA/ARA	7.4 \pm 0.9 ^a	2.6 \pm 0.9 ^b	1.7 \pm 0.7 ^b	1.4 \pm 0.7 ^b	1.3 \pm 0.2 ^b	1.4 \pm 0.1 ^b
EPA/DHA	0.2 \pm 0.1	0.1 \pm 0.1	0.1 \pm 0.1	0.1 \pm 0.1	0.1 \pm 0.0	0.1 \pm 0.1
DHA/ARA	32.9 \pm 2.5 ^a	20.0 \pm 5.6 ^b	7.1 \pm 2.9 ^b	5.8 \pm 2.3 ^b	6.2 \pm 1.7 ^b	6.0 \pm 1.9 ^b
n-3/n-6	3.8 \pm 0.5	3.5 \pm 1.1	2.7 \pm 0.2	2.2 \pm 0.1	2.3 \pm 0.2	2.3 \pm 0.4

Abbreviations as in Table 4.

671
 672
 673
 674

675 **Table 7.** Lipid, fatty acid content and fatty acid composition (% TFA \pm SEM) of liver of female
 676 Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,
 677 $P < 0.05$, $N = 6$)

	A	B	C	D	E	F
TL (mg g ⁻¹ DW)	442 \pm 83	436 \pm 35	419 \pm 27	365 \pm 30	361 \pm 40	406 \pm 33
TFA (μ g mg ⁻¹ L)	482 \pm 94	417 \pm 85	542 \pm 20	534 \pm 27	456 \pm 25	427 \pm 94
Fatty acid composition (%TFA)						
14:0	4.3 \pm 0.6	4.2 \pm 0.7	3.7 \pm 0.7	4.3 \pm 0.7	3.9 \pm 0.6	3.9 \pm 1.0
16:0	21.9 \pm 0.7	19.6 \pm 1.4	18.5 \pm 1.3	21.0 \pm 0.9	21.5 \pm 0.6	19.8 \pm 0.8
18:0	3.7 \pm 1.0	2.8 \pm 0.2	1.9 \pm 0.1	2.1 \pm 0.2	2.9 \pm 0.2	3.2 \pm 0.5
Total SFA	31.1 \pm 1.5	26.9 \pm 2.0	24.2 \pm 1.7	28.3 \pm 1.3	28.8 \pm 1.0	27.0 \pm 2.3
16:1n-7	9.5 \pm 1.7	7.8 \pm 0.6	7.3 \pm 1.6	9.8 \pm 0.4	7.8 \pm 0.4	7.3 \pm 1.3
18:1n-9	18.2 \pm 4.4	19.4 \pm 2.0	21.0 \pm 2.0	21.4 \pm 0.6	22.5 \pm 0.1	20.2 \pm 1.8
18:1n-7	4.2 \pm 0.4	4.1 \pm 0.3	4.7 \pm 0.2	4.7 \pm 0.4	5.1 \pm 0.2	4.9 \pm 0.2
20:1n-9	2.9 \pm 0.5	3.5 \pm 0.8	3.2 \pm 0.6	2.5 \pm 0.2	3.8 \pm 0.4	2.6 \pm 0.5
22:1n-9	2.2 \pm 0.7	2.3 \pm 0.3	2.4 \pm 0.6	2.0 \pm 0.7	2.2 \pm 0.5	2.5 \pm 0.3
Total MUFA	32.2 \pm 5.9	35.2 \pm 3.4	39.8 \pm 2.8	38.8 \pm 1.1	39.7 \pm 0.2	33.7 \pm 3.7
18:2n-6	8.0 \pm 1.1	8.0 \pm 0.1	9.6 \pm 0.5	9.7 \pm 0.6	9.1 \pm 0.2	8.7 \pm 0.6
20:4n-6, ARA	0.6 \pm 0.3 ^c	1.1 \pm 0.2 ^c	1.7 \pm 0.1 ^b	1.8 \pm 0.2 ^b	2.3 \pm 0.1 ^{ab}	3.0 \pm 0.1 ^a
22:4n-6	0.2 \pm 0.1	0.2 \pm 0.1	0.4 \pm 0.0	0.4 \pm 0.0	0.4 \pm 0.1	0.5 \pm 0.1
22:5n-6, DPA	0.3 \pm 0.0	0.3 \pm 0.0	0.5 \pm 0.0	0.5 \pm 0.1	0.5 \pm 0.1	0.6 \pm 0.1
Total n-6 PUFA	9.0 \pm 0.9 ^b	9.6 \pm 0.4 ^b	12.6 \pm 0.4 ^a	12.6 \pm 0.6 ^a	13.9 \pm 0.4 ^a	13.0 \pm 0.6 ^a
18:3n-3	0.9 \pm 0.1	0.9 \pm 0.0	0.8 \pm 0.2	0.9 \pm 0.1	0.9 \pm 0.1	0.8 \pm 0.1
18:4n-3	0.8 \pm 0.2	0.5 \pm 0.0	0.6 \pm 0.2	0.9 \pm 0.1	0.9 \pm 0.2	0.7 \pm 0.1
20:4n-3	0.5 \pm 0.1	0.5 \pm 0.1	0.6 \pm 0.0	0.3 \pm 0.1	0.5 \pm 0.1	0.5 \pm 0.0
20:5n-3, EPA	1.2 \pm 0.3	1.9 \pm 0.4	1.2 \pm 0.2	1.2 \pm 0.2	0.8 \pm 0.3	1.1 \pm 0.1
22:5n-3	2.8 \pm 0.4	3.8 \pm 0.4	3.7 \pm 0.5	3.0 \pm 0.7	3.5 \pm 0.4	3.7 \pm 0.7
22:6n-3, DHA	16.9 \pm 3.9	14.4 \pm 1.4	15.7 \pm 2.3	13.4 \pm 1.9	12.0 \pm 1.3	16.4 \pm 6.0
Total n-3 PUFA	22.6 \pm 4.0	22.9 \pm 1.7	23.1 \pm 3.0	19.5 \pm 1.9	17.2 \pm 0.8	22.3 \pm 5.8
Total PUFA	31.7 \pm 4.6	37.3 \pm 5.4	35.7 \pm 3.3	32.1 \pm 2.3	31.1 \pm 1.2	38.9 \pm 5.9
EPA/ARA	3.3 \pm 1.0 ^a	2.0 \pm 0.6 ^{ab}	0.7 \pm 0.2 ^b	0.6 \pm 0.1 ^b	0.4 \pm 0.1 ^b	0.3 \pm 0.0 ^b
EPA/DHA	0.1 \pm 0.0	0.2 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0	0.1 \pm 0.0
DHA/ARA	45.1 \pm 17 ^a	13.4 \pm 3.6 ^{ab}	9.6 \pm 2.0 ^b	7.5 \pm 1.3 ^b	5.4 \pm 1.1 ^b	4.5 \pm 2.1 ^b
n-3/n-6	2.5 \pm 0.3	2.0 \pm 0.5	1.8 \pm 0.2	1.5 \pm 0.1	1.2 \pm 0.0	0.9 \pm 0.3

678 Abbreviations as in Table 4.
 679
 680
 681