

1 *\*This is an electronic version of an article published in Vol 18 (1) pp 32-43 of*  
2 *Laterality: Asymmetries of Body, Brain and Cognition. Laterality: Asymmetries of*  
3 *Body, Brain and Cognition is available online at:*  
4 *<http://www.tandfonline.com/doi/abs/10.1080/1357650X.2011.626562#.UoJPSpR5NgI>*

5  
6 **Auditory laterality in a nocturnal, fossorial marsupial (*Lasiorhinus latifrons*) in**  
7 **response to bilateral stimuli**

8  
9 Descovich, K.A. <sup>ab</sup>, Reints Bok, T.E. <sup>bc</sup>, Lisle A.T. <sup>a</sup>, Phillips, C.J.C. <sup>b</sup>

10 <sup>a</sup> School of Agriculture and Food Sciences, University of Queensland, Gatton, QLD,  
11 4343, Australia

12 <sup>b</sup> Centre for Animal Welfare and Ethics, University of Queensland, Gatton, QLD, 4343,  
13 Australia

14 <sup>c</sup> Program Emotion and Cognition, Faculty of Veterinary Medicine, Department of Farm  
15 Animal Health, University Utrecht, The Netherlands

16  
17 Behavioural lateralisation is evident across most animal taxa, although few marsupial and  
18 no fossorial species have been studied. Twelve wombats (*Lasiorhinus latifrons*) were  
19 bilaterally presented with eight sounds from different contexts (threat, neutral, food) to  
20 test for auditory laterality. Head turns were recorded prior to and immediately following  
21 sound presentation. Behaviour was recorded for 150 seconds after presentation. Although  
22 sound differentiation was evident by the amount of exploration, vigilance and grooming  
23 performed after different sound types, this did not result in different patterns of head turn  
24 direction. Similarly, left-right proportions of head turns, walking events and food  
25 approaches in the post-sound period were comparable across sound types. A comparison  
26 of head turns performed before and after sound showed a significant change in turn

27 direction ( $\chi^2_1 = 10.65, P = 0.001$ ) from a left preference during the pre-sound period  
28 (mean 58% left head turns, CI 49-66%) to a right preference in the post-sound (mean  
29 43% left head turns, CI 40-45%). This provides evidence of a right auditory bias in  
30 response to the presentation of the sound. This study therefore demonstrates that laterality  
31 is evident in southern hairy-nosed wombats in response to a sound stimulus, although  
32 side biases were not altered by sounds of varying context.

33

#### 34 **Keywords**

35 Wombat; Behaviour; Laterality; Auditory; Marsupial

36

37 Lateralised behaviour is evident in many animal species, including humans (Vallortigara  
38 & Rogers, 2005; Vallortigara, 2006; Corballis, 2007; MacNeilage, Rogers, Vallortigara,  
39 2009), and is apparent in asymmetries of left-right cerebrum use, which manifests as a  
40 side bias (Rogers, 2000). Its expression can be affected by task function, emotion and  
41 perception, social structure, age or gender (de Latude, Demange, Bec, 2009; Pfannkuche,  
42 Bouma, Groothuis, 2009). Laterality assists cognitive processing by reducing reaction  
43 time (Rogers, 2000) and may enhance simultaneous performance of tasks that are  
44 controlled by opposite hemispheres (e.g. feeding and vigilance) (Rogers, Zucca,  
45 Vallortigara, 2004; Ghirlanda, Frasnelli, Vallortigara, 2009). At the population level,  
46 laterality may aid social communication or predator avoidance strategies (Vallortigara,  
47 Chiandetti, Sovrano, 2010).

48

49 Hemispheres appear to control different emotions and therefore the expression of  
50 laterality has the potential to indicate the animal's perception of a stimulus (Rodriguez,  
51 Gomez, Afonso, 1992; Hauser, 1993; Phillips, Llewellyn, Claudia, 2003; de Latude et al.,  
52 2009). A review of lateralization by Rogers (2010) suggests that dominant use of the right

53 hemisphere may indicate an animal that is stressed or has a negative cognitive bias. The  
54 orienting asymmetry paradigm is a non-invasive measure of auditory laterality, using  
55 head turns in response to sounds of varied context (Hauser & Andersson, 1994; Teufel,  
56 Ghazanfar, Fischer, 2010). In dogs (*Canis familiaris*) and Rhesus macaques (*Macaca*  
57 *mulatta*) this test indicated that vocalisations from conspecifics that had been disturbed  
58 by a stranger knocking at the door or isolated in a room were processed with the right ear  
59 / left hemisphere (Hauser & Andersson, 1994; Siniscalchi, Quaranta, Rogers, 2008).  
60 These stimuli did not represent an immediate and severe threat to the dogs. By contrast,  
61 the left ear / right hemisphere was dominant in dogs for thunderstorm sounds and in  
62 Rhesus macaques for heterospecific vocalizations”

63

64 Few marsupial species have been tested for laterality despite cerebral differences from  
65 eutherian mammals (Lippolis, Westman, McAllan, Rogers, 2005). Similar to birds,  
66 marsupials lack a corpus callosum, which bridges the hemispheres and may play a role in  
67 lateralisation (Wiltschko, Traudt, Güntürkün, Prior, Wiltschko, 2002; Josse, Mohamed,  
68 Kherif, Price, 2008), although they do possess an anterior commissure, which  
69 interconnects the auditory fields of the hemispheres (Heath & Jones, 1971; Aitken, 1995).  
70 Unlike marsupials, lateralisation in birds has been extensively studied and the patterns of  
71 hemispheric use are similar to mammals (Rogers, 2008). The wombat presents a useful  
72 marsupial model for such studies due to its laterally placed eyes (Sanderson & Pearson,  
73 1981), largely immobile ears, and its nocturnal, semi-fossorial behaviour. This study  
74 aimed to determine whether wombats are lateralised in their response to auditory stimuli,  
75 and whether this is influenced by sound type and context.

76


77

77 **Methods**

78 This study used twelve adult southern hairy-nosed wombats (*Lasiorhinus latifrons*)  
79 maintained in four groups of one male with two females. Eleven of these were collected  
80 from the wild 5 - 8 years prior to this study and one was born at the facility in 2003. Each  
81 group was provided with a temperature-controlled burrow system, digging chamber,  
82 feeding house, log and an outside enclosure (76 m<sup>2</sup> - 249 m<sup>2</sup>) with native grasses. Carrots,  
83 hay and macropod pellets were provided daily. The wombats were checked during  
84 routine cleaning and weighed fortnightly. Ethical clearance was obtained from the  
85 University of Queensland, Animal Ethics Committee (licence number: SAS/402/09).

86

87 Audio testing occurred in a vacant den (Figure 1) identical to those used by the wombats.  
88 A wooden and wire mesh feeding frame was placed centrally to control the wombat's  
89 position, and two speakers (Logitech, LS11, Dick Smith Electronics) were attached  
90 bilaterally (azimuth = 90° and 270°) to the den roof. Two infrared cameras (Sony IR  
91 Outdoor Night Vision CCIR 507L28) and surveillance software (Skyview Super-series,  
92 Skyview) were used to record behaviour at two frames per second in black and white.  
93 Wombats were habituated to the equipment to avoid confounding results from neophobia  
94 (Robins & Phillips, 2009).



95

96 *Figure 1: Two-dimensional representation of the test den.*

97


98 *Treatment sounds*

99 Eight sound clips were used from three different contexts – threat, neutral and food-  
100 conditioned. Wombat hissing, and vocalisations from two predators (dog and dingo) were  
101 used as threatening sounds. Two sounds believed to be neutral (air-conditioning and  
102 aeroplane), were chosen because they were heard frequently at the test site without  
103 evoking behavioural responses from the wombats. Three novel and biologically irrelevant  
104 sounds (bells, opera and whistle) were positively conditioned to a food reward by  
105 offering a preferred treat immediately after presenting the sound. This was done six times  
106 per day for five days prior to testing.

107

108 All sound files were monophonic and five seconds in length, with a median volume of 62  
109 - 67.5 dB SPL (Digital Sound Level Meter, Q1362, Dick Smith Electronics). The sound  
110 pressure level produced by each speaker was identical (mean difference in minimum level  
111 across 8 sounds was 0 dB SPL and maximum level was 0.5 dB SPL). Frequencies were

112 analysed using Raven Pro (version 1.3) spectral analysis software (Figure 2).  
113 Spectrogram frequency outputs from the two speakers were compared and found to be  
114 less variable than output from the same speaker played two times (mean cross  
115 correlations for eight sounds was 0.958 for within speaker variation, compared with 0.971  
116 for between speakers).  
117



118  
119 *Figure 2: Spectrograms (Raven Pro, version 1.3) of eight test sounds: a) Air-*  
120 *conditioning, b) Bells, c) Dingo, d) Dog, e) Opera, f) Plane, g) Whistle, h) Wombat.*  
121 *Sounds a, b, g and h were recorded on-site using a digital audio recorder (Joybee 110,*  
122 *BenQ). Sounds c - f were sourced from soundboard.com.*

123  
124 Testing began at 0600 h, and was carried out in three blocks of 9, 9 and 6 days, with four  
125 wombats tested individually and in a random order each day. Testing lasted between 566  
126 - 3612 seconds, and all wombats had an inter-test rest period of 3 days.

127

128 Behaviour was recorded over three periods: baseline (head turns prior to sound  
129 presentation), response (head turns within 30 seconds of sound presentation) and post-  
130 sound (all behaviour within 150 seconds of sound presentation). One trained observer  
131 scored all of the video data, and a second trained observer scored 5 % of the videos to  
132 check for inter-rater reliability (94.9 %). Behaviour was categorised into head turns  
133 (vigilance), feeding, locomotion, escape, exploration, resting and grooming. Food was  
134 placed into the bowl using a reaching tool (Nifty Nabber, Crafright) through a hole in the  
135 roof that also allowed discrete viewing of the den. The sound was presented once the  
136 animal was in the feeding frame with its head straight and of equal distance between the  
137 two speakers. Responses were considered invalid if the sound played when these criteria  
138 were not met. The procedure was repeated twice more using the same sound. Wombats  
139 that did not approach the bowl within three minutes (16 % occurrence) were lured to it  
140 using the reaching tool. On two occasions luring did not work within 20 min and the  
141 wombat was released back into its enclosure and re-tested at the end of the session.

142

#### 143 *Statistical analysis*

##### 144 *Head turn in response to sound*

145 The direction of responsive head turns was analysed for the effect of sound type using  
146 Chi-square tests of associations (Freq procedure, SAS®, version 8.2) for left and right  
147 turns, and no response. Head turn preferences for individuals and after each sound type  
148 were calculated using an Exact Binomial Test, using only responses in which a left or  
149 right choice had been made.

150

##### 151 *Behaviour post-sound*

152 Post-sound behaviour (0 – 150 seconds) was categorised into feeding, locomotion,  
153 vigilance, escape, exploratory, resting and grooming. Data were not normally distributed


154 by univariate analysis, therefore behavioural frequencies were analysed using a Genmod  
155 procedure with a Poisson distribution in SAS to determine the influence of sound type.  
156 Resting was too infrequent to include. If an overall effect of sound was indicated, post-  
157 hoc Chi-square tests were conducted to test for differences in behaviour between sounds.  
158 Three post-sound behaviours were directional: walking (clockwise/anti-clockwise), head  
159 turns (left/right) and food approach (bowl on left/right). These were analysed using the  
160 Genmod procedure in SAS with a binomial distribution to determine whether sound type  
161 influenced direction. Pre- and post-sound data were compared using the Genmod  
162 procedure with a binomial distribution in SAS.

163

## 164 **Results**

165 The direction of responsive head turns was not influenced by sound type ( $\chi^2_{14} = 14.24$ ,  $P$   
166  $= 0.43$ ), although one sound (air-conditioning) was significantly skewed to the right  
167 (Figure 3). Only one individual had a significant directional preference (Figure 3). The  
168 direction of post-sound walking, head turning or food approach was similar between  
169 sounds (Table 1). Head turn direction significantly changed ( $\chi^2_1 = 10.65$ ,  $P = 0.001$ ) from  
170 a left preference [proportion left (with 95% CI) = 0.58 (0.49 – 0.66)] in the baseline  
171 period to a right preference in the post-sound period [proportion left (with 95% CI) =  
172 0.43 (0.40 – 0.45)]. This change was not affected by the type of sound presented ( $\chi^2_7 =$   
173 6.77,  $P = 0.45$ ) (Table 2).

174



175

176 *Figure 3. Mean head turn preferences (± SE) for individual wombats and sounds.*

177 *Negative and positive means indicate a left and right preference respectively. Preferences*

178 *were calculated by attributing a left response with -1, right response with +1 and no*

179 *response with 0 then calculating means for each individual or sound. \* indicates a*

180 *significant ( $p < 0.05$ ) preference using an Exact Binomial Test.*

181

182

182 **Table 1. Back-transformed proportions (with 95% CI) of left or anti-clockwise**  
 183 **behaviour after 8 sound types, and the overall sound effect on direction ( $\chi^2$  statistic,**  
 184 **degrees of freedom, probability value).**


Sound type	Head movement left	Approach food left	Walking anti-clockwise
Air-con.	0.40 (0.32 – 0.47)	0.55 (0.28 – 0.79)	0.65 (0.56 – 0.72)
Bells	0.40 (0.33 – 0.48)	0.52 (0.27 – 0.76)	0.60 (0.51 – 0.67)
Dingo	0.46 (0.39 – 0.53)	0.66 (0.43 – 0.83)	0.57 (0.49 – 0.64)
Dog	0.41 (0.34 – 0.49)	0.61 (0.35 – 0.82)	0.66 (0.58 – 0.73)
Opera	0.45 (0.38 – 0.52)	0.46 (0.22 – 0.73)	0.61 (0.52 – 0.69)
Plane	0.39 (0.32 – 0.46)	0.49 (0.25 – 0.73)	0.66 (0.58 – 0.73)
Whistle	0.44 (0.37 – 0.52)	0.39 (0.18 – 0.65)	0.71 (0.63 – 0.78)
Wombat	0.47 (0.40 – 0.54)	0.64 (0.41 – 0.82)	0.68 (0.61 – 0.75)
Sound effect	$\chi^2_7 = 5.37, p = 0.62$	$\chi^2_7 = 3.81, p = 0.80$	$\chi^2_7 = 10.61, p = 0.16$

185

186 **Table 2. Back-transformed proportions (with 95% CI) of left head turns after 8**  
 187 **sound types.**

Sound type	Proportion left pre-sound	Proportion left post-sound
Air-con.	0.47 (0.27 – 0.69)	0.40 (0.32 – 0.47)
Bells	0.72 (0.43 – 0.89)	0.41 (0.33 – 0.49)
Dingo	0.55 (0.31 – 0.78)	0.46 (0.39 – 0.53)
Dog	0.37 (0.19 – 0.61)	0.41 (0.34 – 0.49)
Opera	0.60 (0.46 – 0.73)	0.45 (0.38 – 0.52)
Plane	0.74 (0.44 – 0.92)	0.38 (0.32 – 0.46)
Whistle	0.51 (0.33 – 0.68)	0.44 (0.37 – 0.52)
Wombat	0.60 (0.35 – 0.81)	0.47 (0.40 – 0.81)

188 Sound type significantly affected the expression of three behaviours in the post-sound  
 189 period: exploration ( $\chi^2_7 = 16.8, p = 0.02$ ), vigilance ( $\chi^2_7 = 26.2, p = 0.0005$ ) and grooming  
 190 behaviour ( $\chi^2_7 = 24.1, p = 0.001$ ) (Figure 4). Vigilance behaviour was exhibited most  
 191 frequently after the plane and wombat sounds and least frequently for bells and whistles.  
 192 Exploratory behaviour occurred most commonly after hearing bells, while the whistle  
 193 elicited the most grooming and the wombat the least.  
 194



195  
 196 *Figure 4. Mean counts (± 95% CI) of vigilance, exploration and grooming after the*  
 197 *presentation of eight sound types. Means with different letters are significantly different*  
 198 *( $p < 0.05$ ).*

199 **Discussion**

200 It is evident that the wombats could distinguish between different sounds, as the amount  
201 of vigilance, exploration and grooming was significantly affected by sound type.

202 Vigilance was performed most frequently after plane and wombat sounds and least after  
203 whistles and bells. Wombat hissing represents a threat, while the whistle and bells were  
204 conditioned to food and therefore these results are mostly unsurprising. The plane sound,  
205 however, was believed to be neutral due to its frequency in the test vicinity and usual lack  
206 of response by the wombats. The higher amount of vigilance shown after this sound  
207 suggests that threat perception was influenced by context (Thorson, Morgan, Brown,  
208 Norman, 1998), and habituated sounds encouraged alertness when they were presented in  
209 a novel setting. Grooming was infrequent after all agonistic sounds (wombat, dog, dingo),  
210 which is probably because threatened animals direct less energy towards maintenance  
211 behaviour such as scratching (Hirsch, 2002; Stojan-Dolar & Heymann, 2010).

212 Differences in exploration and grooming were apparent between food-conditioned  
213 sounds, indicating that conditioned responses may be affected by sound type. Despite  
214 sound differentiation, wombats did not demonstrate lateralised behavioural responses to  
215 sounds of varying context. No side preferences were evident for responsive head turns or  
216 post-sound directional behaviour, a result that concurs with Fischer et al.'s (2009)  
217 orienting study in humans.

218

219 Exposure to sound of any type significantly changed the direction of head turns from a  
220 left bias in the baseline period to a right bias after sound presentation. This may be due to  
221 a right auditory bias, as seen in humans (Tallus, Hugdahl, Alho, Medvedev, Hämäläinen,  
222 2007; Devlin, Raley, Tunbridge, Lanary, Floyer-Lea, et al. 2003), which becomes  
223 strengthened after sound presentation due to increased focus on that side (Tallus et al.,  
224 2007). Alternatively the change may result from habituation to the test situation with the

225 wombats displaying vigilance (left side) on first entering the den, and then switching to  
226 the right as they become familiar with the environment. This would concur with previous  
227 studies that show most species (60 – 95%) exhibit a right hemisphere / left side  
228 preference for vigilance (Vallortigara & Rogers, 2005), while familiar objects are usually  
229 processed by the left hemisphere / right side (Robins & Phillips, 2009). Speaker  
230 differences may also have influenced the change in head turn, as speaker positions were  
231 not randomised. However, this is unlikely as turn direction changed significantly while  
232 measured differences between speakers were minimal, and discrepancies would need to  
233 be consistently in the same direction to have caused this change.

234

235 This study concludes that the southern hairy-nosed wombats exhibited lateralised  
236 behaviour in response to sound presentation, although the expression was unaffected by  
237 different types of auditory stimuli. Further studies using this species are recommended to  
238 clarify the functional drivers of this hemispheric specialisation in marsupials.

239

#### 240 **Acknowledgments**

241 The authors would like to acknowledge the Wombat Foundation for financial support,  
242 and the Rockhampton Regional Council for allowing us to use the facilities, animals and  
243 accommodation at the Rockhampton Botanic Gardens and Zoo. Technical support and  
244 advice was kindly provided by Dr Alan Horsup from the QLD Department of  
245 Environmental and Resource Management, Dr Andrew Robins, Dr Wayne Wilson, and  
246 Dr Franz Josef van der Staay.

247

247 **References**

248 Aitkin, L. (1995). The auditory neurobiology of marsupials: A review. *Hearing Research*,  
249 82, 257-266.

250

251 Corballis, M. C. (2007). Cerebral asymmetry and human uniqueness. In W. D. Hopkins  
252 (Ed.), *The evolution of hemispheric specialization in primates: Special topics in*  
253 *primatology* (Vol. 5). Oxford: Academic Press.

254

255 de Latude, M., Demange, M., & Bec, P. (2009). Visual laterality responses to different  
256 emotive stimuli by red-capped mangabeys, *Cercocebus torquatus torquatus*. *Animal*  
257 *Cognition*, 12, 31-42.

258

259 Devlin, J., Raley, J., Tunbridge, E., Lanary, K., Floyer-Lea, A., Narain, C., et al. (2003).  
260 Functional Asymmetry for Auditory Processing in Human Primary Auditory Cortex. *The*  
261 *Journal of Neuroscience*, 23(37), 11516-11522.

262

263 Fischer, J., Teufel, C., Drolet, M., Patzelt, A., Rusamen, R., von Cramon, D. Y., et al.  
264 (2009). Orienting asymmetries and lateralised processing of sounds in humans. *BMC*  
265 *Neuroscience*, 10(14).

266

267 Ghirlanda, S., Frasnelli, E., & Vallortigara, G. (2009). Intraspecific competition and  
268 coordination in the evolution of lateralization. *Philosophical transactions of the Royal*  
269 *Society of London B.*, 364, 861-866.

270

271 Hauser, M. (1993). Right hemisphere dominance for the production of facial expression  
272 in monkeys. *Science*, 261, 475-477.

273

274 Hauser, M. D., & Andersson, K. (1994). Left-hemisphere dominance for processing  
275 vocalisations in adult, but not infant, rhesus monkeys: Field experiments. *Proceedings of*  
276 *the National Academy of Sciences, USA*, 91(9), 3946-3948.

277

278 Heath, C. J., & Jones, E. G. (1971). Interhemispheric pathways in the absence of a corpus  
279 callosum: An experimental study of the commissural connexions in the marsupial  
280 phalanger. *Journal of Anatomy*, 109, 253-270.

281

282 Hirsch, B. T. (2002). Social monitoring and vigilance behaviour in brown capuchin  
283 monkeys (*Cebus apella*). *Behavioral Ecology and Sociobiology*, 52, 458-464.

284

285 Hopkins, W., & Bennett, A. (1994). Handedness and approach-avoidance behaviour in  
286 chimpanzees (*Pan*). *Journal of Experimental Psychology: Behavioural Processes*, 20,  
287 413-418.

288 Josse, G., Mohamed, L. S., Kherif, F., & Price, C. J. (2008). Explaining function with  
289 anatomy: Language lateralization and corpus callosum size. *The Journal of Neuroscience*,  
290 28(52), 14132-14139.

291 Lippolis, G., Westman, W., McAllan, B., & Rogers, L. (2005). Lateralisation of escape  
292 responses in the stripe-faced dunnart, *Sminthopsis macroura* (Dasyuridae: Marsupialia).  
293 *Laterality: Asymmetries of Body, Brain and Cognition*, 10(5), 457-470.

294 MacNeilage, P. F., Rogers, L., & Vallortigara, G. (2009). Origins of the left and right  
295 brain. *Scientific American*, 301(1), 60-67.

296 Pfannkuche, K. A., Bouma, A., & Groothuis, T. G. G. (2009). Does testosterone affect  
297 lateralization of brain and behaviour? A meta-analysis in humans and other animal  
298 species. *Philosophical transactions of the Royal Society of London B.*, 364, 929-942.

299 Phillips, C., Llewellyn, S., & Claudia, A. (2003). Laterality in Bovine Behavior in an  
300 Extensive Partially Suckled Herd and an Intensive Dairy Herd. *Journal of Dairy Science*,  
301 86, 3167-3173.

302 Robins, A., & Phillips, C. (2009). Lateralised visual processing in domestic cattle herds  
303 responding to novel and familiar stimuli. *Laterality: Asymmetries of Body, Brain and*  
304 *Cognition*.

305 Rockhampton Regional Council. (2011). *Rockhampton Airport*. Retrieved 24 February  
306 2011, from [www.rockhamptonairport.com.au](http://www.rockhamptonairport.com.au)

307 Rodriguez, M., Gomez, J. A., & Afonso, D. (1992). Laterality, alternation, and  
308 perserveration relationships on the T-maze test. *Behavioral Neuroscience*, 106(6), 974-  
309 980.

310 Rogers, L. (2000). Evolution of Hemispheric Specialization: Advantages and  
311 Disadvantages. *Brain and Language*, 73, 236-253.

312 Rogers, L. (2008). Development and function of lateralization in the avian brain. *Brain*  
313 *Research Bulletin*, 76, 235-244.

314 Rogers, L. (2010). Relevance of brain and behavioural lateralization to animal welfare.  
315 *Applied Animal Behaviour Science*, 127, 1-11.

316 Rogers, L., Zucca, P., & Vallortigara, G. (2004). Advantage of having a lateralized brain.

- 317 *Proceedings of the Royal Society of London B*, 271, S420-S422.
- 318 Sanderson, K. J., & Pearson, L. J. (1981). Retinal projections in the Hairy-nosed  
319 Wombat, *Lasiorhinus latifrons* (Marsupialia : Vombatidae). *Australian Journal of*  
320 *Zoology*, 29, 473-481.
- 321 Siniscalchi, M., Quaranta, A., & Rogers, L. (2008). Hemispheric Specialization in Dogs  
322 for Processing Different Acoustic Stimuli. *PLoS One*, 3(10), e3349.
- 323 Stojan-Dolar, M., & Heymann, E. W. (2010). Vigilance in a cooperatively breeding  
324 primate. *International Journal of Primatology*, 31, 95-116.
- 325 Tallus, J., Hugdahl, K., Alho, K., Medvedev, S., & Hämäläinen, H. (2007). Interaural  
326 intensity difference and ear advantage in listening to dichotic consonant-vowel syllable  
327 pairs. *Brain Research* 1185, 195-200.
- 328 Teufel, C., Ghazanfar, A. A., & Fischer, J. (2010). On the relationship between  
329 lateralized brain function and orienting asymmetries. *Behavioral Neuroscience*, 124, 437-  
330 445.
- 331
- 332 Thorson, J. M., Morgan, R. A., Brown, J. S., & Norman, J. E. (1998). Direct and indirect  
333 cues of predatory risk and patch use by fox squirrels and thirteen-lined ground squirrels.  
334 *Behavioral Ecology*, 9(2), 151-157.
- 335
- 336 Vallortigara, G. (2006). The evolutionary psychology of left and right: Costs and benefits  
337 of lateralization. *Developmental Psychobiology*, 48(6), 418-427.
- 338 Vallortigara, G., Chiandetti, C., & Sovrano, V. A. (2010). Brain asymmetry (animal).

339 *Wiley Interdisciplinary Reviews: Cognitive Science.*

340 Vallortigara, G., & Rogers, L. (2005). Survival with an asymmetrical brain: Advantages  
341 and disadvantages of cerebral lateralization. *Behavioral and Brain Sciences*, 28, 575-633.

342 Wiltschko, W., Traudt, J., Güntürkün, O., Prior, H., & Wiltschko, R. (2002).

343 Lateralization of magnetic compass orientation in a migratory bird. *Nature*, 419, 467-469.

344