

Accepted refereed manuscript of:

Xie D, Chen F, Lin S, You C, Wang S, Zhang Q, Monroig O, Tocher DR & Li Y (2016) Long-chain polyunsaturated fatty acid biosynthesis in the euryhaline herbivorous teleost *Scatophagus argus*: Functional characterization, tissue expression and nutritional regulation of two fatty acyl elongases, *Comparative Biochemistry and Physiology - Part B: Biochemistry and Molecular Biology*, 198, pp. 37-45.

DOI: [10.1016/j.cbpb.2016.03.009](https://doi.org/10.1016/j.cbpb.2016.03.009)

© 2016, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

1           **Long-chain polyunsaturated fatty acid biosynthesis in the euryhaline**  
2 **herbivorous teleost *Scatophagus argus*: Functional characterization, tissue**  
3 **expression and nutritional regulation of two fatty acyl elongases**

4           Dizhi Xie<sup>1,2,a</sup>, Fang Chen<sup>1,2,a</sup>, Siyuan Lin<sup>1</sup>, Cuihong You<sup>1</sup>, Shuqi Wang<sup>1</sup>, Qinghao Zhang<sup>1</sup>,  
5                           Óscar Monroig<sup>3</sup>, Douglas R. Tocher<sup>3</sup>, Yuanyou Li<sup>1\*</sup>

6  
7           <sup>1</sup> Marine Biology Institute & Guangdong Provincial Key Laboratory of Marine Biotechnology,  
8           Shantou University, Shantou, Guangdong 515063, PR China

9           <sup>2</sup> Fisheries College, Henan Normal University, Xinxiang 453007, P. R. China

10          <sup>3</sup> Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9 4LA,  
11          Scotland, UK

12  
13  
14  
15  
16  
17  
18  
19          \*Correspondence to: Prof. Yuanyou Li, Ph.D.  
20                           Marine Biology Institute  
21                           Shantou University  
22                           Shantou, Guangdong 515063, China  
23                           Tel: +86-754-86503157  
24                           Fax: +86-754-86500614  
25                           E-mail: yyli@stu.edu.cn

26          <sup>a</sup> Joint first authorship.

29 **Abstract:**

30 Both the spotted scat *Scatophagus argus* and rabbitfish *Siganus canaliculatus* belong to the few  
31 cultured herbivorous marine teleost, however, their fatty acyl desaturase (Fad) system involved in  
32 long-chain polyunsaturated fatty acid (LC-PUFA) biosynthesis are different. The *S. argus* has a  $\Delta 6$   
33 Fad, while the rabbitfish has  $\Delta 4$  and  $\Delta 6/\Delta 5$  Fad, which were the first report in vertebrate and  
34 marine teleost, respectively. In order to compare the characteristics of elongases of very long-chain  
35 fatty acids (Elovl) between them, two Elovl cDNAs were cloned from *S. argus* in the present study.  
36 One has 885 bp of open read fragment (ORF) encoding a protein with 294 amino acid (aa) showing  
37 Elovl5 activity functionally characterized by heterologous expression in yeast, which was primarily  
38 active for the elongation of C18 and C20 PUFA. The other has 915 bp of ORF coding for a 305 aa  
39 protein showing Elovl4 activity, which was more efficient in the elongation of C20 and C22 PUFA.  
40 Tissue distribution analyses by RT-PCR showed that *elovl5* was highly expressed in liver compared  
41 to other tissues determined, whereas *elovl4* transcripts were only detected in eye. The expression of  
42 *elovl5* and *elovl4* were significantly affected by dietary fatty acid composition, with highest  
43 expression of mRNA in liver and eye of fish fed a diet with an 18:3n-3/18:2n-6 ratio of 1.7:1. These  
44 results indicated that the *S. argus* has a similar Elovl system in the LC-PUFA biosynthetic pathway  
45 to that of rabbitfish although their Fad system was different, suggesting that the diversification of  
46 fish LC-PUFA biosynthesis specificities is more associated with its Fad system. These new insights  
47 expand our knowledge and understanding of the molecular basis and regulation of LC-PUFA  
48 biosynthesis in fish.

49 **Key words:** Elovl5, Elovl4, LC-PUFA biosynthesis, *Scatophagus argus*

50

51

## 52 **Introduction**

53 The long-chain (C20-24) polyunsaturated fatty acids (LC-PUFA), particularly  
54 eicosapentaenoic (EPA; 20:5n-3) and docosahexaenoic (DHA; 22:6n-3) acids found primarily in  
55 fish and seafood, are regarded as beneficial in a series of human pathologies including metabolic  
56 disorders, cardiovascular, inflammatory and neurological diseases (Muhlhausler & Ailhaud, 2013;  
57 Delgado-Lista et al., 2012; Awada et al., 2013; Campoy et al., 2012). Comparing to freshwater and  
58 salmonids species, marine fish are generally thought to have limited capability or inability for *de*  
59 *novo* LC-PUFA biosynthesis (Sargent et al., 2002; Tocher, 2010). As marine fish are a major source  
60 of n-3 LC-PUFA for humans, the biosynthesis and metabolic regulation of these key nutrients have  
61 become areas of considerable research in recent years (Tocher, 2010; Xie et al., 2014; Zhang et al.,  
62 2014; Kabeya et al., 2015).

63 The biosynthesis of LC-PUFA from the C18 precursors  $\alpha$ -linolenic acid (LNA, 18:3n-3) and  
64 linoleic acid (LA, 18:2n-6) consists of sequential reactions catalyzed by a series of fatty acyl  
65 desaturase (Fads) and elongase of very long-chain fatty acid (Elovl) enzymes such as  $\Delta$ 6 Fad,  $\Delta$ 5  
66 Fad,  $\Delta$ 6/ $\Delta$ 5 Fad,  $\Delta$ 6/ $\Delta$ 8 Fad,  $\Delta$ 4 Fad, Elovl5, Elovl4 and Elovl2 (Torstensen & Tocher, 2010; Li et  
67 al., 2010; Monroig et al., 2011a; Fonseca-Madrugal et al., 2014; Castro et al., 2016). Differences or  
68 absence in the activity of enzymes in one or more steps of the pathways result in differential  
69 LC-PUFA biosynthetic capability in fish (Castro et al., 2016). Currently, existing data show that the  
70 capability for LC-PUFA biosynthesis in marine fish is more diverse than that in other vertebrates  
71 (Fonseca-Madrugal et al., 2014) and the diversity was primarily associated with differences in the  
72 compliment of enzymes/activities in the pathway of LC-PUFA biosynthesis. Among marine fish,  $\Delta$ 6  
73 Fad and Elovl5 cDNAs have been identified in more than a dozen species (Monroig et al., 2011b;  
74 Castro et al., 2016). However, Elovl4 has been investigated to a lesser extent but was reported in  
75 cobia (*Rachycentron canadum*) (Monroig et al., 2011c), rabbitfish (*Siganus canaliculatus*)  
76 (Monroig et al., 2012), Nibe croaker (*Nibea mitsukurii*) (Kabeya et al., 2015) and orange-spotted  
77 grouper (*Epinephelus coioides*) (Li et al., 2015).  $\Delta$ 4 Fad has been identified in two marine fish,  
78 rabbitfish and Senegalese sole (*Solea senegalensis*, 1858) (Li et al., 2010; Morais et al., 2012),  
79 while  $\Delta$ 6/ $\Delta$ 5 Fad was found only in rabbitfish (Li et al., 2010) among the marine teleost. Thus, to  
80 date, rabbitfish is the only marine teleost in which Fad and Elovl enzymes that possess all the  
81 activities required for the production of LC-PUFA from C18 PUFA have been found.

82 With respect to the differences in the capability for LC-PUFA biosynthesis among marine fish,  
83 Castro et al. (2012) hypothesized that the losses and diversifications of crucially important genes in  
84 the LC-PUFA biosynthetic pathway during fish evolution might be linked to habitat-specific food  
85 web characteristics, such as LC-PUFA availability, in different environments. More recently, other  
86 confounding factors including “trophic ecology” and diadromy have been proposed (Morais et al.,  
87 2012; Monroig et al., 2013). Herbivorous rabbitfish has a wide distribution in the coral reefs of the  
88 Indo-Pacific region (Woodland, 1983) and can also live in brackish water (Li et al., 2008), and  
89 feeds on a range of macroalgae including *Enteromorpha prolifera* and *Gracilaria lemaneiformis*  
90 (You et al., 2014).

91 Both the spotted scat (*Scatophagus argus*) and rabbitfish are economically important cultured  
92 teleost. The spotted scat has similar habitat (euryhaline) and feeding (herbivore) habits to rabbitfish,  
93 and is distributed widely in freshwater, brackish and marine habitats of the Indo-Pacific, South and  
94 South East Asia (Barry & Fast, 1992; Gandhi, 2002; Yoshimura et al., 2003). In order to know  
95 whether *S. argus* has an enzymatic complement for LC-PUFA biosynthesis similar to that of  
96 rabbitfish, we aimed to clone and functionally characterize all the genes involved in LC-PUFA  
97 biosynthesis in this species. Our previous study showed that the Fads2 of *S. argus* was a  
98 monofunctional  $\Delta 6$  desaturase enzyme ( $\Delta 6$  Fad), which is in contrast to the more diverse enzymatic  
99 complement ( $\Delta 4$  Fad and  $\Delta 6/\Delta 5$  Fad) found in rabbitfish (Li et al., 2010), suggesting that the above  
100 mentioned diversification also exists within marine herbivorous fish (Xie et al., 2014). Besides, two  
101 elongases including Elov14 and Elov15 were identified in rabbitfish (Oscar et al., 2012). In order to  
102 compare the characteristics of Elov1 system between spotted scat and rabbitfish, and provide basis  
103 for fully understanding the LC-PUFA biosynthetic capability of *S. argus*, the present study reports  
104 the cloning, functional characterization, tissue expression and nutritional regulation of two Elov1  
105 cDNAs encoding putative Elov14 and Elov15, key enzymes with well-demonstrated roles in  
106 LC-PUFA biosynthesis in fish (Castro et al., 2016).

107

## 108 **Materials and Methods**

### 109 ***Experimental fish and sampling***

110 Juvenile *S. argus* (body mass around 4.3 g) were purchased from a commercial hatchery in  
111 Zhuhai (Guangdong, China). Six isoproteic and iso-lipidic experimental diets (D1-D6) were

112 formulated with 32 % crude protein and 8 % crude lipid (soybean oil, perilla oil or fish oil as lipid  
113 source). Diet D2 contained fish oil (FO) as control, and diets D1, D3-D6 contained different  
114 proportions of soybean oil and perilla oil, which resulted in LNA: LA ratios of 0.14, 0.57, 0.84, 1.72  
115 and 2.85, respectively. The detailed dietary formulations, proximate and fatty acid compositions  
116 were shown in [Xie et al. \(2014\)](#).

117 All juvenile *S. argus* were reared in floating cages (0.6×0.6×3.0 m) located on the coast near  
118 Nan'ao Marine Biological Station (NAMBS), Shantou University, and fed an equal mix of the six  
119 experimental diets for two weeks before the start of feeding trial. The feeding trial was conducted in  
120 18 cages at ambient temperature, salinity and photoperiod, with each cage containing 25 fish that  
121 were allocated randomly. Fish in triplicate cages were fed one of the experimental diets twice a day  
122 (at 9:00 and 16:00 h) for 8 weeks. At the end of the feeding trial, fish were anaesthetized with  
123 0.01% 2-phenoxyethanol (Sigma-Aldrich Inc., USA), and livers of 54 fish (3 fish per replicate cage)  
124 were collected, frozen in liquid nitrogen and stored at -80 °C prior to the analysis of *elovl* mRNA  
125 expression by quantitative PCR (qPCR). In order to determine the tissue distribution of *elovl5* and  
126 *elovl4* transcripts, eye, brain, liver, muscle, heart, gills, spleen, kidney and intestine were collected  
127 from wild *S. argus* (50–60 g) captured from the coast near NAMBS, after fish were anaesthetized  
128 with 0.01% 2-phenoxyethanol. Tissue samples were frozen in liquid nitrogen immediately after  
129 collection and stored at -80 °C until RNA extraction.

### 130 ***Molecular cloning of elovl5 and elovl4 cDNAs***

131 Total RNA was extracted from *S. argus* liver and eye using Trizol reagent (Invitrogen, USA)  
132 and reverse transcribed into cDNA using random primers and an appropriate RT-PCR kit  
133 (Invitrogen, USA). For *elovl5*, degenerate primers (E5F1 and E5R1, Table1) were designed on the  
134 basis of alignment of fish *elovl5* including rabbitfish (GU597350), cobia (FJ440239), zebrafish  
135 (*Danio rerio*) (NM\_200453) and rainbow trout (*Oncorhynchus mykiss*) (AY605100), and used for  
136 amplifying partial fragments of putative *elovl5* cDNA from *S. argus* by polymerase chain reaction  
137 (PCR). For *elovl4*, degenerate primers (E4F1 and E4R1, Table1) were designed on the basis on the  
138 alignment of several fish *elovl4* including cobia (HM026361), rabbitfish (JF320823), and zebrafish  
139 *Elov14a* (NM\_200796) and *Elov14b* (NM\_199972), and used for amplifying partial fragments of  
140 putative *elovl4* cDNA fragment by PCR. For both *elovl* cDNAs, PCR (RT-PCR kit, Invitrogen, USA)  
141 consisted of an initial denaturation at 94 °C for 3 min, followed by 35 cycles of denaturation at

142 94 °C for 30 s, annealing at 58 °C for 30 s and extension at 72 °C for 1 min, followed by a final  
143 extension at 72 °C for 10 min. For both genes, the PCR fragments of the expected size were  
144 subsequently cloned into the pMD18-T vector (Takara, Dalian, China) and sequenced (Sangon,  
145 Shang Hai, China). Gene-specific primers were then designed to produce the full-length cDNA by  
146 5' (primers E5R2/E5R3 and E4R2/E4R3 for *elovl5* and *elovl4*, respectively) and 3' (E5F2 and E5F3,  
147 E4F2 and E4F3) rapid amplification of cDNA ends (RACE) PCR (Gene Racer™ Kit, Invitrogen,  
148 USA) (Table1).

#### 149 ***Sequence and phylogenetic analysis of Elov15 and Elov14***

150 The deduced amino acid (aa) sequences of the newly cloned elongases were aligned with their  
151 corresponding orthologues from rabbitfish (Elov15, ADE34561; Elov14, ADZ73580), Nibe croaker  
152 (Elov15, ACR47973; Elov14, AJD80650), cobia (Elov15, ACJ65150; Elov14, ADG59898), Atlantic  
153 salmon (Elov15a, AAO13175; Elov15b, ACI62499; Elov14, ADJ95235) and zebrafish (Elov15,  
154 NP\_956747; Elov14b, NP956266) using ClustalW2 (Higgins & Sharp, 1989). The aa sequence  
155 identities between deduced Elov1 proteins from *S. argus* and other vertebrate homologues were  
156 compared by the EMBOSS Needle Pairwise Sequence Alignment tool ([http://www.  
157 ebi.ac.uk/Tools/psa/emboss\\_needle/](http://www.ebi.ac.uk/Tools/psa/emboss_needle/)). A phylogenetic tree comparing the aa sequence similarities of  
158 different types of elongases (Elov12, Elov14 and Elov15) from a variety of vertebrate lineages was  
159 constructed using the neighbor-joining method (Saitou & Nei, 1987). Confidence in the resulting  
160 phylogenetic tree branch topology was measured through bootstrapping through 1000 iterations.

#### 161 ***Functional characterization of cloned elongase genes in yeast***

162 Functional characterization of the *S. argus* putative elongase genes was conducted by expressing  
163 their open reading frame (ORF) in yeast *Saccharomyces cerevisiae*. Expression primers listed in  
164 Table 1 (*elovl5*: E5F4 and E5R4, *elovl4*: E4F4 and E4R4) containing restriction sites *Bam*HI and  
165 *Xba*I were designed for amplification of the *elovl5* and *elovl4* ORFs from liver and eye cDNA using  
166 high-fidelity DNA polymerase (TianGen, Beijing, China) under the following conditions: initial  
167 denaturation at 94 °C for 5 min, followed by 30 cycles of denaturation at 94 °C for 45 s, annealing  
168 at 58 °C for 45 s and extension at 72 °C for 3 min, and a final extension at 72 °C for 10 min. The  
169 DNA fragments were purified and digested with the corresponding restriction endonucleases (New  
170 England Biolabs, UK) and ligated into the yeast episomal plasmid pYES2 (Invitrogen). The  
171 recombinant plasmids (pYES2-*elovl5* or pYES2-*elovl4*) were transformed into *S. cerevisiae* (strain

172 INVSc1, Invitrogen) using the S.C. Easy Comp Transformation kit (Invitrogen).

173 A single colony expressing either the *elovl5* or *elovl4* ORF was grown on *S. cerevisiae*  
174 minimal medium minus uracil (SCMM<sup>-uracil</sup>). Stearidonic acid (18:4n-3),  $\gamma$ -linolenic acid (18:3n-6),  
175 EPA (20:5n-3), arachidonic acid (ARA, 20:4n-6), docosapentaenoic acid (DPA, 22:5n-3) or  
176 docosatetraenoic acid (DTA, 22:4n-6) were used as substrates for testing the elongase activity of the  
177 *S. argus elovl5* and *elovl4*. All the fatty acids were purchased from Cayman Chemicals Co (Ann  
178 Arbor, MI, USA). The PUFA substrates were added at final concentrations of 0.5 (C18), 0.75 (C20)  
179 and 1.0 (C22) mM (Li et al., 2010). After two days culture, yeast cells were harvested and washed  
180 as described previously (Li et al., 2010).

### 181 ***Lipid extraction and fatty acid analysis***

182 Yeast samples were homogenized in chloroform/methanol (2:1, v/v) containing 0.01 % BHT as  
183 antioxidant, and total lipid extracted according to the Folch method (Folch et al., 1957). Fatty acid  
184 methyl esters (FAME) were prepared by transesterification with boron trifluoride etherate (ca. 48 %,  
185 Acros Organics, NJ, USA) as described previously (Li et al., 2008; Xie et al., 2014). FAME were  
186 purified by TLC, resuspended in hexane (Berry, 2004), and separated using a gas chromatograph  
187 (GC2010-plus, Shimadzu, Japan) as described in detail previously (Li et al., 2010). The activity of  
188 elongase was calculated as the proportion of substrate fatty acid converted to elongated FA products  
189 as follows:  $100 \times [\text{individual product area} / (\text{all product areas} + \text{substrate area})]$  (Li et al., 2010).

### 190 ***Tissue distribution of elovl5 and elovl4 mRNA***

191 In order to determine the distribution of elongase mRNA in *S. argus*, total RNA (1  $\mu$ g) from eye,  
192 brain, liver, muscle, heart, gills, spleen, kidney and intestine was reverse transcribed into cDNA  
193 (Cloned AMV First-Strand cDNA Synthesis Kit, Invitrogen). RT-PCR was carried out with an  
194 initial denaturing step at 94 °C for 5 min, followed by 30 cycles of denaturation at 94 °C for 30 s,  
195 annealing at 58 °C for 30 s, extension at 72 °C for 40 s, and a final extension at 72 °C for 5 min.  
196 The expression of the housekeeping gene *18S rRNA* was used as internal control to check the  
197 efficiency of cDNA synthesis and cDNA integrity. The primer pairs used for RT-PCR are given in  
198 Table 1.

### 199 ***Expression of elovl5 and elovl4 mRNA in liver and eye in response to diets with different*** 200 ***18:3n-3/18:2n-6 ratios***

201 The levels of *elovl5* and *elovl4* mRNA were measured by quantitative real-time PCR (qPCR)  
202 in liver and eye, respectively, from fish fed on the experimental diets D1-D6. QPCR primers were

203 E5qF/E5qR and E4qF/E4qR for *elovl5* and *elovl4*, respectively (Table 1). One µg total RNA from  
204 liver and eye was reverse transcribed into cDNA according to manufacturers instruction (Takara).  
205 The 20 µL reaction system consisted of 2 µL diluted cDNA, 0.4 µL for each primer (10 µmol), 10  
206 uL SYBR Premix, and 7.2 µL sterile double distilled water. PCR amplifications were carried out  
207 using a Lightcycler 480 real-time PCR detection system (Roche, Switzerland) with an initial  
208 denaturing step at 95 °C for 30s, followed by 40 cycles of denaturation at 95 °C for 5s, with a final  
209 step at 60 °C for 31 s. The mRNA levels of *elovl5* and *elovl4* in the liver and eye of *S. argus* in each  
210 dietary groups were normalized relative to the expression of *18S rRNA* calculated by the  
211 comparative threshold cycle (Ct) method (Whelan et al., 2003).

## 212 **Statistics**

213 The *elovl* mRNA expression data were presented as means ± standard error of mean (n = 9).  
214 Differences in the expression of *elovl5* and *elovl4* (tissue distribution and nutritional regulations  
215 experiments) were analyzed by one-way ANOVA followed by Tukey's multiple comparison. All  
216 analyses were conducted using SPSS v17.0 (SPSS Inc., Chicago, IL, USA).

## 217 **Results**

### 218 **3.1. Sequence and phylogenetic analyses of *S. argus elovl5* and *elovl4* cDNAs**

219 The newly cloned *S. argus* elongase cDNAs were 1390 bp (*elovl5*) and 1484 bp (*elovl4*) in  
220 full-length, and deposited in the GenBank database with the accession numbers KF029625 and  
221 KF029624, respectively. The *elovl5*-like cDNA had a 885 bp ORF encoding a peptide of 294 aa,  
222 whereas the *elovl4*-like cDNA had a 918 bp ORF encoding a protein of 304 aa. When compared to  
223 other teleost Elov15 and Elov14 sequences, the *S. argus* Elov15 was 71-85 % identical to teleost  
224 Elov15 including zebrafish, Atlantic salmon, rabbitfish, Nibe croaker and cobia, while *S. argus*  
225 Elov14 shares aa sequence identities of 84-97 % to Elov14 from teleosts including zebrafish, Atlantic  
226 salmon, Nibe croaker, cobia and rabbitfish.

227 Similar to other teleost Elov1-like proteins, both *S. argus* Elov15 and Elov14 deduced proteins  
228 possessed the histidine box motif (HXXHH) conserved in the elongase family (Fig. 1) (Jakobsson et  
229 al., 2006). They have lysine or arginine residues at the carboxyl terminus, more specifically  
230 KXRXX in Elov15 and RXKXX in Elov14, regarded as putative endoplasmic reticulum (ER)  
231 retrieval signals. Five putative transmembrane-spanning regions containing hydrophobic aa  
232 stretches were predicted by comparison with other vertebrate ELOVL proteins (Fig. 1).

233 A neighbor-joining phylogenetic tree was constructed based on the deduced elongase aa  
234 sequences from Elovl2, Elovl4 and Elovl5 retrieved from fish and other vertebrate genomes. Our  
235 results showed that sequences from the same type of elongase (Elovl2, Elovl4 and Elovl5) clustered  
236 together regardless of the vertebrate lineage considered. Thus, the herein characterized *S. argus*  
237 Elovl4 and Elovl5 grouped with other orthologues from other fish and more distantly other  
238 vertebrates (Fig. 2).

### 239 **3.2. Functional characterization**

240 The two putative Elovl isolated from *S. argus* were functionally characterized by heterologous  
241 expression in yeast *S. cerevisiae* grown in the presence of the following FA substrates: 18:4n-3,  
242 18:3n-6, 20:5n-3, 20:4n-6, 22:5n-3 or 22:4n-6. The FA composition of control yeast transformed  
243 with the empty pYES2 vector was 16:0, 16:1 isomers (16:1n-9 and 16:1n-7), 18:0 and 18:1n-9, as  
244 well as any exogenously added PUFA substrate (data not shown). This was consistent with the  
245 earlier observations that *S. cerevisiae* lacks PUFA elongase activity (Agaba et al., 2004; Monroig et  
246 al., 2012). Interestingly, yeast transformed with pYES2-*elovl5* were able to convert C18 to C22  
247 PUFA substrates to corresponding elongated products (Fig. 3). As shown in Table 2, *S. argus* Elovl5  
248 had an apparent preference for C18 and C20 over C22 FA substrates. Moreover, n-3 PUFA were  
249 elongated to a greater extent compared to their corresponding n-6 isomers with, for example, almost  
250 73% of added 18:4n-3 was elongated whereas only 40% of added 18:3n-6 was elongated.

251 When the *S. argus* Elovl4 cDNA was expressed in the yeast cells, evidence of elongation of all  
252 fatty acids was observed (Fig. 4, Table 2). Moreover, *S. argus* Elovl4 was more effectively convert  
253 both C20 and C22 PUFA substrates to C24 products with no obvious preference in terms of fatty  
254 acyl chain length (C20 vs C22) or FA series (n-3 vs n-6).

### 255 **3.3. Tissue expression of Elovl4 and Elovl5**

256 RT-PCR was used to analyze the expression of *elovl5* and *elovl4* in *S. argus* tissues (Fig. 5).  
257 The transcript of *elovl5* was detected in all tissues, with apparent higher expression levels in liver  
258 compared to eye, intestine and brain. However, the expression of *elovl4* was only detected in eye.  
259 As expected, the housekeeping gene *18S rRNA* was expressed in all tissues analyzed (Fig. 5).

### 260 **3.4. Effects of dietary fatty acid composition on the mRNA expression level of elovl4 and elovl5**

261 Nutritional regulation of the newly cloned Elovl5 was analyzed by qPCR in liver (*elovl5*) and  
262 eye (*elovl4*). Compared to fish fed the control diet based on FO (D2), livers from fish fed vegetable  
263 oil-based diets showed higher ( $P < 0.05$ ) expression of *elovl5* except for fish fed diet D3, which had

264 with the same LNA/LA ratio as D2 (Fig. 6B). The highest expression level of *elovl5* was detected in  
265 fish fed diet D5 with a dietary LNA/LA ratio of 1.72 (Fig. 6A). A similar pattern was observed for  
266 the expression of *elovl4* in eye. Thus, fish fed all vegetable oil-based diets, except D3, showed a  
267 significantly higher expression of *elovl4* compared to fish fed FO. Like *elovl5* in liver, the highest  
268 expression of *elovl4* in eye was observed in fish fed diet D5 (Fig. 6B).

269

## 270 **Discussion**

271 Elov1 enzymes account for the condensation step of the elongation reaction resulting in the  
272 addition of 2 carbon atoms to the pre-existing FA substrate (Jakobsson et al., 2006; Guillou et al.,  
273 2010). Investigations in many fish species have demonstrated that Elov15 preferentially elongates  
274 C18 and C20 PUFA, with residual conversion toward C22 substrates (Agaba et al., 2004, 2005;  
275 Monroig et al., 2012). On the other hand, Elov14 has been regarded as participating in the  
276 biosynthesis of very long-chain (> C24) PUFA, although studies in fish have revealed a role in the  
277 biosynthetic pathways of long-chain (C20-24) PUFA (Castro et al., 2016). To date, *elovl5* cDNAs  
278 have been identified in numerous fish species, while Elov14 has been studied in a lesser number of  
279 species (Castro et al., 2016). In the present study, we provide evidence for the existence of both  
280 Elov15 and Elov14 encoding cDNAs and demonstrate their role in the biosynthesis of long-chain  
281 PUFA in the euryhaline teleost *S. argus*.

282 The deduced aa sequences of *S. argus* Elov15 and Elov14, containing all the main structural  
283 features common for Elov1 protein family members (Jakobsson et al., 2006), shared high similarity  
284 to other fish orthologues. Consistent with this, the functional characterization of the newly cloned  
285 Elov1-like encoding cDNAs showed similar substrate specificities as those described in rabbitfish.  
286 Thus the *S. argus* Elov15 has high activity towards C18 and C20 PUFA substrates, and relatively  
287 low activity towards C22 PUFA. In addition to rabbitfish (Monroig et al., 2012), these results are  
288 consistent with previously reported activities in other Elov15 proteins characterized in cobia (Zheng  
289 et al., 2009), southern bluefin tuna (*Thunnus maccoyii*) (Gregory et al., 2010), Asian sea bass (*Lates  
290 calcarifer*) (Mohd-Yusof et al., 2010), Atlantic bluefin tuna (*Thunnus thynnus* L.) (Morais et al.,  
291 2011), rabbitfish (Monroig et al., 2012), Nibe croaker (Kabeya et al., 2015). In contrast to the  
292 Elov15, *S. argus* Elov14 may effectively elongate the tested C20 and C22 PUFA, generating  
293 products up to C24 in length. However, previously reported Elov14 from zebrafish, Atlantic salmon,  
294 cobia, rabbitfish and orange-spotted grouper Elov14, have the ability to catalyze the conversion of

295 C20 and C22 PUFA up to C36 PUFA (Monroig et al., 2010; Carmona-Antoñanzas et al., 2011;  
296 Monroig et al., 2011c, 2012; Li et al., 2015). The capability of the *S. argus* Elovl4 for elongation of  
297 C22 and C24 PUFA substrates up to C36 products was not found, and a recent investigation on the  
298 Nibe croaker Elovl4 (Kabeya et al., 2015) reported similar activities as those obtained for *S. argus*  
299 Elovl4, i.e. elongation products up to C24. Although elongation products longer than C24 could  
300 have been produced in yeast at amounts below the detection level, this results were similar to  
301 those reported in other fish species (Monroig et al., 2011b) and mammals (Agbaga et al., 2008).  
302 Taken together, it is still reasonable to believe that the *S. argus* Elovl4 plays a prominent role in the  
303 biosynthesis of VLC-PUFA. This is consistent with the mRNA tissue distribution showing that the *S.*  
304 *argus elovl4* was highly expressed in eye, a major metabolic site for VLC-PUFA (Agbaga et al.,  
305 2010) where these compounds accumulate in photoreceptor cell phospholipids (Aveldaño, 1988;  
306 Agbaga et al., 2010; Harkewicz et al., 2012).

307 The restricted pattern of *elovl4* mRNA found in *S. argus* tissues is largely consistent with that of  
308 rabbitfish, although brain also showed expression of *elovl4* in the latter (Monroig et al., 2012). In  
309 contrast, more widespread distribution of *elovl4* mRNA has been observed in the marine teleosts  
310 cobia and orange-spotted grouper, where *elovl4* transcripts were detected in eye, brain, testis, liver,  
311 kidney, muscle and stomach (Monroig et al., 2011c; Li et al., 2015). Although further studies are  
312 required to draw a firm conclusion, the difference in the distribution pattern of *elovl4* mRNAs of *S.*  
313 *argus* and rabbitfish compared to other teleost fish may be linked to their feeding habits. For *S.*  
314 *argus elovl5* mRNA, a wide spread distribution pattern was obtained, with greatest expression level  
315 in liver, eye, intestine and brain. This is similar to the tissue expression pattern obtained from  
316 rabbitfish, in which the expression of an *elovl5* was greatest in liver, followed by intestine and brain.  
317 In contrast, studies on carnivorous marine fish, including cobia, Asian sea bass, Nibe croaker,  
318 meagre, Japanese eel and Northern pike, showed that the expression of *elovl5* transcript was  
319 substantially higher in brain than other tissues (Zheng et al., 2009; Mohd-Yusof et al., 2010;  
320 Yamamoto et al., 2010; Monroig et al., 2013; Carmona-Antoñanzas et al., 2013; Wang et al., 2014).

321 From the functional and phylogenetic analysis, and tissue distribution of *elovl5* and *elovl4*  
322 transcripts, it is possible to conclude that the elongase complement involved in LC-PUFA  
323 biosynthesis in *S. argus* is similar to that characterized in rabbitfish (Monroig et al., 2012). While it  
324 is unclear if phylogeny and/or feeding habits can partly explain such similarity between *S. argus*  
325 and rabbitfish elongation capability, it is clear that such resemblance does not extend to Fads

326 complement. Thus, the sole *Fads2* found in *S. argus* (Xie et al., 2014), as observed in many fish  
327 species, such as cobia, Nibe croaker, Japanese eel (*Anguilla japonica*) and common carp (*Cyprinus*  
328 *carpio* var. Jian) (Zheng et al., 2009; Yamamoto et al., 2010; Wang et al., 2014; Kabeya et al., 2015;  
329 Ren et al., 2013), was characterized with  $\Delta 6$ -desaturase activity but no  $\Delta 5$ - and  $\Delta 4$ -desaturase (Xie  
330 et al., 2014). In contrast, the rabbitfish possess at least two *Fads2* desaturases, a dual  $\Delta 6/\Delta 5$   
331 desaturase and a  $\Delta 4$  desaturase, the latter being the first record of a  $\Delta 4$  desaturation activity in  
332 vertebrates (Li et al., 2010). Additionally, the distribution of *S. argus fads2* mRNA, with highest  
333 expression in liver, followed a pattern typically found in freshwater/salmonid species in contrast to  
334 carnivorous marine species whereby brain has shown the highest levels of *fads2* transcription  
335 (Monroig et al., 2011b). These results further confirm the enormous diversification of fish  
336 LC-PUFA biosynthesis specificities that has been previously hypothesized to be associated with  
337 factors including habitat, trophic level and ecology, as well as species-specific evolutionary history  
338 (Fonseca-Madrigal et al., 2014).

339 The ability of fish to regulate LC-PUFA biosynthesis has been extensively investigated in  
340 commercially important species in order to understand the metabolic impact of replacing FO by VO  
341 devoid of LC-PUFA in aquafeed (Ling et al., 2006; Jordal et al., 2005; Li et al., 2008; Thanuthong  
342 et al., 2011; Navarro-Guillen et al., 2014; Xie et al., 2014, 2015; Kuah et al., 2015). We herein  
343 showed that dietary lipid resource also affected the expression of *elovl5* and *elovl4* in *S. argus*.  
344 Generally speaking, both *elovl4* and *elovl5* were up-regulated in *S. argus* in response to low dietary  
345 LC-PUFA (high VO) input. While nutritional regulation of *elovl5* in liver has often been reported  
346 (Ling et al., 2006; Morais et al., 2009; Yamamoto et al., 2010; Thanuthong et al., 2011;  
347 Navarro-Guillen et al., 2014; Xie et al., 2015; Kuah et al., 2015), regulatory mechanisms of  
348 LC-PUFA in eye have less been investigated despite eye accumulating large amounts of LC-PUFA  
349 in some species (Aveldaño, 1988; Agbaga et al., 2010; Harkewicz et al., 2012). Li et al. (2015)  
350 recently described an up-regulation of *elovl4* in viscera of orange-spotted grouper larvae. Although  
351 the specific mechanism remains to elucidated, it is largely accepted that the increased expression of  
352 key enzymes including *elovl* and *fads* involved in LC-PUFA biosynthesis pathway stimulated by  
353 dietary VO is a biochemical/molecular mechanism that can at least partially compensate dietary  
354 essential fatty acid deficiencies (Tocher et al., 2003).

355 In addition to the dietary lipid source (FO v. VO), the ratio of dietary LNA and LA also  
356 influenced the expression of key enzymes involved in LC-PUFA biosynthesis. Functional

357 characterization of Fads and Elovl isolated from various teleosts species have often revealed a  
358 higher activity towards n-3 compared to n-6 PUFA substrates (Morais et al., 2009; Zheng et al.,  
359 2009; Li et al., 2010; Monroig et al., 2012; Monroig et al., 2013; Xie et al., 2014; Kabeya et al.,  
360 2015). Previous studies have revealed that Fad gene expression and enzymatic activity varied with  
361 dietary LNA/LA ratio. For example, up-regulation of  $\Delta 6$  *fads2* gene expression was measured in  
362 rabbitfish, Murray cod (*Maccullochella peelii peelii*), rainbow trout and *S. argus* fed high dietary  
363 ratios of LNA/LA (Li et al., 2008; Senadheera et al., 2011; Thanuthong et al., 2011; Xie et al.,  
364 2014). An excess of LNA in the diet could also block  $\Delta 6$  *fads* gene expression (Izquierdo et al.,  
365 2008; Xie et al., 2014). Unlike desaturases, few studies have reported the influence of dietary  
366 LNA/LA ratio on elongase gene expression. In the present study, the mRNA expression of *elovl5*  
367 and *elovl4* was highest in liver and eye of *S. argus* fed a diet with an LNA/LA ratio of around 1.72.  
368 As prior to the elongase activity, the  $\Delta 6$  desaturase acting on LNA and LA increased with the elevation  
369 of the dietary ALA/LA ratio (Thanuthong et al., 2011; Xie et al., 2014). In our previous study, the  
370 highest  $\Delta 6$  *fads* mRNA expression was detected in liver of *S. argus* fed a diet with an LNA/LA ratio of  
371 1.72 (Xie et al., 2014), which means the high level Elovl substrate are available to Elovl5 and Elovl4 in  
372 the dietary treatment with an LNA/LA ratio of 1.72. Therefore, it appears that dietary ratio of  
373 LNA/LA can influence the expression of *fads* and *elovl* and LC-PUFA biosynthesis efficiency  
374 could be optimized with particular dietary levels of C18 PUFA.

375 In summary, the present study showed that *S. argus* has at least two Elovl with high sequence,  
376 function and distribution homology to Elovl4 and Elovl5 reported previously in another herbivorous  
377 species, the rabbitfish. However, the Fads complement of *S. argus* and rabbitfish is remarkably  
378 different and this suggests that the diversification of fish LC-PUFA biosynthesis specificities is  
379 highly varied. Replacing FO with a VO blend with a dietary ratio of LNA/LA of 1.72 resulted in  
380 highest expression of *elovl5* and *elovl4* in liver and eye of *S. argus*, respectively. These discoveries  
381 will expand our knowledge in understanding the molecular basis and regulation of LC-PUFA  
382 biosynthesis in fish.

383

#### 384 **Acknowledgements**

385 This work was financially supported by the Major International Joint Research Project from  
386 the National Natural Science Foundation of China (NSFC) (31110103913), NSFC General Projects  
387 (No. 41276179) and Youth Projects (Nos. 31202011, 31202012).

388 **References**

- 389 Agaba, M., Tocher, D.R., Dickson, C., Dick, J.R., Teale, A.J., 2004. Zebrafish cDNA encoding multifunctional  
390 fatty acid elongase involved in production of eicosapentaenoic (20:5n-3) and docosahexaenoic (22:6n-3)  
391 acids. *Mar. Biotechnol.* 6, 251–261.
- 392 Agaba, M.K., Tocher, D.R., Dickson, C.A., Zheng, X., Dick, J.R., Teale, A.J., 2005. Cloning and functional  
393 characterisation of polyunsaturated fatty acid elongases from marine and freshwater teleost fish. *Comp.*  
394 *Biochem. Physiol. B* 142, 342–352.
- 395 Agbaga, M.P., Mandal, M.N.A., Anderson, R.E., 2010. Retinal very long chain polyunsaturated fatty acids: new  
396 insights from studies on ELOVL4 protein. *J. Lipid Res.* 51, 1624-1642.
- 397 Aveldaño, M.I., 1988. Phospholipid species containing long and very long polyenoic fatty acids remain with  
398 rhodopsin after hexane extraction of photoreceptor membranes. *Biochemistry* 27, 1229-1239.
- 399 Awada, M., Meynier, A., Soulage, C. O., Hadji, L., Gélöën, A., Viau, M., Lagard, M., 2013. N-3 PUFA added to  
400 high-fat diets affect differently adiposity and inflammation when carried by phospholipids or triacylglycerols  
401 in mice. *Nutr. Metab.* 10, 220-230.
- 402 Barry, T.P., Fast, A.W., 1992. Biology of the spotted scat (*Scatophagus argus*) in the Philippines. *Asian Fish Sci.* 5,  
403 163-179.
- 404 Campoy, C., Escolano-Margarit, V., Anjos, T., Szajewska, H., Uauy, R., 2012. Omega 3 fatty acids on child  
405 growth, visual acuity and neurodevelopment. *Br. J. Nutr.* 107, 85-106.
- 406 Carmona-Antoñanzas, G., Monroig Ó., Dick J.R., Davie, A., Tocher, D.R., 2011. Biosynthesis of very long-chain  
407 fatty acids (C>24) in Atlantic salmon: cloning, functional characterisation, and tissue distribution of an  
408 Elov14 elongase. *Comp. Biochem. Physiol. B* 159, 122-129.
- 409 Carmona-Antoñanzas, G., Tocher, D.R., Taggart, J.B., Leaver, M.J., 2013. An evolutionary perspective on Elov15  
410 fatty acid elongase: comparison of Northern pike and duplicated paralogs from Atlantic salmon. *BMC*  
411 *evolutionary biology*, 13, 85.
- 412 Castro, L.F.C., Monroig, Ó., Leaver, M.J., Wilson, J., Cunha, I., Tocher, D.R., 2012. Functional desaturase Fads1  
413 ( $\Delta 5$ ) and Fads2 ( $\Delta 6$ ) orthologues evolved before the origin of jawed vertebrates. *PLoS One* 7: e31950.
- 414 Castro, L.F.C., Tocher, D.R., Monroig, Ó., 2016. Long-chain polyunsaturated fatty acid biosynthesis in chordates:  
415 insights into the evolution of Fads and Elovl gene repertoire. *Prog. Lipid Res.* doi:10.1016/j.plipres.2016.  
416 01.001.
- 417 Delgado-Lista, J., Perez-Martinez, P., Lopez-Miranda, J., Perez-Jimenez, F., 2012. Long chain omega-3 fatty acids  
418 and cardiovascular disease: a systematic review. *Br. J. Nutr.* 107, 201-213.

419 Folch, J., Lees, M., Stanley, S., 1957. A simple method for the isolation and purification of the total lipids from  
420 animal tissues. *Biol. Chem.* 226, 497-509.

421 Fonseca-Madrigal, J., Navarro, J. C., Hontoria, F., Tocher, D. R., Martínez-Palacios, C. A., Monroig, Ó., 2014.  
422 Diversification of substrate specificities in teleostei Fads2: characterization of  $\Delta 4$  and  $\Delta 6/\Delta 5$  desaturases of  
423 *Chirostoma estor*. *J. Lipid Res.* 55, 1408-1419.

424 Gandhi, V., 2002. Studies on the food and feeding habits of cultivable butterfish *Scatophagus argus* (Cuv. and  
425 Val.). *J. Mar. Biol. Ass India* 44: 115-121.

426 Gregory, M., See, V.H.L., Gibson, R.A., Shuller, K.A., 2010. Cloning and functional characterisation of a fatty  
427 acyl elongase from southern bluefin tuna (*Thunnus maccoyii*). *Comp. Biochem. Physiol. B* 155, 178-185.

428 Guillou, H., Zadavec, D., Martin, P.G.P., Jacobsson, A., 2010. The key roles of elongases and desaturases in  
429 mammalian fatty acid metabolism: Insights from transgenic mice. *Prog. Lipid Res.* 49, 186-99.

430 Harkewicz, R., Du, H., Tong, Z., Alkuraya, H., Bedell, M., Sun, W., Su, Z., 2012. Essential role of ELOVL4  
431 protein in very long chain fatty acid synthesis and retinal function. *J. Biol. Chem.* 287, 11469-11480.

432 Higgins, D., Sharp, P., 1989. Fast and sensitive multiple sequence alignments on a microcomputer. *J. Comput.*  
433 *Anal. Appl.* 5, 151-153.

434 Izquierdo, M.S., Robaina, L., Juárez-Carrillo, E., Oliva, V., Hernandez-Cruz, C.M., Afonso, J.M., 2008.  
435 Regulation of growth, fatty acid composition and delta-6 desaturase expression by dietary lipids in gilthead  
436 seabream larvae (*Sparus aurata*). *Fish Physiol. Biochem.* 34, 117-127.

437 Jakobsson, A., Westerberg, R., Jacobsson, A., 2006. Fatty acid elongases in mammals: Their regulation and roles  
438 in metabolism. *Prog. Lipid Res.* 45, 237-49.

439 Jordal, A.-E.O., Torstensen, B.E., Tsoi, S., Tocher, D.R., Lall, S.P., Douglas, S.E., 2005. Dietary rapeseed oil  
440 affects the expression of genes involved in hepatic lipid metabolism in Atlantic salmon (*Salmo salar* L.). *J.*  
441 *Nutr.* 135, 2355–2361.

442 Kabeya, N., Yamamoto, Y., Cummins, S.F., Elizur, A., Yazawa, R., Takeuch, Y., Haga, Y., Satoh, S., Yoshizaki, G.,  
443 2015. Polyunsaturated fatty acid metabolism in a marine teleost, Nibe croaker *Nibea mitsukurii*: Functional  
444 characterization of Fads2 desaturase and Elov15 and Elov14 elongases. *Comp. Biochem. Physiol. B* 188,  
445 37-45.

446 Kuah, M.K., Jaya-Ram, A., Shu-Chien, A.C., 2015. The capacity for long-chain polyunsaturated fatty acid  
447 synthesis in a carnivorous vertebrate: Functional characterisation and nutritional regulation of a Fads2 fatty  
448 acyl desaturase with  $\Delta 4$  activity and an Elov15 elongase in striped snakehead (*Channa striata*). *Biochim.*  
449 *Biophys. Acta* 1851, 248-260.

- 450 Li, S., Monroig, Ó., Navarro, J.C., Yuan, Y., Xu, W., Mai, K., Tocher, D.R., Ai, Q., 2015. Molecular cloning and  
451 functional characterization of a putative Elovl4 gene and its expression in response to dietary fatty acid  
452 profiles in orange-spotted grouper *Epinephelus coioides*. *Aquac. Res.* DOI: 10.1111/are.12902.
- 453 Li, Y.Y., Hu, C.B., Zheng, Y.J., Xia, X.A., Xu, W.J., Wang, S.Q., Chen, W.Z., Sun, Z.W., Huang, J.H., 2008. The  
454 effects of dietary fatty acids on liver fatty acid composition and  $\Delta^6$ -desaturase expression differ with ambient  
455 salinities in *Siganus canaliculatus*. *Comp. Biochem. Physiol. B* 151, 183-190.
- 456 Li, Y., Monroig, Ó., Zhang, L., Wang, S., Zheng, X., Dick, J.R., You, C., Tocher, D.R., 2010. Vertebrate fatty acyl  
457 desaturase with  $\Delta^4$  activity. *Proc. Natl. Acad. Sci. U. S. A.* 107,16840-16845.
- 458 Ling, S., Kuah, M.K., Muhammad, T.S.T., Kolkovski, S., Shu-Chien, A.C., 2006. Effect of dietary HUFA on  
459 reproductive performance, tissue fatty acid profile and desaturase and elongase mRNAs in female swordtail  
460 *Xiphophorus helleri*. *Aquaculture* 261, 204-214.
- 461 Mohd-Yusof, N.Y., Monroig, Ó., Mohd-Adnan, A., Wan, K.L., Tocher, D.R., 2010. Investigation of highly  
462 unsaturated fatty acid metabolism in the Asian sea bass, *Lates calcarifer*. *Fish Physiol. Biochem.* 3,827-843.
- 463 Monroig, Ó., Rotllant, J., Reverter, J.M.C, Dick, J.R., Figueras, A., Tocher, D.R., 2010. Expression and role of  
464 Elovl4 elongases in biosynthesis of very long-chain fatty acids during zebrafish *Danio rerio* early embryonic  
465 development. *Biochim. Biophys. Acta* 1801, 1145-1154.
- 466 Monroig, Ó., Li, Y., Tocher, D. R. 2011a. Delta-8 desaturation activity varies among fatty acyl desaturases of  
467 teleost fish: high activity in delta-6 desaturases of marine species. *Comp. Biochem. Physiol. B* 159, 206-213.
- 468 Monroig, Ó., Navarro, J.C., Tocher, D.R., 2011b. Long-chain poly unsaturated fatty acids in fish: recent advances  
469 on desaturases and elongases involved in their biosynthesis. *Avances. Nutr. Acuicola*, 11, 257-283.
- 470 Monroig, Ó., Webb, K., Castro, L.I., Holt, G.J., Tocher, D.R., 2011c. Biosynthesis of long-chain polyunsaturated  
471 fatty acids in marine fish: Characterization of an Elovl4-like elongase from cobia *Rachycentron canadum* and  
472 activation of the pathway during early life stages. *Aquaculture* 312, 145-153.
- 473 Monroig, Ó., Wang, S.Q., Zhang, L., You, C.H., Tocher, D.R, Li, Y.Y., 2012. Elongation of long-chain fatty acids  
474 in rabbitfish *Siganus canaliculatus*: Cloning, functional characterization and tissue distribution of Elovl5-  
475 and Elovl4-like elongases. *Aquaculture* 350-353, 63-70.
- 476 Monroig, Ó., Tocher, D.R., Hontoria, F., Navarro, J.C., 2013. Functional characterisation of a Fads2 fatty acyl  
477 desaturase with  $\Delta^6/\Delta^8$  activity and an Elovl5 with C16, C18 and C20 elongase activity in the anadromous  
478 teleost meagre (*Argyrosomus regius*). *Aquaculture* 412-413, 14-22.
- 479 Morais, S., Monroig, Ó., Zheng, X., Leaver, M.J., Tocher, D.R, 2009. Highly unsaturated fatty acid synthesis in  
480 Atlantic salmon: characterization of Elovl5- and Elovl2-like elongases. *Mar. Biotechnol.* 11, 627-639.

481 Morais, S., Mourente, G., Ortega, A., Tocher, J.A., Tocher, D.R., 2011. Expression of fatty acyl desaturase and  
482 elongase genes, and evolution of DHA:EPA ratio during development of Atlantic bluefin tuna (*Thunnus*  
483 *thynnus* L.). *Aquaculture* 313, 129–139.

484 Morais, S., Castanheira, F., Martínez-Rubio, L., Conceição, L.E.C., Tocher, D.R., 2012. Long-chain  
485 polyunsaturated fatty acid synthesis in a marine vertebrate: ontogenetic and nutritional regulation of a fatty  
486 acyl desaturase with  $\Delta 4$  activity. *Biochim. Biophys. Acta* 1821, 660-671.

487 Muhlhausler, B.S., Ailhaud, G.P., 2013. Omega-6 polyunsaturated fatty acids and the early origins of obesity. *Curr*  
488 *Opin Endocrinol.* 20, 56–61,

489 Navarro-Guillen, C., Engrola, S., Castanheira, F., Bandarra, N., Hachero-Cruzado, I., Tocher, D.R., Conceicao,  
490 L.E., Morais, S., 2014. Effect of varying dietary levels of LC-PUFA and vegetable oil sources on  
491 performance and fatty acids of Senegalese sole post larvae: puzzling results suggest complete biosynthesis  
492 pathway from C18 PUFA to DHA. *Comp. Biochem. Physiol. B* 167:51–58.

493 Óscar Monroig, Shuqi Wang, Liang Zhang, Cuihong You, Douglas R. Tocher, Yuanyou Li. 2012. Elongation of  
494 long-chain fatty acids in rabbitfish *Siganus canaliculatus*: Cloning, functional characterization and tissue  
495 distribution of Elovl5-and Elovl4-like elongases. *Aquaculture*, 350-353: 63-70.

496 Ren, H.T., Zhang, G.Q., Li, J.L., Tang, Y.K., Li, H.X., Yu, J. H., Xu, P., 2013. Two  $\Delta 6$ -desaturase like genes in  
497 common carp (*Cyprinus carpio* var. Jian): Structure characterization, mRNA expression, temperature and  
498 nutritional regulation. *Gene* 525: 11–17.

499 Saitou, N., Nei, M., 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol.*  
500 *Biol. Evol.* 4, 406-425.

501 Sargent, J.R., Tocher, D.R., Bell, J.G., 2002. The lipids. In: *Fish Nutrition* (Halver, J.E. & Hardy, R.W. eds), pp.  
502 181–257. Academic Press, San Diego.

503 Senadheera, S.D., Turchini, G.M., Thanuthong, T., Francis, D.S., 2011. Effects of dietary  $\alpha$ -linolenic acid (18:  
504 3n-3)/linoleic acid (18: 2n-6) ratio on fatty acid metabolism in Murray cod (*Maccullochella peelii peelii*).  
505 *Journal of agricultural and food chemistry*, 59(3), 1020-1030.

506 Thanuthong, T., Francis, D.S., Senadheera, S.P.S.D., Jones, P. L., Turchini, G.M. 2011. LC-PUFA biosynthesis in  
507 rainbow trout is substrate limited: use of the whole body fatty acid balance method and different 18: 3n-3/18:  
508 2n-6 ratios. *Lipids*, 46, 1111-1127.

509 Thanuthong, T., Francis, D.S., Manickam, E., Senadheera, S.D., Cameron-Smith, D., Turchini, G.M., 2011. Fish  
510 oil replacement in rainbow trout diets and total dietary PUFA content: II) Effects on fatty acid metabolism  
511 and in vivo fatty acid bioconversion. *Aquaculture* 322, 99–108.

512 Torstensen, B.E., Tocher, D.R., 2010. The effects of fish oil replacement on lipid metabolism of fish. In: Turchini,  
513 G.M., Ng, W.K., Tocher (eds), Fish oil replacement and alternative lipid sources in aquaculture feeds, pp.  
514 405–437. CRC Press, Taylor & Francis group, Boca Raton, Florida, USA.

515 Tocher, D.R., Bell, J.G., Dick, J.R., Crampton, V.O., 2003. Effects of dietary vegetable oil on Atlantic salmon  
516 hepatocyte fatty acid desaturation and liver fatty acid compositions. *Lipids* 38, 723–732.

517 Tocher, D.R., 2010. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquac. Res.* 41, 717-732.

518 Wang, S., Monroig, Ó., Tang, G., Zhang, L., You, C., Tocher, D.R., Li, Y. 2014. Investigating long-chain  
519 polyunsaturated fatty acid biosynthesis in teleost fish: Functional characterization of fatty acyl desaturase  
520 (Fads2) and Elovl5 elongase in the catadromous species, Japanese eel *Anguilla japonica*. *Aquaculture* 434,  
521 57-65.

522 Woodland, D.J., 1983. Zoogeography of the Siganidae (Pisces): An interpretation of distribution and richness  
523 patterns. *B. Mar. Sci.* 33, 713-717.

524 Xie, D., Chen, F., Lin, S., Wang, S., You, C., Monroig, Ó., Tocher, D.R., Li, Y., 2014. Cloning, functional  
525 characterization and nutritional regulation of  $\Delta 6$  fatty acyl desaturase in the herbivorous euryhaline teleost  
526 *Scatophagus Argus*. *PloS one*, 9, e90200.

527 Xie, D., Wang, S., You, C., Chen, F., Monroig, Ó., Tocher, D.R., Li, Y. 2015. Characteristics of LC - PUFA  
528 biosynthesis in marine herbivorous teleost *Siganus canaliculatus* under different ambient salinities. *Aquacult.*  
529 *Nutr.* 21, 541-551.

530 You, C., Zeng, F., Wang, S., Li, Y., 2014. Preference of the Herbivorous Marine Teleost *Siganus canaliculatus* for  
531 Different Macroalgae. *J. Ocean Univ. China* 13, 516- 522.

532 Yoshimura, K., Yamane, T., Utsugi, K., Kohno, H., 2003. Seasonal occurrence and abundance of the spotted scat,  
533 *Scatophagus argus*, in surf zones and rivers of the northern coast of Bali, Indonesia. *Mer.* 41, 82–85.

534 Yamamoto, Y., Kabeya, N., Takeuchi, Y., Alimuddin, Haga, Y., Satoh, S., Takeuchi, T., Yoshizaki, G., 2010.  
535 Cloning and nutritional regulation of polyunsaturated fatty acid desaturase and elongase of a marine teleost,  
536 the nibe croaker *Nibea mitsukurii*. *Fish Sci.* 76, 463-472.

537 Zhang, Q., Xie, D., Wang, S., You, C., Monroig, Ó., Tocher, D.R., Li, Y., 2014. miR-17 is involved in the  
538 regulation of LC-PUFA biosynthesis in vertebrates: Effects on liver expression of a fatty acyl desaturase in  
539 the marine teleost *Siganus canaliculatus*. *Biochim. Biophys. Acta* 1841, 934-943.

540 Zheng, X., Tocher, D.R., Dickson, C.A., Bell, J.G., Teale, A.J., 2005. Highly unsaturated fatty acid synthesis in  
541 vertebrates: new insights with the cloning and characterization of a  $\Delta 6$  desaturase of Atlantic salmon. *Lipids*  
542 40, 13–24.

543 Zheng, X., Ding, Z., Xu, Y., Monroig, O., Morais, S., Tocher, D.R., 2009. Physiological roles of fatty acyl  
544 desaturase and elongase in marine fish: characterisation of cDNAs of fatty acyl  $\Delta$ 6 desaturase and Elovl5  
545 elongase of cobia (*Rachycentron canadum*). Aquaculture 290, 122–131.

546

547

548 Table 1

549 Primers used for cDNA cloning or determining gene expression of *Scatophagus argus* elongases

Aim	Primer	Primer sequence	Accession No <sup>1</sup>
First fragment cloning	E5F1	5'-GGTACTACTTCTCCAAGCTCAT-3'	KF029625
	E5R1	5'-GTGATGTATCTCTTCCACC-3'	
	E4F1	5'-GTCTACAACCTTCAGCATGGTG-3'	KF029624
	E4R1	5'-GGAACCTGGATCATCTGAATAA-3'	
3'RACE	E5F2	5'-ACAGCTTCGTCCACGTCGTGATGTA-3'	KF029625
	E5F3	5'-TTCGTTATGAACTGGCAACCCTGTG-3'	
	E4F2	5'-TGGCAGCCTTGGGACCTCAG-3'	KF029624
	E4F3	5'-GTGGATTGGCATCAAATGGGTC-3'	
5'RACE	E5R2	5'-TTCAGCATGGTAGCGTGGTGGTAGA-3'	KF029625
	E5R3	5'-TGTTTATGGCGGCACCGAAGTATGA-3'	
	E4R2	5'-GCCAGGGATGTAAGGGTTTCTTCAGAC-3'	KF029624
	E4R3	5'-GTGGATGGAAGAGTTGATGGTTGC-3'	
ORF cloning	E5F4	5'-CCCAAGCTTCAAATGGAGACCATCAATC-3'	KF029625
	E5R4	5'-CCGCTCGAGTCAATCCATCCTCAGCTT-3'	
	E4F4	5'-CCCAAGCTTGCCATGGAGGTTGTAACAC-3'	KF029624
	E4R4	5'-CCGCTCGAGTTACTCTCTTTTTGCTCT-3'	
RT-PCR and qPCR	E5qF	5'-ATGAACTGGCAACCCTGTGG-3'	KF029625
	E5qR	5'-ATATGGCTGCACACATCGTCTG-3'	
	E4qF	5'-TAGCAGACAAGAGGGTGGAGAA-3'	KF029624
	E4qR	5'-CTATGAGGGTCTTCCTGAGTGTA-3'	
	18SF	5'-CGCCGAGAAGACGATCAAAC-3'	AJ427629
	18SR	5'-TGATCCTTCCGCAGGTTTAC-3'	

550 <sup>1</sup> GenBank(<http://www.ncbi.nlm.nih.gov/>)

551

552

553

554

555

556

557

558

559

560

561

562

563 Table 2

564 Functional characterization of the *Scatophagu argus* elongases in *Saccharomyces cerevisiae* yeast.  
565 Results are expressed as a percentage of total substrate fatty acid converted to elongated products.

Elongase	Substrate fatty acid	Product	%Conversion	Activity
Elov15	18:3n-6	20:3n-6	43.9	C18→20
		22:3n-6	10.3	C20→22
	18:4n-3	20:4n-3	72.9	C18→20
		22:4n-3	20.8	C20→22
	20:4n-6	22:4n-6	32.0	C20→22
		24:4n-6	6.9	C22→24
	20:5n-3	22:5n-3	35.8	C20→22
		24:5n-3	10.1	C22→24
	22:4n-6	24:4n-6	7.5	C22→24
		22:5n-3	24:5n-3	11.5
Elov14	18:3n-6	20:3n-6	7.6	C18→20
		20:4n-3	12.3	C18→20
	20:4n-6	22:4n-6	37.3	C20→22
		24:4n-6	19.3	C22→24
	20:5n-3	22:5n-3	35.2	C20→22
		24:5n-3	21.6	C22→24
	22:4n-6	24:4n-6	26.5	C22→24
		22:5n-3	24:5n-3	34.8

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581 SaE5 METINLKLNAQLETWIGPR---DQRVVGWLLLDNYPPTFALTVIYLLIVWMGPKYMKYRQPYSCRGLLVFYNLGLTLL 75

582 ScE5 MEDFNRLNSYFESWIGPR---DQRLQGWLLLDNYPPTFALTVVYLLIVWLGPKYMKNRPAYSCRGLMVIYNLGLTLL 75

583 RcE5 METFNHKLNAYIESWMGPR---DQRVKGWLLLDNYPPTFALTVMYLLIVWMGPKYMKHRQPYSCRGLLVLYNLGLTLL 75

584 NmE5 METFNHKLNTYLESWMGPR---DQRVVGWLLLDNYPPTFALTVMYLIVWMGPKYMKHRQPYSCRGLLVLYNLGLTLL 75

585 SsE5a METFNKYLNMYIDSWMGPR---DERVQGWLLLDNYPPTFALTVMYLLIVWLGPKYMRHRQPVSCRGLLVLYNLGLTIL 75

586 SsE5b MEAFNHKLNTYIDSWMGPR---DERVQGWLLLDNYPPTFALTLMYLLIVWLGPKYMRHRQPVSCQGLLVLYNLALTLL 75

587 DrE5 METFSHRVNSYIDSWMGPR---DLRVTVGWFLDDYIPTFIFTVMYLLIVWMGPKYMKNRQAYSCRALLVPYNLCLTLL 75

588 SaE4 MEVVTHFVNDTVEFYKWSLTIADKRVEKWPMMSSPLPTLAISCLYLLFLWAGPRYMQRDRQPCTLRKTLIVYNFSMVVL 78

589 ScE4 MEVVTHFVNDTVEFYKWSLTIADKRVEKWPMMSSPLPTLAISCLYLLFLWAGPRYMQRDRQPFTLRKTLIVYNFSMVVL 78

590 RcE4 MEVVTHFVNDTVEFYKWSLTIADKRVENWPMMASPLPTLAISCLYLLFLWVGPYMQDRQPYTLRRTLIVYNFSMVVL 78

591 NmE4 MEAVTHFVNDTVEFYKWGLTIADKRVENWPMMSPLPTLAISCLYLLFLWAGPRYMQRDRQPFTLRKTLIVYNFSMVVL 78

592 SsE4 MEAVTHFMNDTVEFYRWSLTIADKRVEKWPMMSSPAPTLAISCLYLLFLWAGPKYMQNREPFQLRKTLIVYNFSMVL 78

593 DrE4b METVVHLMNDSVEFYKWSLTIADKRVEKWPMMSSPLPTLGISVLYLLFLWAGPLYMQNREPFQLRKTLIVYNFSMVLL 78

594 \*\* . :\* .. :.: : \* \*: \* : : . \*\* : : : \*\* : : \* \* \* : \* : : : \*\* : : : : \* : : : \*

595 I II

596 SaE5 SFYMFYELVTAVWYGGYNFYCQNS-HSAEEADNKIMNVLWVWYFSKLI EFMDTFFFILRKNNHQISFLH **VYHH**ATMLN 150

597 ScE5 SFYMFYELGSAIWFGGYHFYCQNT-HSLPEMDNKVMRALWVWYFSKLI EFMDTFFFILRKNNHQITFLH **IYHH**ASMFN 150

598 RcE5 SFYMFYELVTAVWHGGYNFYCQDT-HSAEEVDNKI INVLWVWYFSKLI EFMDTFFFILRKNNHQITFLH **IYHH**ATMLN 150

599 NmE5 SFYMFYELVTAVWHGGYNFYCQDI-HSAQEVDNKI INVLWVWYFSKLI EFMDTFFFILRKNNHQITFLH **IYHH**ASMLN 150

600 SsE5a SFYMFYEMVSAVWHGDYNFYCQDT-HSAGETDTKI INVLWVWYFSKLI EFMDTFFFILRKNNHQITFLH **IYHH**ASMLN 150

601 SsE5b SFYMFYEMVSAVWQGGYNFYCQDT-HSAGETDTKI INVLWVWYFSKVI EFMDTFFFILRKNNHQITFLH **IYHH**ASMLN 150

602 DrE5 SLYMFYELVMSVYQGGYNFFCQNT-HSGGDADNRMMNVLWVWYFSKLI EFMDTFFFILRKNNHQITFLH **VYHH**ATMLN 150

603 SaE4 NFYIAKELLGSRAGYSYLCQPVNYSNDVNEVRIASALWVWYISKGV EFLDVTFFFILRKKFNQVSFLH **VYHH**CTMFI 154

604 ScE4 NFYIAKELLGSRAGYSYLCQPVNYSNDVNEVRIASALWVWYISKGV EFLDVTFFFILRKKFNQVSFLH **VYHH**CTMFI 154

605 RcE4 NFYIAKELLIATRAAGYSYLCQPVNYSNDVNEVRIASALWVWYISKGV EFLDVTFFFILRKKFNQVSFLH **VYHH**CTMFI 154

606 NmE4 NFYIAKELLGSRAGYSYLCQPVNYSNDVNEVRIASALWVWYISKGV EFLDVTFFFIMRKKFNQVSFLH **VYHH**CTMFI 154

607 SsE4 NFYIAKELLGARAAGYSYLCQPVSYSDVNEVRIASALWVWYISKGV EYLDVTFFFILRKKINQVSFLH **VYHH**CTMFI 154

608 DrE4b NFYICKELLGSRAGYSYLCQPVNYSNDVNEVRIASALWVWYISKGV EFLDVTFFFIMRKKFNQVSFLH **VYHH**CTMFI 154

609 :.: : \* : . . \* : \* \* : \* : : . \* \* \* : : \* \* : \* : \* \* \* \* \* : \* : : \* \* \* \* \* : \* :

610 II

611 SaE5 IWWFVMNWQPCGHSYFGAAINSFVHVVMYSYYGLSAI-PGIRPYLWKKYITQLQMIQFFLTMCQTMCAAIWPCGVVP 225

612 ScE5 IWWFVMNWIPCGHSYFGASLNSFVHVVMYSYYGLSAV-PSLRPYLWKKYITQLQLVQFFLTMFQTYCAVLWPCGFPI 225

613 RcE5 IWWFVMNWIPCGHSYFGASLNSFVHVVMYSYYGLSAI-PAMRPYLWKKYITQLQLIQFFLTMSQTMCAVIWPCDFPR 225

614 NmE5 IWWFVMNWVPCGHSYFGASLNSFVHVVMYSYYGLSAI-PAMRPYLWKKRYITQLQLVQFFLTMSQTMCAVVWPCGFPM 225

615 SsE5a IWWFVMNWVPCGHSYFGASLNSFIHVLMSYYGLSAV-PALRPYLWKKYITQGGLIQFFLTMSQTICAVIWPCGFPR 225

616 SsE5b IWWFVMNWVPCGHSYFGASLNSFVHVLMSYYGLSAV-PAIRPYLWKKYITQGGLIQFFLTMSQTICAVIWPCGFPR 225

617 DrE5 IWWFVMNWVPCGHSYFGATFNSFIHVLMSYYGLSAV-PALRPYLWKKYITQGGLVQFVLTMFQTSCAVVWPCGFPM 225

618 SaE4 LWWIGIKWVPGGQSFFGATINSSIHVLMYGYGLAALGPQM **KYLW**KKYLTIIQMIQFHV TIGHAGHS **SLY**TGCPFPA 229

619 ScE4 LWWIGIKWVPGGQSFFGATINSSIHVLMYGYGLAALGPQM **KYLW**KKYLTIIQMIQFHV TIGHAGHS **SLY**TGCPFPA 229

620 RcE4 LWWIGIKWVPGGQAFFGATINSSIHVLMYGYGLAALGPQM **KYLW**KKYLTIIQMIQFHV TIGHAGHS **SLY**TGCPFPC 229


```

621 NmE4      LWWIGIKWVPGGQSFFGATINSSIHVLMYGYGLAALGPQMQLWKKYLTIIQMIQFHVTTIGHAGHSlyTGCPFFA 229
622 SsE4      LWWIGIKWVPGGQSFFGAGINSSIHVLMYGYGLAAFgPKIQKFLWKKYLTIIQMIQFHVTTIGHAGHSlyTGCPFFA 229
623 DrE4b     LWWIGIKWVPGGQSFFGATINSGIHVLMYGYGLAAFgPKIQKFLWKKYLTIIQMIQFHVTTIGHAAHSlyTGCPFFA 229
624          :*: . :* . *::** *:: **:* ***:*. * :: :***:*:* *::** *:: : : : * . *
625          ----- III ----- IV ----- V
626 SaE5      GWLYLQIGYTitLMIFFLNFYFQTYKKRSPS-RQK----EhMNGSPVSTNGHANGTPSMEHSG-----HKKLrMD 294
627 ScE5      GWLYFQISYMVTLVLFNSFYIQTyKKRSSS-RKT----DHQNGSPLSTNGHANGK---ESAA-----HKKLrVD 291
628 RcE5      GWLYFQISYMVTLIILFSNFYIQTyKKHSGS-LKK----EHQNGSPVSTNGHANGTPSMEYNV-----HKKLrVD 294
629 NmE5      GWLYFQISYMVTLIFLFSNFYVQTYKKHSVS-LKK----EHQNGSPVSPNGHANGTPSLEHAA-----HKKLrVD 294
630 SsE5a     GWLYFQIFyVVTLiAlFSNFYIQTyKKHLVSQKKE----CHQNGSVASLNGHVNGVTPtETIT-----HRKVRGD 295
631 SsE5b     GWLFFQIFyMASLIaFFSNFYIQTyKKHRVS-QKE----YHQNGSVDSLNGHANGVTPtETIT-----HRKVRVD 294
632 DrE5      GWLYFQISYMVTLILLFSNFYIQTyKKRSGS-RKS----DYPNGS---VNGHTNGVMSSEKIK-----HRKARAD 291
633 SaE4      WMQWALIGYAVTFIILFANFYyHAYRRKpSSMQKGDkP--VANGTSTVTNG-HSKVEEVDdNK-KRQKKGRakRE 305
634 ScE4      WMQWALIGYAVTFIILFANFYyHAYRRKpSSGQKGGkP--VTNGTSTVTNG-HSKVEEEE---KRQKKGRakRE 302
635 RcE4      WMQWALIGYAVTFIILFANFYyHAYRGKpSSSQKGGkP--IANGTSVVTNG-HSKVEEVEDNG-KRQKKGRakRE 305
636 NmE4      WMQWALIGYAVTFIILFANFYyHAYRRKpSSAQKGGkP--AVNGTSMVTNG-HSKAEVEEDNG-KRQKKGRakRE 305
637 SsE4      WMQWALIGYAVTFIILFGNFYyQTYRRTPrSAHKVAKP--VTNGVSMATNG-YNKLQDVEENGLKQKKGRakRE 306
638 DrE4b     WMQWALIGYAVTFIILFANFYyQTYRRQPR--LKTAKS--AVNGVSMSTNG-TSKTAeVTENG-KKQKKGKGGkHD 303
639          :... :* * ::: :* *** :*:..          **      **          :. : : :
640          ----- V -----

```


641 Figure 1. Alignment of the deduced amino acid (aa) sequences of elongases Elov15 (E5) and Elov14  
642 (E4) isolated from *Scatophagus argus* (Sa) with their corresponding orthologues, including  
643 rabbitfish (*Siganus canaliculatus*, Sc) Elov15 (ADE34561) and Elov14 (ADZ73580), cobia  
644 (*Rachycentron canadum*, Rc) Elov15 (ACJ65150) and Elov14 (ADG59898), Nibe croaker (*Nibea*  
645 *mitsukurii*, Nm) Elov15 (ACR47973) and Elov14 (AJD80650), Atlantic salmon (*Salmo salar*, Ss)  
646 Elov15a (AAO13175), Elov15b (ACI62499) and Elov14 (ADJ95235), zebrafish (*Danio rerio*, Dr)  
647 Elov15 (NP\_956747) and Elov14b (NP956266). Deduced aa sequences were aligned using  
648 ClustalW2. Identical and similar residues are marked with ‘\*’ and ‘.’, respectively. The conserved  
649 histidine box HXXHH is shaded grey, five putative transmembrane domains are dash-underlined,  
650 and the putative ER retrieval signal is solid underlined.

651  
652  
653  
654  
655  
656  
657



658  
 659 Figure 2. Phylogenetic tree comparing the deduced amino acids of *Scatophagus argus* Elov14 and  
 660 Elov15 with other Elov1 members from fish and mammals. The tree was constructed using the  
 661 neighbor-joining method (Saitou and Nei,1987 ) with MEGA6. The horizontal branch length is  
 662 proportional to the substitution rate per site. Numbers represent the frequencies with which the tree  
 663 topology presented was replicated after 1000 bootstrap iterations.

664  
 665  
 666  
 667  
 668  
 669  
 670  
 671  
 672



673

674 Figure 3 Functional characterization of *Scatophagus argus* putative Elov15 in yeast *Saccharomyces cerevisiae*.

675 FAMES were extracted from yeast transformed with the pYES2-*elov15*, and grown in the presence of PUFA

676 substrates 18:3n-6 (A), 18:4n-3 (B), 20:4n-6(C), 20:5n-3 (D), 22:4n-6 (E) and 22:5n-3 (F). Based on retention

677 times, substrates (\*) and their corresponding elongated products (↓) are indicated accordingly. Peaks 1–4 represent

678 the main endogenous FAs of *S. cerevisiae*, namely 16:0, 16:1 isomers, 18:0 and 18:1n-9, respectively. Vertical

679 axis, FID response; horizontal axis, retention time.

680

681

682

683

684

685

686


687

688

689

690

691



692

693 Figure 4 Functional characterization of *Scatophagus argus* putative Elovl4 in yeast *Saccharomyces cerevisiae*.

694 FAME were extracted from yeast transformed with the pYES2-*elovl4*, and grown in the presence of PUFA

695 substrates 18:3n-6 (A), 18:4n-3 (B), 20:4n-6(C), 20:5n-3 (D), 22:4n-6 (E) and 22:5n-3 (F). Based on retention

696 times, substrates (\*) and their corresponding elongated products (↓) are indicated accordingly. Peaks 1–4 represent

697 the main endogenous FAs of *S. cerevisiae*, namely 16:0, 16:1 isomers, 18:0 and 18:1n-9, respectively. Vertical

698 axis, FID response; horizontal axis, retention time.

699

700

701

702

703

704

705


706

707

708

709

710



711

712 Figure 5. Tissue-specific expression of *elov15* and *elov14* from *S. argus*. Expression of the housekeeping gene *18S*

713 *rRNA* is also shown. NTC: no template control.

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729


730

731

732

733

734



735

736 Figure 6. Relative expression levels of *elovl5* (A) and *elovl4* (B) in livers and eyes collected from *S. argus* fed six  
 737 experimental diets. Expression values were normalized to those of *18S rRNA*. Data are means  $\pm$  SEM (n = 6). Bars  
 738 with different superscripts are significantly different (P < 0.05, one-way ANOVA and Tukey's tests). D2: control  
 739 diet with fish oil as lipid source; D1, D3–D6: diets with blended vegetable oils as lipid source with dietary  
 740 LNA/LA ratios of 0.14, 0.57, 0.84, 1.72, and 2.85, respectively.