

Accepted refereed manuscript of:

Arif M, Ali K, Jan MT, Shah Z, Jones DL & Quilliam R (2016) Integration of biochar with animal manure and nitrogen for improving maize yields and soil properties in calcareous semi-arid agroecosystems, *Field Crops Research*, 195, pp. 28-35.

DOI: [10.1016/j.fcr.2016.05.011](https://doi.org/10.1016/j.fcr.2016.05.011)

© 2016, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

1 **Integration of biochar with animal manure and nitrogen for improving maize yields and**
2 **soil properties in calcareous semi-arid agroecosystems**

3

4 Muhammad Arif^a, Kawsar Ali^a, M.T. Jan^a, Z. Shah^b, Davey L. Jones^c, Richard S. Quilliam^{d*}

5

6 ^a*Department of Agronomy, University of Agriculture, Peshawar, Pakistan*

7 ^b*Department of Soil and Environmental Sciences, University of Agriculture, Peshawar,*

8 *Pakistan*

9 ^c*Environment Centre Wales, Bangor University, Bangor, LL57 2UW, UK*

10 ^d*Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK*

11

12 * Corresponding author: Richard Quilliam

13 Email: Richard.Quilliam@stir.ac.uk

14 Tel: 0044 1786 467769

15 **ABSTRACT**

16 Declining soil quality is commonplace throughout Southern Asia and sustainable strategies
17 are required to reverse this trend to ensure food security for future generations. One
18 potential solution to halt this decline is the implementation of integrated nutrient
19 management whereby inorganic fertilisers are added together with organic wastes. These
20 organic materials, however, are often quickly broken down in soil and provide only a
21 transitory improvement in soil quality. Biochar, which can potentially persist in soil for
22 centuries, may offer a more permanent solution to this problem. To address this, we
23 undertook a 2-year field trial to investigate the interactions between conventional NPK
24 fertilisers, farmyard manure (FYM) and biochar in a maize cropping system. Biochar
25 application to the nutrient poor soil increased maize yields after year one by approximately
26 20% although the yield increase was lower in the second year (ca. 12.5%). Overall, there was
27 little difference in grain yield between the 25 t ha⁻¹ and the 50 t ha⁻¹ biochar treatments. In
28 terms of soil quality, biochar addition increased levels of soil organic carbon, inorganic N, P
29 and base cations and had no detrimental impact on pH and salinity in this calcareous soil.
30 Overall, this field trial demonstrated the potential of biochar to induce short-term benefits
31 in crop yield and soil quality in maize cropping systems although the long-term benefits
32 remain to be quantified. From a management perspective, we also highlight potential
33 conflicts in biochar availability and use, which may limit its adoption by small scale farming
34 systems typical of Southern Asia.

35

36 *Keywords:* calcareous soil; crop production; integrated nutrient management; Pakistan; soil
37 organic matter

38 1. Introduction

39 Progressive declines in soil quality and poor nutrient use efficiency continue to hamper
40 agricultural productivity and food security in many developing countries (Vagen et al., 2005;
41 Jones et al., 2013). These problems are further exacerbated by increasing pressures on
42 agronomic systems posed by increases in human population growth and urbanization,
43 uncertainties in the global climate and the need for agriculture to deliver a range of other
44 ecosystem services in addition to food production (e.g. carbon sequestration, biodiversity,
45 flood risk mitigation, water quality; Lal, 2009). There is therefore an urgent need to redesign
46 agroecosystems to rectify the wide range of inefficiencies that exist in the system including
47 disconnects in nutrient supply, demand and recycling as well as those in water use efficiency
48 (Lal et al., 2013). One potential solution includes the recycling of organic nutrients back to
49 land which can help sustain soil organic matter levels which in turn typically brings about
50 improvements in soil biological functioning, aeration, moisture retention, reduced
51 compaction, pollutant attenuation and nutrient supply (Girmay et al., 2008). The types of
52 organic matter that can be potentially added to soil are diverse ranging from crop residues,
53 green manures, industrial wastes, animal wastes and household waste (Ali et al., 2011;
54 Quilty and Cattle, 2011). However, their addition can have a range of benefits or even
55 negative effects depending on the quality of waste added and the level of contaminants
56 present (Jones and Healey, 2010). It is also likely that synergies may exist between the
57 different organic wastes and thus co-application may represent the best option for
58 maximizing the delivery of a range of ecosystem services.

59 The application of pyrolysed organic matter (biochar) to soils is currently gaining
60 considerable interest worldwide due to its potential to improve soil nutrient retention
61 capacity (through the sorption or stabilisation of nutrient ions), water holding capacity and

62 to sequester carbon in a largely recalcitrant form from decades to possibly thousands of
63 years (Downie et al., 2009; Spokas et al., 2012). Although there is strong economic and
64 social competition from the use of charcoal as a domestic fuel source (Maes and Verbist,
65 2012), there is no doubt that it is applicable for use in arable systems where it can be readily
66 incorporated into soil. However, before we can advocate the wide-scale adoption of biochar
67 to resource poor farmers in developing countries, we must first provide the evidence base
68 to show that it is beneficial in both agronomic and economic terms. A number of studies
69 have reported positive effects of biochar amendments on maize yields and soil properties
70 (Cornelissen et al., 2013; Zhang et al., 2012), whilst others have reported no net effect
71 (Jones et al., 2012) suggesting that the response may be to some extent specific to
72 particular environmental conditions and soil types, or agronomic practices, e.g. differences
73 in crop cultivar or fertiliser and pesticide applications. Compared to biochar research in the
74 temperate soils of Europe and North America, relatively little work has been undertaken on
75 the potential use of biochar and its effects on the behaviour of organic and inorganic
76 nutrients in semi-arid regions of the world where improvements in soil quality and food
77 security remain critical. Although there are a growing number of studies investigating the
78 effect of biochar application to tropical soils, many of these focus on acidic soils and the
79 liming effect of biochar (Major et al., 2010). Subsequently, there is a significant lack of data
80 on biochar amendment of agronomic calcareous soils in semi-arid areas such as regions of
81 northern Pakistan.

82 As the supply of fertilizers in Pakistan is limited by a range of socioeconomic, political
83 and geographical constraints, alternative sustainable strategies are required to optimize
84 fertiliser integration (Gandah et al., 2003; Schlecht et al., 2006). Low fertilizer-use-efficiency
85 and losses to the environment, e.g. through leaching, are major environmental problems

86 both in Pakistan and globally, and there is an urgent need for research that aims to improve
87 fundamental efficiencies of crop nutrient use (Tilman et al., 2002; Sanchez, 2002; Arif et al.,
88 2015). The aim of the present study was therefore to determine the effectiveness of
89 biochar, farmyard manure (FYM) and mineral nitrogen alone and in various combinations on
90 aspects of crop yield and soil quality in maize cropping systems. Maize was chosen as the
91 trial crop as it contributes >10% of the total agricultural produce and 15% of agricultural
92 employment in Pakistan, the major share of which (over 50%) originates from small land-
93 holding farmers, who produce mostly for their own food needs (FAO, 2014). Within these
94 farming systems, the intrinsically low fertility of the soil and increasing prices of chemical
95 fertilizers represent the major constraints to increasing maize yields (Khan and Shah, 2011).
96 The need to simultaneously increase yields, decrease production costs and maintain soil
97 health has therefore become a major challenge in semi-arid agroecosystems (Anjum et al.,
98 2010).

99

100 **2. Materials and methods**

101 *2.1. Experimental site*

102 The trial site was located at the New Developmental Farm of the University of
103 Agriculture, Peshawar (34°1'21"N, 71°28'5"E) and the experiment was started in the
104 summer of 2011. The site has a warm to hot, semi-arid, sub-tropical, continental climate
105 with mean annual rainfall of 360 mm. Summer (May–September) has a mean maximum
106 temperature of 40°C and mean minimum temperature of 25°C. Winter (December to the
107 end of March) has mean minimum temperature of 4°C and a maximum of 18.4°C. The
108 average winter rainfall is higher than that of the summer. The highest winter rainfall has
109 been recorded in March, while the highest summer rainfall is in August. %. The soil is a silty

110 clay loam, well drained and strongly calcareous (pH 8.23 \pm 0.09), with an electrical
111 conductivity (EC) of 166 \pm 28.5 μ S cm⁻¹ and an organic matter content of less than 1%. The
112 soil is deficient in nitrogen (23.72 \pm 1.75 mg kg⁻¹) and phosphorus (3.20 \pm 0.50 mg kg⁻¹) but
113 has adequate potassium (85.80 \pm 6.56 mg kg⁻¹).

114

115 2.2. *Experimental design*

116 The study consisted of three levels of biochar (0, 25 and 50 t ha⁻¹), two levels of FYM
117 (5 and 10 t ha⁻¹) and two levels of fertilizer-N (urea) (75 and 150 kg ha⁻¹) together with a
118 control treatment (no biochar, FYM or fertilizer-N). A summary of the treatments and their
119 abbreviations are provided in Table 1. Biochar and FYM were applied at the time of sowing
120 at the beginning of year 1, and reflected typical FYM doses for the region. Half of the
121 fertilizer-N was applied at sowing and the remaining half applied at the 8 leaf stage (V8).
122 Single super phosphate (SSP) was applied at the rate of 90 kg ha⁻¹ as a basal dose. Dairy
123 cattle FYM was obtained from the Peshawar University of Agriculture dairy farm and the
124 biochar was produced from *Acacia* (e.g. *A. nilotica* (Linn.) Delile) using traditional methods
125 employed in the region (Amur and Bhattacharya, 1999). No commercial biochar production
126 takes place in the Khyber Pakhtunkhwa region of Pakistan; however, a limited amount is
127 produced domestically using small biochar furnaces. The biochar was prepared in an
128 enclosed dome shaped room, with several small holes made in the roof which were sealed
129 after about 12 h burning. The feedstock was composed of cuttings from the main stem and
130 branches of > 3 y old *Acacia* trees with a trunk diameter greater than 15 cm. The highest
131 temperature reached during pyrolysis was between 400 to 500 °C, and the final ash content
132 of the biochar was 27 %. Characteristics of the FYM and biochar are shown in Table 2.

133 The experiment had four replicates per treatment, and was laid out in a randomized

134 complete block design. The treatment plots were 4.0 m x 4.5 m in size with strong ridges
135 placed around each plot for delineation and to prevent biochar migration. Between row and
136 within row distance was 75 cm and 20 cm, respectively. The field was ploughed twice down
137 to a depth of 30 cm, followed by planking to break the clods and level the field taking care
138 not to disturb the ridges and to facilitate biochar movement from one plot to another.
139 Biochar was crushed and sieved to pass 2 cm, spread uniformly on the surface of the soil of
140 each sub plot and then ploughed-in with a rotivator, which thoroughly mixed the biochar
141 into the soil surface to a depth of about 15 cm. Maize (*Zea mays* L.) cv. 'Azam' (Cereal Crops
142 Research Institute, Nowshera, Pakistan) was sown at a rate of 30 kg ha⁻¹ on July 1st, 2011
143 and thinned about 15 days after emergence to maintain plant to plant distance of 20 cm and
144 a density of 60,000 to 70,000 plants ha⁻¹. The crop was irrigated ten days after sowing and
145 then again usually every 15 days with adjustment according to rainfall. The crop was
146 specifically irrigated at the critical growth stages of tasseling, silking, cob and grain
147 development. The volume of water applied during irrigation was 340 m³ per ha⁻¹. Weeds
148 were controlled manually by hoeing between the ridges with a blade digger about 20 days
149 post emergence. Pesticides were applied at the eight leaf stage (Lorsban® 40EC-
150 (Chlorpyrifos, OP at 5 ml l⁻¹) to protect against stem borer.

151

152 2.3. Crop harvest

153 At harvest (Oct 1st, 2011), the following maize yield components were recorded:
154 total aboveground biomass, grain yield, number of ears m⁻², number of grains per ear and
155 the thousand grain weight. To determine total above-ground yield (t ha⁻¹), the plants from
156 the four central rows in each plot were harvested, sun dried (until constant weight) and
157 weighed. The ears from these harvested plants were then removed, threshed and grain

158 yield (t ha^{-1}) calculated. Ears were counted in the four central rows of the standing maize
159 crop in each plot. Thousand grain weight was calculated from a sub-sample from of each
160 plot.

161

162 *2.4. Soil quality analysis*

163 Three replicate soil samples were taken from 0-15 cm depth within a week of
164 harvest. Soil carbon was determined by the Walkley-Black procedure (Nelson and Sommers,
165 1996). Carbonates were not removed before soil C determination, but an excess amount of
166 dichromates was used to oxidize all possible organic C. Total mineral N in the soil samples
167 was determined after KCl extraction by the steam distillation method as described in
168 Mulvaney (1996). Soil pH and EC were measured in a saturated soil-water (1:1 w:v) paste
169 extract under vacuum (Rhoades, 1996), using a pH meter (InoLab pH 720, WTW Series,
170 Germany) and an EC meter (EC Meter 4510, Jenway, UK). Plant-available P and K in soil were
171 determined in an ammonium bicarbonate-DTPA extract (1 M NH_4HCO_3 , 0.005 M DTPA; pH
172 7.6) either colorimetrically (P) or by flame photometry (K) according to the procedure
173 outlined in Soltanpour and Schwab (1977). Ca and Mg were determined in the saturation
174 paste extracts by Atomic Absorption Spectrophotometry (Model 2380, Perkin Elmer Corp.,
175 Waltham, MA, USA).

176

177 *2.5. Statistical analysis*

178 Differences between each treatment (biochar, FYM and N fertiliser) in each year were
179 compared by analysis of variance (three-way ANOVA) for each yield and soil quality
180 parameter. The difference between year 1 and year 2 for yield and each soil quality
181 parameter was compared by Student's t-test (Minitab 12.0 software, Minitab Inc., PA, USA).

182 3. Results

183 3.1. Yield and yield components

184 The addition of FYM and N fertiliser significantly increased the yield of maize
185 compared to the unamended control plots (Fig. 1; Tables 3 and 4). Biochar application
186 significantly increased the grain yield in both years ($P < 0.001$), although there was little
187 difference in grain yield between the 25 t ha⁻¹ and the 50 t ha⁻¹ biochar treatments (Fig. 1;
188 Tables 3 and 4). Biological yield was significantly higher in both years in plots treated with
189 biochar, although the number of grains per ear was only higher in the first year ($P < 0.001$)
190 and an increase in the thousand grain weight was only significantly higher in the second
191 year (Table 5). The addition of FYM in the treated plots made no significant difference to
192 grain yield in either year (Table 5), although it did significantly increase the grains per ear,
193 the thousand grain weight and the biological yield in year 1. Nitrogen fertiliser significantly
194 increased the grain yield and grains per ear in the first year ($P < 0.001$), but this was not
195 repeated in the second year (Table 5). Two-way interactions between the biochar, FYM and
196 the N fertiliser significantly increased grain yield in the first year ($P < 0.05$), but not the
197 second year (Table 5), when there was no significant interaction between all three
198 treatments on any of the yield parameters measured.

199

200 3.2. Soil properties

201 Overall, the addition of biochar made a significant difference to soil quality
202 parameters in both cropping cycles (Table 6). There was a significant increase in soil pH ($P <$
203 0.05) following biochar application, i.e. 7.18 ± 0.11 ; 7.43 ± 0.10 ; 7.65 ± 0.20 for 0, 25 and 50 t
204 ha⁻¹ biochar addition respectively (data from both cropping cycles combined).

205 By year 2, soil organic carbon was significantly higher ($P < 0.05$) in plots amended

206 with biochar in year 1 (Tables 3 and 4;), with between 40 – 75 % more soil organic carbon in
207 the plots containing 50 t ha⁻¹ biochar compared to the plots containing 25 t ha⁻¹ (Fig. 2a).
208 Soil mineral N remained at a similar concentration from year 1 to year 2 for each treatment
209 (Fig. 2b), and was not affected by the rate of N fertiliser that had been applied (half rate, 75
210 kg ha⁻¹ or full rate, 150kg ha⁻¹). Although the concentration of soil N after the second year
211 was significantly higher in plots amended with biochar at both 25 and 50 t ha⁻¹ compared to
212 the unamended plots (Table 4), overall, there was no significant interaction between
213 biochar and the application of N fertiliser (Table 6). The addition of biochar at both rates
214 increased the concentration of soil P in the first year (Fig. 2c; Table 3). In the 50 t ha⁻¹
215 biochar plots there was significantly more soil P compared to the plots containing 25 t ha⁻¹
216 ($P < 0.01$), and in the plots with 50 t ha⁻¹ biochar the highest concentration of soil P was
217 coupled with the full rate of FYM (Table 6). By year 2 however, in the biochar-amended
218 plots the concentration of soil P had significantly declined ($P < 0.01$) compared to the
219 concentration in year 1 (Fig 2c). In contrast, the increase in soil Ca/Mg was significantly
220 higher after year 2 in plots amended with 50 t ha⁻¹ biochar (Fig. 3a). Although there was a
221 significant interaction effect between biochar, the FYM and the N (either singly or in
222 combination with biochar) in year 1; by year 2 the concentration of soil Ca/Mg was not
223 affected by either organic or inorganic fertilisers (Table 6). For K, the application of FYM and
224 inorganic N fertiliser to the non-biochar-amended soil was no different to the control soil
225 which contained neither fertiliser nor biochar (Fig. 3b), although there were significantly
226 higher levels after the second year ($P < 0.01$). The application of 50 t ha⁻¹ biochar
227 significantly increased the concentration of K in the soil (Fig. 3b); particularly in the first year
228 (Table 3) when there was a significant interaction between the biochar and the FYM and the
229 N fertiliser (Table 6). Consequently, the effect of an increased concentration of ions in the

230 biochar-amended soil generated a significant increase in soil EC (Fig. 3c) in both year 1 and
231 year 2 (Tables 3 and 4).

232

233 **4. Discussion**

234 There is a significant lack of data on biochar amendment of agronomic calcareous soils in
235 semi-arid areas such as regions of northern Pakistan, but this study has shown that the
236 application of FYM and synthetic N in combination with biochar had an overall positive
237 effect on soil properties, and increased maize yield in the first year after application. While
238 the short term impacts of biochar application are becoming clearer for temperate
239 agricultural soils, we absolutely lack an adequate understanding of the longer-term impacts
240 and implications of biochar use in the cereal cropping systems commonly used in South
241 Asia. Following biochar application to temperate soils an initial transient flush of labile
242 compounds into the rhizosphere can enhance nutrient cycling and increase crop yield
243 (Quilliam et al., 2012). Similarly, biochar application to the nutrient poor soils of Pakistan
244 used in these field trials increased maize yields after year one by approximately 20%
245 although this magnitude of yield increase was not replicated in the second year, and the
246 potential benefits of biochar addition to this semi-arid calcareous agricultural soil appears
247 to be short term or transient.

248 In tropical acidic soils, biochar application can have a liming effect which is often
249 associated with increased nutrient availability, e.g. phosphorus, and ultimately improved
250 crop yield. Applying biochar to the alkaline soils used in this study increased the pH from
251 7.18 to 7.43 and 7.63 respectively for the two biochar applications, which may have
252 influenced the availability of some soil nutrients. In applied terms however, the increase of
253 0.30 to 0.45 pH units probably made little difference to the availability of soil nutrients at

254 this near neutral pH. None of the nutrients we measured decreased with the increasing pH,
255 and as the total yield was not negatively affected our data also suggests that the increasing
256 pH did not facilitate plant toxicity of any other soil nutrients.

257 Biochar application to agricultural soil can facilitate the sorption or stabilisation of
258 solutes and nutrient ions, and reduce nutrient loss from leaching (Asai et al., 2009; Laird et
259 al., 2010), and the maintenance of elevated levels of soil P and N after the second year
260 harvest suggests that biochar can mediate the slow release of these nutrients (Mukherjee
261 and Zimmerman, 2013). Depending on pyrolysis conditions, the total surface area and pore
262 volume of biochar can be orders of magnitude greater than soil (Calvelo Pereira et al., 2011;
263 Quilliam et al., 2013). Subsequently, biochar can provide multiple planar sites to strongly
264 sorb soil mineral and organic compounds (Joseph et al., 2010), although cation exchange
265 capacity and the hydrophobicity of the biochar surface can also significantly affect its
266 sorptive ability (Pignatello, 2013). Absorption of nutrients contained within the inorganic N
267 fertiliser and the FYM onto the surface of the biochar would effectively reduce
268 bioavailability for microbial utilisation and prevent bound nutrients from being leached
269 away following rainfall or irrigation and may reduce volatilization of NH₃.

270 After the second year, the biochar amended plots (at both application rates) had higher
271 concentrations of P and N. Therefore, these macronutrients are not being retained in the
272 soil for as long when applied in just a mix of FYM and synthetic N compared with when they
273 were applied in tandem with biochar. As the yield was higher (or no different) in the
274 biochar-amended soil compared to the soil containing the FYM and N, it is not plant uptake
275 and subsequent harvest that is removing these nutrients in the non biochar-amended soils.
276 Reports from tropical acidic soil show that biochar can bind nutrients to its surface, which
277 allows them to remain in the soil for longer, e.g. not being leached away after a single

278 cropping season, and despite the higher pH of the calcareous soil used in this study, our
279 results also suggest that biochar can retain nutrients such as P and N. Over time, these
280 nutrients will slowly be released back into the soil resulting in a more sustainable use of the
281 farmer's original investment in synthetic fertiliser (Asai et al., 2009). In addition to the
282 increased efficiency of nutrient input, incorporating biochar into agroecosystems has the
283 potential to enhance wider ecosystem service delivery, for example, by reducing nutrient
284 and pesticide mobilisation and transfer from soil into aquatic systems (Jeffery et al., 2013).

285 For this study we have applied fairly high rates of biochar in order to clearly
286 demarcate potential differences between our treatments; however, there are also recent
287 reports of lower biochar application rates being beneficial in calcareous soils (Zhang et al.,
288 2012; Ippolito et al., 2014). To produce such high quantities of biochar requires large
289 volumes of feedstock, and there is justifiable concern about the implications of
290 overharvesting existing forests for biochar production, as progressive deforestation in semi-
291 arid ecosystems has already led to the deterioration of a range of ecosystem services. In
292 Pakistan, nearly 62% of the population live in rural areas and are reliant on agriculture for
293 their livelihoods. Consequently, there is a significant dependence on fuelwood as a source of
294 energy, and in a country that already has low forest cover (of about 4.80%), the high
295 consumption rate of fuelwood per household per day (6.70 kg) is contributing to the
296 unsustainable use of the country's wood resources (Butt et al., 2013). In the rain-fed areas
297 of Pakistan, e.g. the southern districts of Khyber Pakhtunkhwa, wild-growing Acacia is
298 already seasonally pruned to make charcoal; however, any potential benefits of biochar
299 application to agricultural soil are accompanied by some important trade-offs, such as the
300 potential for deforestation and land degradation (Anjum et al., 2010), together with the
301 behavioural and cultural implications associated with using a primary source of fuel as a soil

302 amendment (Maes and Verbist, 2012).

303 Environmental degradation in semi-arid regions, as a consequence of biochar
304 production, is obviously not a sustainable strategy for improving soil nutrient use efficiency
305 and delivering increased food security (Woolf et al., 2010). However, biochar can be
306 produced from any organic material, and the pyrolysis of non-virgin feedstocks would allow
307 the production of significant volumes of biochar without exacerbating the existing pressures
308 on forest resources. Whilst there is the potential to produce biochar from 'on-farm' organic
309 wastes, e.g. stover or maize cobs, in semi-arid agricultural systems much of this 'waste'
310 biomass is already fully utilised, for example as animal feed, mulch or for constructing
311 fences and roofs. Thus, short-term cycling of these streams of organic matter back through
312 the agricultural chain is probably more beneficial than taking them out of the loop by
313 converting them into biochar (Jones et al., 2013).

314 Our results have demonstrated that the integration of biochar with inorganic N
315 fertiliser and FYM application at the field-scale can improve the productivity of maize and
316 could provide a more sustainable input of N and P to soil. The soil used in this study has low
317 levels of organic matter (Arif et al., 2015) therefore, augmenting the soil organic matter
318 content with FYM can also promote nutrient cycling and the water holding capacity, and
319 adding biochar to soil in Pakistan could improve yield responses to inorganic N and P
320 fertilizers. For resource-poor farmers living with soil of intrinsically low fertility, the cost and
321 availability of chemical fertilizers is often the most prohibitive constraint to increasing crop
322 yields; therefore the sustainable management of nutrients is critical for maximising the
323 efficiency of crop nutrient use. Incorporating FYM and biochar into an integrated nutrient
324 management regime could be an important strategy for improving the overall farm
325 productivity of cereal-based cropping systems in Pakistan. However, this needs critical

326 evaluation in a sustainable agricultural context. Central to this are participatory-based
327 approaches to assess whether biochar can really make a practical contribution to agriculture
328 in Pakistan by providing farmers with a sustainable solution to help alleviate the constraints
329 driven by poor soil fertility (Arif et al., 2015). Crucially, an evaluation of the wider ecosystem
330 services linked to the trade-offs associated with producing biochar in semi-arid ecosystems
331 needs both careful consideration and robust evidence before it can be promoted as a
332 sustainable option for optimising fertiliser use efficiency.

333

334 **Acknowledgments**

335 Funding for this project was provided by The Higher Education Commission, Pakistan and
336 the Welsh European Funding Office under the SEREN programme.

337

338 **References**

339 Ali, K., Munsif, F., Zubair, M., Hussain, Z., Shahid, M., Din, U.I., Khan, N., 2011. Management
340 of organic and inorganic nitrogen for different maize varieties. *Sarhad Journal of*
341 *Agriculture* 27, 525-529.

342 Amur, G.Q., Bhattacharya, S.C., 1999. A study of biomass as a source of energy in Pakistan.
343 *International Energy Journal* 21, 25-36.

344 Anjum, S.A., Wang, L.C., Xue, L.L., Saleem, M.F., Wang, G.X., Zou, C.M., 2010. Desertification
345 in Pakistan: causes, impacts and management. *Journal of Food Agriculture and*
346 *Environment* 8, 1203-1208.

347 Arif M., Jalal, F., Jan, M.T., Muhammad, D., Quilliam, R.S., (2015). Incorporation of biochar
348 and legumes into the summer gap: improving productivity of cereal-based cropping
349 systems in Pakistan. *Agroecology and Sustainable Food Systems* 39, 391-398.

350 Asai, H., Samson, B. K., Stephan, H. M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., Horie,
351 T., 2009. Biochar amendment techniques for upland rice production in Northern
352 Laos: 1. Soil physical properties, leaf SPAD & grain yield. *Field Crops Research* 111,
353 81-84.

354 Butt, S., Hartmann, I., Lenz, V., 2013. Bioenergy potential and consumption in Pakistan.
355 *Biomass and Bioenergy* 58, 379-389.

356 Calvelo Pereira, R., Kaal, J., Camps Arbestain, M., Pardo Lorenzo, R., Aitkenhead, W., Hedley,
357 M., Macías, F., Hindmarsh, J., Maciá-Agulló, J.A., 2011. Contribution to
358 characterisation of biochar to estimate the labile fraction of carbon. *Organic*
359 *Geochemistry* 42, 1331-1342.

360 Cornelissen, G., Martinsen, V., Shitumbanuma, V., Alling, V., Breedveld, G.D., Rutherford,
361 D.W., Sparrevik, M., Hale, S.E., Obia, A. Mulder, J., 2013. Biochar effect on maize
362 yield and soil characteristics in five conservation farming sites in Zambia. *Agronomy*
363 3, 256-274.

364 Downie, A., Crosky, A. Munroe, P., 2009. Physical properties of biochar. In 'Biochar for
365 Environmental Management: Science and Technology'. Eds J. Lehmann, S. Joseph.
366 pp. 13-32. Earthscan, London, UK.

367 FAO, 2014. Pakistan. www.fao.org/countryprofiles/index/en/?iso3=PAK (accessed May 14th,
368 2014).

369 Gandah, M., Bouma, J., Brouwer, J., Hiernaux, P., van Duivenbooden, N., 2003. Strategies to
370 optimize allocation of limited nutrients to sandy soils of the Sahel: a case study from
371 Niger, West Africa. *Agriculture Ecosystems and Environment* 94, 311-319.

372 Girmay, G., Singh, B.R., Mitiku, H., Borresen, T., Lal, R., 2008. Carbon stocks in Ethiopian soils in
373 relation to land use and soil management. *Land Degradation & Development* 19, 351-367.

374 Ippolito, J.A., Stromberger, M.E., Lentz, R.D., Dungan, R.S., 2014. Hardwood biochar
375 influences calcareous soil physicochemical and microbiological status. *Journal of*
376 *Environmental Quality* 43, 681-689.

377 Jeffery, S., Bezemer, T.M., Cornelissen, G., Kuyper, T.H.W., Lehmann, J., Mom-mer, L., Sohi,
378 S., van de Voorde, T.F.J., Wardle, D.A. and van Groenigen, J.W., 2015. The way
379 forward in biochar research: targeting trade-offs between the potential wins. *Global*
380 *Change Biology: Bioenergy* 7, 1-13.

381 Jones, D.L., Healey, J.R., 2010. Organic amendments for remediation: putting waste to good
382 use. *Elements* 6, 369-374.

383 Jones, D.L., Rousk, J., Edwards-Jones, G., DeLuca, T.H., Murphy, D.V., 2012. Biochar-
384 mediated changes in soil quality and plant growth in a three year field trial. *Soil*
385 *Biology & Biochemistry* 45, 113-124.

386 Jones, D.L., Cross, P., Withers, P.J., DeLuca, T.H., Robinson, D.A., Quilliam, R.S., Harris I,
387 Chadwick D.R., Edwards-Jones, G., 2013. Nutrient stripping: the global disparity
388 between food security and soil nutrient stocks. *Journal of Applied Ecology* 50, 851-
389 862.

390 Joseph, S.D., Camps-Arbestain, M., Lin, Y., Munroe, P., Chia, C.H., Hook, J., van Zwieten, L.,
391 Kimber, S., Cowie, A., Singh, B.P., Lehmann, J., Foidl, N., Smernik, R.J., Amonette, J.E.,
392 2010. An investigation into the reactions of biochar in soil. *Australian Journal of Soil*
393 *Research* 48, 501-515.

394 Khan, M.A., Shah, S.A.A., 2011. Food insecurity in Pakistan: causes and policy response.
395 *Journal of Agricultural and Environmental Ethics* 24, 493-509.

396 Laird, D.A., Fleming, P., Davis, D.D., Horton, R., Wang, B., Karlen, D.L., 2010. Impact of
397 biochar amendments on the quality of a typical Midwestern agricultural soil.

398 Geoderma 158, 443-449.

399 Lal, R., 2009. Soils and food sufficiency: a review. *Agronomy and Sustainable Development*
400 29, 113-118

401 Lal, R., 2013. Food security in a changing climate. *Ecohydrology and Hydrobiology* 13, 8-21.

402 Lentz, R.D., Ippolito, J.A., 2012. Biochar and manure affect calcareous soil and corn silage
403 nutrient concentrations and uptake. *Journal of Environmental Quality* 41, 1033-1043.

404 Maes, W.H., Verbist, B., 2012. Increasing the sustainability of household cooking in
405 developing countries: policy implications. *Renewable and Sustainable Energy*
406 *Reviews* 16, 4204-4221.

407 Major, J., Rondon, M., Molina, D., Riha, S.J. Lehmann, J., 2010. Maize yield and nutrition
408 during 4 years after biochar application to a Colombian savanna oxisol. *Plant & Soil*
409 333, 117-128.

410 Mukherjee, A., Zimmerman, A.R. 2013. Organic carbon and nutrient release from a range of
411 laboratory-produced biochars and biochar-soil mixtures. *Geoderma*, 193, 122-130.

412 Mulvaney, R.L., 1996. Nitrogen – Inorganic forms. p. 1123-1184. In D. L. Sparks et al. (ed.)
413 *Methods of soil analysis. Part 2. Chemical properties. SSSA Book Series 5., Soil*
414 *Science Society of America, Madison, WI, USA.*

415 Nelson, D.W., Sommers, L.E., 1996. Total carbon, organic carbon, and organic matter. In:
416 *Methods of Soil Analysis, Part 2, 2nd ed., A.L. Page et al., Ed. Agronomy. 9:961-1010.*
417 *American Society of Agronomy Inc., Madison, WI, USA.*

418 Pignatello, J.J., 2013. Adsorption of Dissolved Organic Compounds by Black Carbon. In: Xu, J.,
419 Sparks, D.L. (Eds.), *Molecular Environmental Soil Science. Springer, Dordrecht, The*
420 *Netherlands, pp. 359-385.*

421 Quilliam, R.S., Marsden, K., Gertler, G., Rousk, J., DeLuca, T.H., Jones, D.L., 2012. Nutrient

422 dynamics, microbial growth and weed emergence in biochar amended soil are
423 influenced by time since application and reapplication rate. *Agriculture, Ecosystems*
424 *and Environment* 158, 192-199.

425 Quilliam, R.S., Glanville, H.C., Wade, S.C., Jones, D.L., 2013. Life in the 'charosphere' – does
426 biochar in agricultural soil provide a significant habitat for microorganisms? *Soil*
427 *Biology & Biochemistry* 65, 287-293.

428 Quilty, J.R., Cattle, S.R., 2011. Use and understanding of organic amendments in Australian
429 agriculture: a review. *Soil Research* 49, 1-26.

430 Rhoades, J.D., 1996. Salinity: electrical conductivity and total dissolved solids. In *Methods of*
431 *Soil Analysis. Part 3. Chemical Methods. 3rd Ed.*, D.L. Sparks., Ed., pp 417-435.
432 American Society of Agronomy Inc. Madison, WI. and Soil Science Society of America,
433 Madison, WI,

434 Sanchez, P.A., 2002. Soil fertility and hunger in Africa. *Science* 295, 2019-2020.

435 Schlecht, E., Buerkert, A., Tielkes, E., Bationo, A., 2006. A critical analysis of challenges and
436 opportunities for soil fertility restoration in Sudano-Sahelian West Africa. *Nutrient*
437 *Cycling in Agroecosystems* 76, 109-136.

438 Soltanpour, P.N., Schwab, A.P., 1977. A new soil test for simultaneous extraction of macro-
439 and micro-nutrients in alkaline soils. *Communications in Soil Science and Plant*
440 *Analysis* 8, 195–207.

441 Spokas, K.A., Cantrell, K.B., Novak, J.M., Archer, D.W., Ippolito, J.A., Collins, H.P., Boateng,
442 A.A., Lima, I.M., Lamb, M.C., McAloon, A.J., Lentz, R.D., Nichols, K.A., 2012. Biochar: a
443 synthesis of its agronomic impact beyond carbon sequestration. *Journal of*
444 *Environmental Quality* 41, 973-989.

445 Vagen, T.G., Lal, R., Singh, B.R., 2005. Soil carbon sequestration in sub-Saharan Africa: A

446 review. *Land Degradation and Development* 16, 53-71

447 Whalen, J.K., Chang, C., Clayton, G.W., Carefoot, J.P., 2000. Cattle manure amendments can
448 increase the pH of acid soils. *Soil Science Society of America Journal* 64, 962-966.


449 Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural
450 sustainability and intensive production practices. *Nature* 418, 671-677.

451 Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., Joseph, S., 2010. Sustainable
452 biochar to mitigate global climate change. *Nature Communications* 1, 56.


453 Zhang, A., Liu, Y., Pan, G., Hussain, Q., Li, L., Zheng, J., Zhang, X., 2011. Effect of biochar
454 amendment on maize yield and greenhouse gas emissions from a soil organic carbon
455 poor calcareous loamy soil from Central China Plain. *Plant Soil* 351, 263–275.

456 Zingore, S., Delve, R.J., Nyamangara, J., Giller, K.E., 2008. Multiple benefits of manure: The
457 key to maintenance of soil fertility and restoration of depleted sandy soils on African
458 smallholder farms. *Nutrient Cycling in Agroecosystems* 80, 267-282.

459




471 **Fig. 1.** Yield of maize in year 1 (grey bars) and year 2 (black bars) fertilised with FYM at either
 472 5 t ha⁻¹ (half manure; HM) or 10 t ha⁻¹ (full manure; FM) and N fertiliser, at either 75 kg ha⁻¹
 473 (half fertiliser; HF) or 150 kg ha⁻¹ (full fertiliser; FF). All plots were amended with biochar at
 474 the application rates of 0, 25 or 50 t ha⁻¹. Control, 0 t ha⁻¹ FYM, 0 kg ha⁻¹ N fertiliser, and 0 t
 475 ha⁻¹ biochar. Asterisks indicate a significant difference between year 1 and year 2 data for
 476 each treatment at the *P < 0.05, **P < 0.01 and ***P < 0.001 level (T-test). Data points
 477 represent the mean of three replicates +SE.



478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493

494 **Fig. 2.** Soil organic carbon (a), mineral nitrogen (b) and extractable phosphorus following the
 495 harvest of maize in year 1 (grey bars) and year 2 (black bars). Plots had been fertilised with
 496 FYM at either 5 t ha⁻¹ (half manure; HM) or 10 t ha⁻¹ (full manure; FM) and N fertiliser, at
 497 either 75 kg ha⁻¹ (half fertiliser; HF) or 150 kg ha⁻¹ (full fertiliser; FF). All plots were amended
 498 with biochar at the application rates of 0, 25 or 50 t ha⁻¹. Control, 0 t ha⁻¹ FYM, 0 kg ha⁻¹ N
 499 fertiliser, and 0 t ha⁻¹ biochar. Asterisks indicate a significant difference between year 1 and
 500 year 2 data for each treatment at the *P < 0.05, **P < 0.01 and ***P < 0.001 level (T-test).
 501 Data points represent the mean of three replicates +SE.



518 **Fig. 3.** Soil extractable Ca/Mg (a), extractable potassium (b) and soil electrical conductivity
 519 following the harvest of maize in year 1 (grey bars) and year 2 (black bars). Plots had been
 520 fertilised with FYM at either 5 t ha⁻¹ (half manure; HM) or 10 t ha⁻¹ (full manure; FM) and N
 521 fertiliser, at either 75 kg ha⁻¹ (half fertiliser; HF) or 150 kg ha⁻¹ (full fertiliser; FF). Plots were
 522 amended with biochar at the application rates of 0, 25 or 50 t ha⁻¹. Control, 0 t ha⁻¹ FYM, 0
 523 kg ha⁻¹ N fertiliser, and 0 t ha⁻¹ biochar. Asterisks indicate a significant difference between
 524 year 1 and year 2 data for each treatment at the *P < 0.05, **P < 0.01 and ***P < 0.001
 525 level (T-test). Data points represent the mean of three replicates +SE.

526 **Table 1:**

527 Description of treatment combinations used for each replicated ($n = 3$) experimental plot.

Biochar (t ha ⁻¹)	FYM (t ha ⁻¹)	Fertiliser N (kg ha ⁻¹)	Abbreviation ^a
0	0	0	Control
0	5	75	B0-HM-HF
0	5	150	B0-HM-FF
0	10	75	B0-FM-HF
0	10	150	B0-FM-FF
25	5	75	B25-HM-HF
25	5	150	B25-HM-FF
25	10	75	B25-FM-HF
25	10	150	B25-FM-FF
50	5	75	B50-HM-HF
50	5	150	B50-HM-FF
50	10	75	B50-FM-HF
50	10	150	B50-FM-FF

528 ^aHM, half manure rate (5 t ha⁻¹); FM, full manure rate (10 t ha⁻¹);

529 HF, half fertiliser rate (75 t ha⁻¹); FF, full fertiliser rate (150 t ha⁻¹)

530

531 **Table 2:**

532 Chemical properties of the fresh biochar and Farmyard manure (FYM) prior to application to
533 soil.

	Biochar	Farmyard manure
pH	7.01	8.65
EC (dS m ⁻¹) ^a	1.57	2.44
C (g kg ⁻¹)	578	486
P (g kg ⁻¹)	11.4	35.2
N (g kg ⁻¹)	10.2	15.6
Ca (g kg ⁻¹)	2.68	1.86
Mg (mg kg ⁻¹)	10.0	112.6

534 ^aEC, electrical conductivity
535

536 **Table 3:** Multiple pairwise comparisons (Tukey’s HSD) for each treatment for year 1 data

	Yield	Soil C	Soil N	Soil P	Ca/Mg	Soil K	Soil EC
Control	e	b	a	d	g	e	538
B0-HM-HF	d	b	a	d	f	e	539
B0-HM+FF	b,c,d	a,b	a	d	f	e	540
B0-M+HF	d	b	a	d	d,e	d	541
B0-M+FF	a	b	a	d	e,f	e	542
B25-HM-HF	a,b,c	b	a	c	d	c	543
B25-HM+FF	a,b,c	b	a	c	d	c	544
B25-M+HF	c,d	a,b	a	c	b,c,d	c	545
B25-M+FF	a,b	a,b	a	c	c,d	c	546
B50-HM-HF	a,b,c	a,b	a	b	b,c	b	547
B50-HM+FF	a,b,c	a,b	a	b	b,c	a	548
B50-M+HF	a,b,c	a	a	a	b	a	549
B50-M+FF	a,b,c	a	a	a	a	a	550
							551
							552
							553

554

555 Different letters within the same column indicates that the mean significantly differs from
 556 each other (one-way ANOVA, $P < 0.001$; Tukey multiple comparison test, $P < 0.05$).

557

558

559

560

561

562 **Table 4:** Multiple pairwise comparisons (Tukey’s HSD) for each treatment for year 2 data

	Yield	Soil C	Soil N	Soil P	Ca/Mg	Soil K	Soil S
Control	e	f	e	f	e	e	e
B0-HM-HF	d	f	e	d,e	e	c,d,e	d
B0-HM+FF	b,c,d	f	e	d,e	d,e	d,e	d
B0-M+HF	c,d	e,f	d,e	e	d,e	c,d,e	d
B0-M+FF	c,d	d,e,f	c,d	c,d,e	c,d,e	b,c,d,e	d
B25-HM-HF	a,b,c,d	d,e	a,b	c,d,e	b,c,d	c,d,e	a,b,c
B25-HM+FF	a,b,c,d	d	a,b,c	c,d	a,b,c	c,d,e	a,b
B25-M+HF	a,b,c,d	c	b,c	c,d	b,c	b,c,d,e	b,c,d
B25-M+FF	a,b	c	a,b	c	b,c	a,b,c,d	b,c,d
B50-HM-HF	a,b,c	a,b	a,b	b	a,b	a,b,c,d	a
B50-HM+FF	a	a,b	a,b	a,b	a,b	a,b	a,b,c
B50-M+HF	a,b	a	a,b	a	a,b	a	a,b,c,d
B50-M+FF	a,b,c	b	a	a,b	a	a,b,c	a,b

594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605

Different letters within the same column indicates that the mean significantly differs from each other (one-way ANOVA, $P < 0.001$; Tukey multiple comparison test, $P < 0.05$).

606 **Table 5:**

607 Statistical *P* values for three-way ANOVA comparing differences in yield parameters.

	Grain yield		Grains per ear		Thousand grain weight		Biological yield	
	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2
Biochar	***	***	***	NS	NS	***	*	***
FYM	NS	NS	***	NS	**	NS	**	NS
N fertiliser	***	NS	***	NS	*	NS	***	NS
Biochar*FYM	*	NS	NS	NS	***	NS	NS	NS
Biochar*N fertiliser	**	NS	NS	NS	NS	NS	NS	NS
FYM*N fertiliser	**	NS	NS	NS	NS	NS	NS	NS
Biochar*FYM*N fertiliser	NS	NS	NS	NS	NS	NS	NS	NS

608 Asterisks indicate a significant difference at the **P* < 0.05, ***P* < 0.01 and ****P* < 0.001 level; NS, not-significant.

609

610

611 **Table 6:**

612 Statistical *P* values for three-way ANOVA comparing differences in soil quality parameters.

	Organic C		Mineral N		Phosphorus		Ca/Mg		Potassium		EC		pH	
	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2
Biochar	***	***	*	***	***	***	***	***	***	***	***	***	***	*
FYM	**	***	**	***	***	**	***	NS	***	*	NS	*	***	***
N fertiliser	NS	NS	NS	***	NS	NS	NS	NS	NS	NS	NS	NS	**	NS
Biochar*FYM	**	***	*	*	*	*	*	NS	***	NS	NS	***	***	***
Biochar*N fertiliser	NS	NS	NS	NS	NS	NS	***	NS	***	NS	NS	NS	NS	NS
FYM*N fertiliser	NS	**	NS	**	NS	NS	NS	NS	***	NS	NS	NS	NS	NS
Biochar*FYM*N fertiliser	NS	*	NS	*	NS	NS	***	NS	NS	NS	NS	NS	NS	NS

613 Asterisks indicate a significant difference at the **P* < 0.05, ***P* < 0.01 and ****P* < 0.001 level; NS, not-significant. EC, electrical conductivity.

614

615

616

617

618