

Accepted refereed manuscript of:

Glencross B, Irvin S, Arnold S, Blyth D, Bourne N & Preston NP (2014) Effective use of microbial biomass products to facilitate the complete replacement of fishery resources in diets for the black tiger shrimp, *Penaeus monodon*, *Aquaculture*, 431, pp. 12-19.

DOI: [10.1016/j.aquaculture.2014.02.033](https://doi.org/10.1016/j.aquaculture.2014.02.033)

© 2014, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

1 Effective use of microbial biomass products to facilitate the complete replacement of fishery  
2 resources in diets for the black tiger shrimp, *Penaeus monodon*

3

4

5 Brett Glencross<sup>1,2</sup>, Simon Irvin<sup>1,3</sup>, Stuart Arnold<sup>1,3</sup>, David Blyth<sup>1,3</sup>, Nicholas Bourne<sup>1,2</sup> and Nigel  
6 Preston<sup>1,2</sup>

7

8

9 1. CSIRO Food Futures Flagship, 41 Boggo Rd, Dutton Park, QLD 4102, Australia

10 2. CSIRO Marine and Atmospheric Research, GPO Box 2583, Brisbane, QLD 4001, Australia

11 3. CSIRO Marine and Atmospheric Research, PO Box 2066, Woorim, QLD 4507, Australia

12

13

14 (p) 61-7-3826-7236

15 (f) 61-7-3826-7281

16 (e) Brett.Glencross@csiro.au

17

18 Keywords : prawns, fishmeal, alternative ingredients,

19

20 Submitted to : Aquaculture

21

22 **Abstract**

23

24           A series of experiments were conducted with black tiger shrimp (*Penaeus monodon*) juveniles  
25 to firstly determine the effects of reducing fishmeal inclusion in a diet and then to evaluate the  
26 potential for a microbial bioactive to support complete replacement of both fishmeal and fish oil in  
27 feeds when fed under clear-water and green-water conditions. The isoproteic and isoenergetic  
28 replacement of fishmeal resulted in a consistent decline in growth performance indicating that at  
29 every decrease in fishmeal below an inclusion level of 45% there was a decline in performance. In a  
30 subsequent trial undertaken in a clear-water tank system diets devoid of both fishmeal and fish oil fed  
31 to shrimp were demonstrated to produce poorer performance than a fishmeal and fish oil reference  
32 diet. However the addition of a microbial bioactive to the diet resulted in not only a compensation for  
33 the replacement of these ingredients but additional growth. Replication of the clear-water trial in a  
34 green-water tank system produced similar results, but also showed that the green-water system largely  
35 compensated for the performance lost through replacement of fishmeal and fish oil. However it was  
36 also shown that the use of the microbial bioactive in the diets still resulted in improved growth  
37 performance of shrimp. This study has effectively demonstrated a viable strategy for not only a  
38 complete replacement of all fishery products in shrimp diets, but an improved performance strategy.

39 **1. Introduction**

40 There has been a considerable amount of research in the past few decades to improve the  
41 capacity to utilize alternative raw materials in diets for many aquaculture species, including shrimp  
42 (Gatlin et al., 2007; Glencross et al., 2007). Recent progress in the use of raw materials other than  
43 fishmeal and fish oils in diets for shrimp has resulted in significant advancements in the ability to  
44 replace fishmeal and fish oil with terrestrial grain and animal resources (Davis and Arnold, 2000;  
45 Davis et al 2002; Alvarez et al., 2007; Cruz-Suarez et al., 2001; 2007; Smith et al., 2007). However,  
46 most of these studies still have a certain amount of fishmeal or fish oil in the diet, typically rarely  
47 lower than 10% for fishmeal (Cruz-Suarez et al., 2007; Smith et al., 2007). Replacement of fish oil in  
48 many instances has proven even more difficult with few other lipid resources able to provide the  
49 necessary long-chain polyunsaturated fatty acids or provide the required short-chain polyunsaturates  
50 suitable for trophic upgrading (Deering et al., 1997; Lim et al., 1997; Glencross and Smith 1999).

51 However, studies with *Litopenaeus vannamei* have suggested that, when replaced with co-  
52 extruded poultry and soybean meals, fishmeal inclusion could be decreased to 0%, though shrimp  
53 performance (based on shrimp weight gain) improved with increasing replacement of fishmeal and  
54 this effect was not explained (Samocha et al., 2004). In another study it was shown that canola and  
55 soybean meals could be used effectively with as little as 6% fishmeal (Suarez et al., 2009). Pond  
56 studies by Amaya et al. (2007) indicated that all of the fishmeal in diets for *L. vannamei* could be  
57 replaced when soybean and poultry-byproduct meals were used. Indeed in that study neither growth  
58 nor feed use efficiency was compromised through the reduction in fishmeal content. This appears to  
59 indicate that under pond production systems the endogenous feed sources in the pond can help  
60 ameliorate this loss in performance seen with a reduction in fishmeal (Amaya et al., 2007), which has  
61 not been seen in laboratory tank trials (Suarez et al., 2009). None of these successes in achieving very  
62 low fishmeal inclusion in the diets of more carnivorous shrimp species, like *Penaeus monodon*, have  
63 been reported.

64 The recent invention of a microbial biomass based growth promoter (microbial bioactive)  
65 (Novacq™, CSIRO, Dutton Park, QLD, Australia) has resulted in the ability to stimulate shrimp  
66 growth in excess of 50% above that of a standard reference diet of the same basic nutritional  
67 specifications (Glencross et al., 2013). This product therefore also offers some potential in terms of  
68 being able to off-set poorer performance due to a range of formulation changes, including the  
69 complete replacement of any fishery derived resources in the diet of shrimp. Similar such products  
70 appear to have been reported by Burford et al. (2004) and Kuhn et al. (2008, 2009), who used another  
71 microbial biomass products to achieve significant improvements in performance in *L. vannamei*.

72 In the present study, a series of experiments were undertaken with black tiger shrimp  
73 (*Penaeus monodon*) in an attempt to define critical inclusion limits of fishmeal and fish oil for shrimp  
74 diets. In addition to this replacement of fishmeal and fish oil the use of a microbial bioactive was also  
75 examined for its ability stimulate growth performance and to be able to potentially aid the complete

76 replacement of fishmeal and fish oil. The work was undertaken in both indoor laboratory 'clear-water'  
77 and outdoor, zero-exchange 'green-water' conditions. This study aimed to test the hypotheses' that at  
78 a critical threshold of fishmeal and fish oil inclusion that feed intake and growth would decline, but  
79 that the use of a microbial bioactive would ameliorate those declines.

## 80 **2. Materials and Methods**

### 81 *2.1 Study design*

82 A series of three experiments were undertaken to define; 1) the thresholds to replacement of  
83 fish meal in a clear-water tank experiment, 2) the capacity of a microbial bioactive to support the  
84 complete replacement of fishmeal and fish oil in a clear-water tank experiment, and 3) the capacity of  
85 a microbial bioactive to support the complete replacement of fishmeal and fish oil in a green-water  
86 tank experiment.

87

### 88 *2.2 Diet manufacture*

89 Each diet was based on using a standard reference diet of 42% protein and 7% lipid which  
90 was a mimic of the commercial feeds typically used in the Australian shrimp farming industry and  
91 which acts as our industry equivalent performance benchmark (Glencross et al., 1999a). Variants of  
92 this diet were then made by increasing inclusion of both poultry offal meal and a lupin kernel meal.  
93 Details and composition of all ingredients used in this study are presented in Table 1. Each diet was  
94 prepared by ensuring all ingredients were milled to <750 µm, prior to mixing in an upright planetary  
95 mixer (Hobart, Sydney, NSW, Australia). Water was then added during the mixing to form a dough  
96 which was subsequently screw-pressed (Dolly, La Monferrina, Castell'Alfero, Italy) through a 3mm  
97 die and cut to pellet lengths of about 6mm. The pellets were then steamed at 100°C for 5 minutes  
98 before being oven dried at 60°C for 24h. Diets were kept at -20°C when not being fed.

99

### 100 *2.3 Experiment 1*

101 In experiment 1 a series of diets were formulated to the same specifications as the reference  
102 diet but with the fishmeal progressively diluted out of the formulations from 20% to 0% inclusion at  
103 5% increments. In addition to the series of diets with fishmeal diluted out, two additional diets with  
104 5% and 0% fishmeal were formulated which also included 10% of a microbial bioactive (Novacq™,  
105 CSIRO, Dutton Park, QLD, Australia) as a replacement for the wheat component of the diet. An  
106 additional two diets maintained high levels of fishmeal, but also had 5% and 10% inclusion levels of  
107 the microbial bioactive in replacement of wheat as a reference (Table 2).

108

#### 109 *2.3.1 Shrimp collection and trial management*

110 Several hundred individuals (~8 g) of a wild-type genotype of black tiger shrimp (*Penaeus*  
111 *monodon*) were collected from a grow-out pond at Truloff's Prawn Farm (Woolgoolga, QLD,  
112 Australia) by cast-netting and transferred to a holding tank (10,000 L) where they were held pending  
113 allocation to trial tanks. During the holding period (~7 days) they were fed a standard commercial  
114 grower diet (Prawn Grower, Ridley Aquafeeds, Narangba, QLD, Australia).

115 Six shrimp were then allocated to each of 50 x 100 L tanks in an indoor laboratory system.

116 The mean initial weight across all tanks was  $8.19 \pm 0.72$  g. Tanks of the shrimp were maintained with

117 flow-through seawater at a rate of 500 mL/min. Temperatures of each tank were maintained at  $29.2 \pm$   
118  $0.28^\circ\text{C}$  and dissolved oxygen maintained at  $6.4 \pm 0.14$  mg/L. Light was maintained on a 12:12  
119 light:dark cycle. Shrimp were individually weighed at day 0, 14, day 28 and again at day 42. The  
120 mean weight of each tank was determined at each assessment point to calculate the mean weight for  
121 each treatment, with tanks used as the replicate ( $n = 5$  per treatment). During this period the shrimp  
122 were manually fed the diets twice daily to marginal excess and the amount of feed remaining the  
123 following day scored and used to adjust the next day's ration (increase or decrease) according to a  
124 feed intake score. Uneaten feed was siphoned from each tank daily after scoring. The assessment was  
125 also used to provide a quantitative measurement of uneaten feed in each tank. This method was also  
126 used to estimate as accurately as possible feed intake within each tank on each day (Smith et al.,  
127 2007b).

128

## 129 2.4 *Experiment 2*

130 In Experiment 2, a series of diets were formulated to the same specifications as the reference  
131 diet but with the fishmeal reduced in the formulations to either 10% or 0% inclusion. In addition to  
132 the series of diets with the reduced fishmeal, a corresponding series of diets (with 10% and 0%  
133 fishmeal) were formulated with linseed oil replacing all fish oil. A further additional corresponding  
134 series of diets (with 10% and 0% fishmeal) were formulated with the microbial bioactive in  
135 replacement of the wheat component of the diet. A final additional corresponding series of diets was  
136 formulated with linseed oil replacing all fish oil and also including the microbial bioactive (Table 3).

137

### 138 2.4.1 *Shrimp collection and trial management*

139 Several hundred individuals ( $\sim 4$  g) of a wild-type genotype of shrimp were collected from a  
140 grow-out pond at the Bribie Island Research Centre (DEEDI, Woorim, QLD, Australia) by cast-  
141 netting and transferred to a holding tank (10,000 L) where they were held pending allocation to trial  
142 tanks. During the holding period ( $\sim 7$  days) they were fed a standard commercial grower diet (Prawn  
143 Grower, Ridley Aquafeeds, Narangba, QLD, Australia).

144 Six shrimp were then allocated to each of  $40 \times 100$  L tanks in an indoor laboratory system.  
145 The mean initial weight across all tanks was  $4.35 \pm 0.04$  g. Tanks of the shrimp were maintained with  
146 flow-through water at a flow rate of 500 mL/min. Temperatures of each tank were maintained at  $28.9$   
147  $\pm 0.15^\circ\text{C}$  and dissolved oxygen maintained at  $5.3 \pm 0.11$  mg/L. Light was maintained on a 12:12  
148 light:dark cycle. Shrimp were individually weighed at day 0, 14, day 28 and again at day 42. The  
149 mean weight of each tank was determined at each assessment point to calculate the mean weight for  
150 each treatment, with tanks used as the replicate ( $n = 4$  per treatment). During this experiment feeding  
151 of the shrimp was managed the same as detailed for Experiment 1.

152

## 153 2.5 *Experiment 3*

154 In Experiment 3 the same series of diets as used in Experiment 2 were fed to shrimp kept in  
155 green-water conditions. Green-water conditions were established in each of 30 x 2400 L tanks about 7  
156 days before being stocked with shrimp and maintained as a zero-water-exchange system for the  
157 duration of the experiment. Freshwater was added to tanks as required to maintain water level and  
158 salinity change from evaporation. To generate a green-water culture each tank was filled with  
159 seawater filtered through a 100µm screen and then fertilised with 70g of Aquasol™ (Yates, Padstow,  
160 NSW, Australia). Each of the 30 tanks was then recirculated through a common sump to ensure  
161 homogeneity in the tank conditions. After seven days each tank had established a consistent  
162 microalgal bloom within and this was consistent across all 30 tanks.

163

### 164 2.5.1 *Shrimp collection and trial management*

165 Several hundred individuals (~3 g) of a wild-type genotype of shrimp were collected from a  
166 grow-out pond at the Pacific Reef Farms Pty Ltd (Ayr, QLD, Australia) by cast-netting and  
167 transferred by air-freight to Brisbane where they were placed in several holding tanks (10,000 L) and  
168 held pending allocation to trial tanks. During the holding period (~7 days) they were fed a standard  
169 commercial grower diet (Prawn Grower, Ridley Aquafeeds, Narangba, QLD, Australia).

170 Thirty shrimp were then allocated to each of 30 x 2400 L tanks in an outdoor tank system  
171 kept in a glasshouse. The mean initial weight across all tanks was  $3.89 \pm 0.07$  g (mean  $\pm$  S.D.). Tanks  
172 were maintained on a static water basis with regular aeration and a top up with fresh seawater to  
173 maintain volumes after evaporation. Temperatures of each tank were maintained at  $28.1 \pm 1.8^\circ\text{C}$  and  
174 dissolved oxygen averaged  $7.1 \pm 0.25$  mg/L. Light was ambient over the period of the study (8<sup>th</sup>  
175 November 2012 to 10<sup>th</sup> January 2013). Water salinity  $38.1 \pm 1.01$  ppt and pH  $8.1 \pm 0.11^\circ\text{C}$ . Secchi  
176 depth was also measured weekly and over the 63 days averaged  $75 \pm 10.6$  cm. Shrimp were  
177 individually weighed at day 21, day 42 and again at day 63. The mean weight of each tank was  
178 determined at each assessment point to calculate the mean weight for each treatment, with tanks used  
179 as the replicate (n = 3 per treatment). During this period the shrimp were manually fed the diets twice  
180 daily on either of two feed trays within each tank to marginal excess and the amount of feed  
181 remaining the following day scored and used to adjust the next day's ration (increase or decrease)  
182 according to feed intake score. This method was also used to estimate as accurately as possible feed  
183 intake within each tank on each day. Uneaten feed was emptied from the feed tray after scoring each  
184 day. This feeding method was adapted from Smith et al (2002).

185

### 186 2.6 *Chemical analysis*

187 Ingredient samples were analysed for dry matter, ash, nitrogen, amino acids, total lipid,  
188 carbohydrate and gross energy content. Diets and whole shrimp samples were also analysed for fatty  
189 acid composition. Dry matter of the samples was calculated by gravimetric analysis of a milled  
190 sample following oven drying at  $105^\circ\text{C}$  for 24 h. Protein levels were calculated from the



191 determination of total nitrogen by CHNOS auto-analyser, based on N x 6.25. Amino acid analysis  
192 involved the samples being hydrolyzed at 110°C for 24 h in 6M HCl with 0.05% Phenol. Cystine was  
193 derivatized during hydrolysis by the addition of 0.05% 3-3-dithiodipropionic acid. The acid  
194 hydrolysis destroyed tryptophan making it unable to be determined. Separation was by HPLC on a  
195 Hypersil AA-ODS 5µm column using an 1100 series Hewlett Packard HPLC system. Gross ash  
196 content was determined gravimetrically following loss of mass after combustion of a sample in a  
197 muffle furnace at 550°C for 12 h. Total lipid content of the diets was determined gravimetrically  
198 following extraction of the lipids using chloroform:methanol (2:1). Fatty acids were analysed as  
199 methyl ester derivatives. Lipids were esterified by the method of O'Fallon et al. (2007) and analysed  
200 by gas chromatography (GC) using flame ionization detection. Peaks were identified by comparing  
201 relative retention times to standards. Carbohydrates were estimated based on dry matter content of the  
202 feed minus the lipid, ash and protein contents. Gross energy was determined by adiabatic bomb  
203 calorimetry. All methods were consistent with those recommended by AOAC (2005).

204

## 205 2.7 *Statistical analysis*

206 All values are means ± standard deviations unless otherwise specified. Significant differences  
207 were determined using an ANOVA followed by a Tukey's HSD test with critical ranges were set at P  
208 < 0.05. These tests were undertaken using the Statistica™ v6.0 software (Statsoft®, Tulsa, OA, USA).  
209 Linear regression analysis and line fitting of those relationships was undertaken using the data  
210 analysis tools and graphics elements of Microsoft Excel.

211 **3. Results**

212 *3.1 Experiment 1*

213 There were significant effects on shrimp growth due to the reduction in fishmeal inclusion  
214 (Table 4). Growth (as defined by final weight, weight gain, gain rate) of those shrimp fed the  
215 reference diet was typical for that of this species in a clear-water tank system over a six week period  
216 ( $0.91 \pm 0.11$  g/wk/shrimp) (Smith et al., 2007b). The progressive replacement of fishmeal with a  
217 balance of poultry offal meal and lupin kernel meal resulted in a significant decline in growth  
218 ( $P=0.026$  as regression; Fig. 1) to a lowest performance at 10% inclusion of fishmeal, though this was  
219 not significantly different from the growth of shrimp fed diets with either 5% or 0% fishmeal. Based  
220 on an ANOVA analysis the critical inclusion level where growth was poorer than the reference diet  
221 (F45:M0) was the diet with 10% fishmeal (F10:M0). The addition of the microbial bioactive resulted  
222 in a numerical, but not a significant ( $P=0.063$ ) improvement in growth. The addition of the microbial  
223 bioactive to diets with either low or no fishmeal resulted in those shrimp subsequently fed those diets  
224 performing equal to the reference diet (F45:M0). Feed conversion ranged from 3.92 to 6.42 and but  
225 was not significantly affected ( $P=0.116$ ) by fishmeal inclusion level.

226

227 *3.2 Experiment 2*

228 In clear-water indoor laboratory conditions the complete replacement of both fish meal and  
229 fish oil resulted in a progressive decline in growth (as defined by final weight, weight gain and gain  
230 rate) of those shrimp in experiment 2 (Table 5). Growth of shrimp fed the diet completely free of fish  
231 meal and fish oil (F0:O0:M0) was significantly ( $P<0.001$ ) poorer than that of those shrimp fed the  
232 reference diet (F50:O2:M0). This effect was despite that the reduction in fish meal and fish oil  
233 resulted in increased feed intake by the shrimp (Table 5). The addition of the microbial bioactive  
234 (F50:O2:M10) resulted in a significant increase in growth ( $P<0.001$ ) and feed intake ( $P=0.034$ ) by the  
235 shrimp. When added to diets with low or no fish meal and/or fish oil (diets F10:O0:M10, F0:O0:M10,  
236 F10:O2:M10 and F0:O2:M10) this increase in growth induced by the microbial bioactive exceeded  
237 the losses in growth observed with the complete replacement of fish meal and fish oil (Table 5). Feed  
238 conversion was not significantly affected by any treatment, largely due to the inherent variability  
239 associated with this parameter.

240 There were a range of significant effects of the dietary treatments on the composition of the  
241 shrimp (Table 7). Notable changes included a decrease in the LC-PUFA with a decrease of either (or  
242 both) fishmeal and fish oil in the diet. This was caused principally by a decline in the percentage of  
243 C20:5n-3, C22:5n-3 and C22:6n-3. This decline in the LC-PUFA and notably the Total n-3 also  
244 resulted in a significant change to the n-3 : n-6 ratio in the fatty acid composition of the shrimp, which  
245 was further exacerbated by an increase in the Total n-6 fatty acids.

246

247 *3.3 Experiment 3*

248           In green-water zero-exchange outdoor tank conditions the complete replacement of both fish  
249 meal and fish oil had no significant ( $P=0.994$ ) effect on growth (as defined by final weight, weight  
250 gain and gain rate) of those shrimp in experiment 3 (Table 6). However, the addition of the microbial  
251 bioactive to the diet resulted in a significant ( $P=0.005$ ) increase in growth and feed intake by the  
252 shrimp. When added to diets with low or no fish meal and/or fish oil this increase in growth induced  
253 by the microbial bioactive was greater than that observed when the diets had high levels of fish meal  
254 and fish oil (F50:O2:M10) (Table 6).

255           As with experiment 2 there were also a range of significant effects on the composition of the  
256 shrimp due to the dietary treatments in experiment 3 (Table 8). In contrast to experiment 2, there were  
257 no significant differences among treatments in the percentages of Total LC-PUFA in experiment 3.  
258 Although there was some variability in the percentages of C20:5n-3 and C22:6n-3 with changes in  
259 dietary treatments in this experiment, it was the substantial increase in the levels of C20:4n-6 that  
260 caused the absence of a significant difference in the Total LC-PUFA.

261

## 4. Discussion

The use of fishmeal in shrimp diets has long been identified as one of the critical issues affecting sustainability of feed use in this aquaculture sector (Tacon and Metian, 2008). However, despite this apparent reliance on the use of fishmeal there was little data in the literature on the consequences of diluting the use of fishmeal through the inclusion of other protein sources. In addition to attempts to define critical inclusion limits of fishmeal for shrimp diets, the development of a microbial bioactive (Novacq™; CSIRO, Brisbane, QLD, Australia) also holds promise for being able to potentially aid the complete replacement of both fishmeal and fish oil (Ju et al., 2008; Kuhn et al., 2009). This study therefore aimed to test a series of hypotheses. Firstly, that at a critical threshold of fishmeal inclusion feed intake and growth would decline. Secondly, that the use of a microbial bioactive would ameliorate those declines and those also attributable to complete replacement of fish oil as well. It was also hypothesised that these effects would be seen in both clear-water and green-water conditions.

### 4.1 Thresholds to fishmeal inclusion

From experiment 1 it was clear that reducing the fishmeal inclusion in diets for *P. monodon* had a negative effect on performance. Based on the regression analysis a decline in growth was observed with every level of reduction in fishmeal. Though, based on an ANOVA analysis though this difference was only significant at the 10% inclusion level of fishmeal and below. This observation supports the fact that although regression analysis presents a case for decline in performance with any replacement of fishmeal, it tangibly only becomes an issue once fishmeal is as low as 10%. However, the results from experiment 1 provide clear support to the claims of importance of fishmeal use in diets for shrimp (Tacon and Metian, 2008).

Other researchers who have examined the replacement of fishmeal with other alternative protein sources have also noted a similar response. Lim and Dominy (1990) replaced fishmeal (mixed with squid and shrimp meal) in series of diets for *L. vannamei* using soybean meal and achieve diets with 0 g/kg fishmeal. However, these authors observed that below 213 g/kg inclusion of the fishmeal mix that growth began to significantly deteriorate. Other studies using co-extruded poultry and soybean meals as replacements found that they could achieve equal performance with 0g/kg fishmeal inclusion, and interestingly shrimp performance appeared to improve with increasing replacement of fishmeal, though this observation wasn't rationalised. Notably, though the addition of 10 g/kg of krill meal made a notable improvement to a diet with 0 g/kg of fishmeal (Samocho et al., 2004). In other studies it has been demonstrated that use of both rapeseed and soybean meals can be effective with as little as 6% fishmeal inclusion in the diet (Suarez et al., 2009). Though notably, most of these studies have relied on the addition of a krill or squid meal supplement to achieve equal or slightly poorer performance of shrimp fed fishmeal free diets.

299 4.2 *Growth stimulation by microbial bioactive inclusion*

300 In experiment 1 the addition of the microbial bioactive to the diet was also observed to  
301 improve the growth of the shrimp (both in the high and low fishmeal based diets; diets F45:M5,  
302 F45:M10, F5:M10 and F0:M10). Indeed if the performance of the shrimp fed diets F5:M10 and  
303 F0:M10 which had 5% and 0% fishmeal respectively is compared against their analogous diets  
304 (F5:M0 and F0:M0) without the microbial bioactive it can be noted that their performance is 40% to  
305 46% better. This contrasts the diet fortified with microbial bioactive that still had high fishmeal (diets  
306 F45:M5 and F45:M10), but the growth improvement was only 11% to 18% better than that of the  
307 reference (F45:M0). However, in experiment 2 the inclusion of 10% of the microbial bioactive in to a  
308 similar diet with high levels of fishmeal increased growth rates of shrimp in indoor lab conditions  
309 from 0.87 g/wk to 1.30 g/wk, an increase of 50%. We postulate that this variability in response may  
310 be an effect of shrimp stock used, with different stocks used for each of the different experiments.

311 Other researchers have also recently reported the use of microbial biomass products in shrimp  
312 feeds with varying degrees of success (De Schryver et al., 2008; Ju et al., 2008; 2009; Kuhn et al.,  
313 2008; 2009). In the work of Ju et al (2008), the authors included a microbial biomass in at 20%  
314 inclusion level and observed an increase in growth rate of *L. vannamei* from 0.85 g/wk to 1.03 g/wk in  
315 indoor lab conditions, a 21% increase in growth rate. These authors also went on to fractionate the  
316 microbial biomass into various chemical fractions and identified that the prospective bioactive  
317 component was in the acetone soluble fraction, possibly a carotenoid of some sort, though none of the  
318 fractions produced in their study were significantly different from each other in terms of growth rates  
319 achieved.

320

321 4.3 *Complete replacement of fishmeal and fish oil in clear-water by using microbial bioactive*

322 The use of clear-water experimental systems provides the most stringent manner in which to  
323 assess the impact of a feed on shrimp performance because it isolates other exogenous factors other  
324 than the feed provided (Cuzon et al., 2004). In the second experiment, consistent with the  
325 observations of the first, there was a consistent decline in growth of the shrimp with reduction in  
326 fishmeal content of the diet. However, significant improvements were seen to both the 10% fishmeal  
327 (diet F10:O2:M0) and 0% fishmeal (diet F0:O2:M0) treatments with the addition of the microbial  
328 bioactive. In each case the relative biomass gains for each diet with its “paired” microbial bioactive  
329 fortified analogue (diet F10:O2:M0 cf. F10:O2:M10 and diet F0:O2:M0 cf. F0:O2:M10) were close to  
330 or over 100% better.

331 The replacement of fish oil with linseed oil was arguably more successful and is consistent  
332 with earlier observations and established knowledge on the management of essential fatty acid  
333 demands in this species (Deering et al., 1997; Glencross and Smith, 1999; Glencross et al., 2002b). In  
334 those studies it was shown that provided the n-3 and n-6 fatty acids were kept in ‘balance’ then the  
335 animal had sufficient capacity to synthesise their own LC-PUFA and in fact that if this optimal

336 balance was maintained and nominal amounts of LC-PUFA supplied then this actually promoted  
337 superior growth. These earlier observations are again confirmed where it can be seen that replacement  
338 of the fish oil with linseed oil (diet F10:O0:M0) and addition to the diet with 10% fishmeal (which  
339 introduces nominal amounts of LC-PUFA) actually resulted in numerically better growth than that  
340 achieved with the comparative diet with fish oil (diet F10:O2:M0).

341

#### 342 4.4 Complete replacement of fishmeal and fish oil in green-water by using microbial bioactive

343 It has been suggested that the culture system used in experimentation with shrimp can have a  
344 significant impact on the results achieved (Tacon et al., 2002; Cuzon et al., 2004). Cuzon et al. (2004)  
345 proposed that it was more appropriate to conduct shrimp trials in outdoor green-water tanks that better  
346 mimicked pond conditions. In the present study there is some indication to support this, in that the  
347 observation of reduced growth with shrimp fed no fishery products in their diet from the clear-water  
348 trial was not apparent in the green-water trial, suggesting that the shrimp were obtaining some level of  
349 nutrition from the green-water system that helped offset those growth losses observed in the clear-  
350 water trial. However, the growth stimulatory effect of the microbial bioactive was apparent in both  
351 experimental systems. Additionally, the use of this microbial bioactive and the complete replacement  
352 of fishery resources had no significant impact on the total LC-PUFA content of the shrimp produced  
353 using the green-water system.

354 There are few studies where there has been complete replacement of fish meal and/or fish oil,  
355 in green-water or pond conditions. One such study that did attempt complete replacement of fish  
356 meals was that by Amaya et al. (2007), who fed diets based on poultry by-product and soybean meals  
357 to *L. vannamei* in pond trials with fishmeal levels ranging from 90 g/kg to 0 g/kg. These authors  
358 observed after 81 days that there was no significant difference ( $P=0.072$ ) in growth between those  
359 shrimp fed the different treatments, although the one with 0 g/kg fishmeal was the poorest of all  
360 treatments. In this same study Amaya et al. (2007) also examined a diet made solely with plant protein  
361 sources, but also supplemented with squid meal (10 g/kg). All experimental diets also performed  
362 numerically poorer than the commercial reference used. In the green-water trial in the present study,  
363 equal or superior growth was achieved with no aquatic animal meals or oils included in the diet. In  
364 addition, the inclusion of a microbial bioactive resulted in superior performance of those same  
365 fishmeal and fish oil free diets into which it was included. Therefore we contest that the present study  
366 is the first reported that has achieved complete replacement of all fishery derived products and  
367 demonstrated equal or superior performance and therefore represents a major step forward in terms of  
368 complete independence from reliance of fishery products to underpin shrimp production.

369

#### 370 4.5 Conclusions

371 The findings of this study demonstrate that through the use of a microbial bioactive product  
372 containing growth promoting properties that it is possible to completely off-set the need for both fish

373 meal and fish oil and produce diets for shrimp that have no reliance on any fishery products. While  
374 other studies have shown the ability replace fish meal or fish oil, in most cases noted this has only  
375 been achieved through the use of other fishery resources like squid or crustacean meals (Amaya et al.,  
376 2007). The findings from this study are a major progression in the sustainability of the shrimp farming  
377 industry in that it demonstrates clear potential for complete independence from fishery resources. This  
378 outcome is yet to be achieved with any other marine species.  
379

380 **Acknowledgements**

381 Some of the chemical analyses work undertaken by Malcolm McGrath of the Department of  
382 Food and Agriculture Western Australia is gratefully acknowledged. Special thanks to Mat Cook for  
383 constructive comment and editorial on an early draft.

384

385

386 **References**

387 AOAC (Association of Official Analytical Chemists). 2005. Official Methods of Analysis of the  
388 Association of Official Analytical Chemists. 15th edition. Association of Official Analytical  
389 Chemists. Washington, DC, USA.

390

391 Alvarez, J. S., Hernández-Llamas, A., Galindo, J., Fraga, I., García, T., Villareal, H., 2007.  
392 Substitution of fishmeal with soybean meal in practical diets for juvenile white shrimp *Litopenaeus*  
393 *schmitti*. Aquaculture Research 38, 689–695.

394

395 Amaya, E.A., Davis, D.A., Rouse, D.B., 2007. Replacement of fish meal in practical diets for the  
396 Pacific white shrimp (*Litopenaeus vannamei*) reared under pond conditions. Aquaculture 262, 393-  
397 401.

398

399 Burford, M.A., Thompson, P.J., McIntosh, R.P., Bauman, R.H., Pearson, D.C., 2004. The contribution  
400 of flocculated material to shrimp (*Litopenaeus vannamei*) nutrition in a high-intensity zero-exchange  
401 system. Aquaculture 232, 525-537.

402

403 Cruz-Suarez, L.E., Guillaume, J., Van Wormhoudt, A., 1987. Effect of various levels of squid protein  
404 on growth and some biochemical parameters of *Penaeus japonicus* juveniles. Bull. Japan. Soc. Sci.  
405 Fish, 53 (11), 2083-2088.

406

407 Cruz-Suarez, L.E., Ricque-Marie, D., Tapia-Salazar, M., McCallum, I.M., Hickling, D., 2001.  
408 Assessment of differently processed feed pea (*Pisum sativum*) meals and canola meal (*Brassica* sp.) in  
409 diets for blue shrimp (*Litopenaeus stylirostris*). Aquaculture 196, 87–101.

410

411 Cruz-Suárez, L.E., Nieto-López, M., Guajardo-Barbosa, C., Tapia-Salazar, M., Scholz, U., Ricque-  
412 Marie, D., 2007. Replacement of fish meal with poultry by-product meal in practical diets for  
413 *Litopenaeus vannamei*, and digestibility of the tested ingredients and diets. Aquaculture 272, 466-476.

414

415 Cuzon, G., Lawrence, G., Gaxiola, C., Rosas, C. and Guillaume, J., 2004. Nutrition of *Litopenaeus*  
416 *vannamei* reared in tanks or ponds. Aquaculture, 235, 513-551.



417  
418 Davis, D.A., Arnold, C.R., 2000. Replacement of fish meal in practical diets for the Pacific white  
419 shrimp, *Litopenaeus vannamei*. Aquaculture 185, 291–298.  
420  
421 Davis, D.A., Arnold, C.R., McCallum, I., 2002. Nutritional value of feed peas (*Pisum sativum*) in  
422 practical diet formulations for *Litopenaeus vannamei*. Aquac. Nutr. 8, 87–94.  
423  
424 Deering, M.J., Fielder D.R., Hewitt, D.R., 1997. Growth and fatty acid composition of juvenile leader  
425 prawns, *Penaeus monodon*, fed different lipids. Aquaculture 151, 131-141.  
426  
427 De Schryver, P., Crab, R., Defoirdt, T., Boon, N., Verstraete, W., 2008. The basics of bio-flocs  
428 technology: the added value for aquaculture. Aquaculture 277, 125-137.  
429  
430 Gatlin, D.M., Barrows, F.T., Brown, P., Dabrowski, K., Gaylord, T.G., Hardy, R.W., Herman, E., Hu,  
431 G., Krogdahl, A., Nelson, R., Overturf, K., Rust, M., Sealy, W., Skonberg, D., Souza, E.J., Stone, D.,  
432 Wilson, R., Wurtele, E., 2007. Expanding the utilisation of sustainable plant products in aquafeeds: a  
433 review. Aquaculture Research 38, 551-579.  
434  
435 Glencross, B.D., Smith, D.M., Tonks, M.L., Tabrett, S.M., Williams, K.C., 1999. A reference diet for  
436 nutritional studies of the prawn, *Penaeus monodon*. Aquaculture Nutrition 5, 33-40.  
437  
438 Glencross, B.D., Smith, D.M., 1999. The linoleic and linolenic acids requirements of the prawn,  
439 *Penaeus monodon*. Aquaculture Nutrition 5, 53-64  
440  
441 Glencross, B.D., Smith, D.M., Thomas, M.R., Williams, K.C. 2002b. Optimising the essential fatty  
442 acid and total neutral lipid requirements for weight gain of the prawn, *Penaeus monodon*. Aquaculture  
443 204, 85-99.  
444  
445 Glencross, B.D., Booth, M., Allan, G.L. 2007. A feed is only as good as its ingredients – A review of  
446 ingredient evaluation for aquaculture feeds. Aquaculture Nutrition 13, 17 – 34.  
447  
448 Glencross, B.D., Tabrett, S.J., Irvin, S., Wade, N., Smith, D.M., Coman, G.E., Preston, N.P., 2013. An  
449 examination of the genotype by diet interaction effects in the Black Tiger shrimp, *Penaeus monodon*.  
450 – why do selected shrimp grow faster ? Aquaculture Nutrition doi: 10.1111/j.1365-2095.2012.00941.x  
451

452 Ju, Z.Y., Forster, I., Conquest, L., Dominy, W., 2008. Enhanced growth effects on shrimp  
453 (*Litopenaeus vannamei*) from inclusion of whole shrimp floc or floc fractions to a formulated diet.  
454 Aquaculture Nutrition, 14, 533-543.  
455

456 Ju, Z. Y., Forster, I.P., Dominy, W.G., 2009. Effects of supplementing two species of marine algae or  
457 their fraction to a formulated diet on growth, survival and composition of shrimp (*Litopenaeus*  
458 *vannamei*). Aquaculture, 292, 237-243.  
459

460 Kuhn, D.D., Boardman, G.D., Lawrence, A.L., Marsh, L., Flick Jr. G.L., 2008. Use of microbial flocs  
461 generated from Tilapia effluent as a nutritional supplement for shrimp *Litopenaeus vannamei*, in  
462 recirculating aquaculture systems. Journal of the World Aquaculture Society, 39, 72-82.  
463

464 Kuhn, D.D., Boardman, G.D., Lawrence, A.L., Marsh, L., Flick Jr. G.L., 2009. Microbial floc meal as  
465 a replacement ingredient for fish meal and soybean protein in shrimp feed. Aquaculture, 296, 51-57.  
466

467 Lim, C., Dominy, W., 1990. Evaluation of soybean meal as a replacement for marine animal protein  
468 in diets for shrimp *Penaeus vannamei*. Aquaculture 87, 53–64.  
469

470 Lim, C., Ako, H., Brown, C.L. Hahn, K.O., 1997. Growth response and fatty acid composition of  
471 juvenile *Penaeus vannamei* fed different sources of dietary lipid. Aquaculture 151, 143-153.  
472

473 O'Fallon, J.V., Busboom, J.R., Nelson, M.L. & Gaskins, C.T., 2007. A direct method for fatty acid  
474 methyl ester (FAME) synthesis: Application to wet meat, tissues, oils & feedstuffs. Journal of Animal  
475 Science 85, 1511-1522  
476

477 Samocha, T.M., Davis, A.D., Soud, P.I., de Bault, K., 2004. Substitution of fish meal by coextruded  
478 soybean poultry by-product meal in practical diets for the Pacific white shrimp, *Litopenaeus*  
479 *vannamei*. Aquaculture 231, 197–203.  
480

481 Smith, D.M., Burford, M.A., Tabrett, S.J., Irvin, S.J., Ward, L. 2002. The effect of feeding frequency  
482 on water quality and growth of the black tiger shrimp (*Penaeus monodon*). Aquaculture 207, 125–  
483 136.  
484

485 Smith, D.M., Tabrett, S.J., Glencross, B.D., Irvin, S.J., Barclay, MC., 2007a. Digestibility of lupin  
486 kernel meals in feeds for the black tiger shrimp, *Penaeus monodon*. Aquaculture 264, 353-362.  
487

488 Smith, D.M., Tabrett, S.J., Glencross, B.D. 2007b. Growth response of the black tiger shrimp,  
489 *Penaeus monodon* fed diets containing different lupin cultivars. Aquaculture 269, 236-246.  
490

491 Suarez, J.A., Gaxiola, G., Mendoza, R., Cadavid, S., Garcia, D., Alanis, G., Suarez, A., Faillace, J.,  
492 Cuzon, G., 2009. Substitution of fish meal with plant protein sources and energy budget for white  
493 shrimp *Litopenaeus vannamei* (Boone 1931). Aquaculture 289, 118-123.  
494

495 Tacon, A.G.J., Cody, J.J., Conquest, L.D., Divakaran, S., Forster, I.P., Decamp, O.E., 2002. Effect of  
496 culture system on the nutrition and growth performance of Pacific white shrimp *Litopenaeus*  
497 *vannamei* (Boone) fed different diets. Aquaculture Nutrition, 8, 121-137.

## Tables and Figures

Table 1. Composition of the key experimental ingredients (all values are g/kg dry basis - unless otherwise specified).

	Fishmeal	Gluten	Wheat	Lupin	POM	MB
Dry matter (g/kg)	912	904	900	921	906	917
Protein	753	807	129	418	680	42
Lipid	102	22	22	55	182	6
Ash	159	8	839	30	151	269
Carbohydrates	0	163	10	497	69	683
Energy (kJ/g DM)	21.5	22.1	18.4	20.0	21.3	13.0
Alanine	45	19	4	16	48	2
Arginine	40	26	6	55	45	1
Aspartic acid	66	25	7	46	54	4
Cystine	9	20	1	7	8	0
Glutamate	92	299	40	82	87	3
Glycine	42	25	5	18	60	2
Histidine	23	13	1	11	12	0
Isoleucine	32	28	4	18	25	2
Leucine	55	53	9	31	46	2
Lysine	55	11	5	18	37	1
Methionine	23	15	2	4	18	1
Phenylalanine	29	43	6	18	26	1
Proline	30	115	25	21	46	6
Serine	30	40	6	23	28	2
Taurine	7	0	0	0	4	0
Threonine	32	21	5	17	28	3
Tyrosine	24	27	4	18	20	1
Valine	37	28	5	16	27	2

POM: Poultry offal meal. MB : Microbial bioactive

Table 2. Formulations and composition of diets from Experiment 1. Data are percent values.

	F45:M0	F45:M5	F45:M10	F20:M0	F15:M0	F10:M0	F5:M0	F0:M0	F5:M10	F0:M10
Fish meal (anchovetta) <sup>a</sup>	45.00	45.00	45.00	20.00	15.00	10.00	5.00	0.00	5.00	0.00
Gluten (wheat) <sup>b</sup>	5.00	5.00	5.00	3.00	4.75	6.50	8.25	10.00	8.25	10.00
Wheat flour <sup>b</sup>	47.03	42.03	37.03	35.03	34.53	34.03	33.53	33.03	23.53	23.03
Lecithin <sup>a</sup>	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Fish Oil <sup>a</sup>	1.50	1.50	1.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Microbial bioactive <sup>c</sup>	0.00	5.00	10.00	0.00	0.00	0.00	0.00	0.00	10.00	10.00
Lupin kernel meal <sup>d</sup>	0.00	0.00	0.00	20.00	22.50	25.00	27.50	30.00	27.50	30.00
Poultry offal meal <sup>a</sup>	0.00	0.00	0.00	20.00	21.25	22.50	23.75	25.00	23.75	25.00
Astaxanthin (10%) <sup>g</sup>	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Cholesterol <sup>e</sup>	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Banox E <sup>f</sup>	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Vitamin C (Stay C) <sup>g</sup>	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Vitamin premix <sup>h</sup>	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
<i>Composition (% DM unless otherwise stated)</i>										
Dry matter (% as is)	92.6	93.0	92.0	92.9	93.3	93.5	92.7	92.1	92.2	93.0
Protein	43.8	43.8	44.5	45.0	44.5	44.0	44.0	43.9	44.0	44.1
Lipid	7.0	7.0	7.1	7.9	7.7	7.6	7.6	7.5	7.6	7.5
Ash	7.2	8.4	9.5	6.7	6.1	5.6	5.1	4.6	7.4	6.9
Carbohydrate	41.9	40.9	38.9	40.5	41.7	42.8	43.3	44.0	40.9	41.5
Gross Energy (MJ/kg DM)	20.3	20.1	19.9	20.6	20.7	20.7	20.8	20.8	20.4	20.5

<sup>a</sup> Fish (Peruvian anchovetta) meal, Poultry offal meal, Lecithin and Fish (Peruvian anchovetta) oil : Ridley Aquafeeds, Narangba, QLD, Australia. <sup>b</sup> Wheat gluten and flour : Manildra, Auburn, NSW, Australia. <sup>c</sup> Microbial bioactive: Novacq™ : CSIRO, Cleveland, QLD, Australia, PCT Patent AU 2008201886. <sup>d</sup>Lupin (*L. angustifolius*) kernel meal: Coorow Seed Cleaners, Coorow, WA, Australia. <sup>e</sup> Cholesterol : MP Bio, Aurora, OH, USA. <sup>f</sup> Banox-E™ : BEC Feed Solutions, Carole Park, QLD, Australia. <sup>g</sup> Astaxanthin (10%) as Carophyll Pink™ and Stay C™: DSM, Wagga Wagga, NSW, Australia. <sup>h</sup> Vitamin and mineral premix : Rabar, Beadesert, QLD, Australia; includes (IU/kg or g/kg of premix): Vitamin A, 2.5MIU; Vitamin D3, 1.25 MIU; Vitamin E, 100 g; Vitamin K3, 10 g; Vitamin B1, 25 g; Vitamin B2, 20 g; Vitamin B3, 100 g; Vitamin B5, 100; Vitamin B6, 30 g; Vitamin B9, 5; Vitamin B12, 0.05 g; Biotin, 1 g; Vitamin C, 250 g; Banox-E, 13 g; <sup>h</sup>Yttrium oxide: Stanford Materials, Aliso Viejo, CA, USA.

Table 3 Formulations and composition of the diets from Experiments 2 and 3. Data are percent values.

	F50:O2:M0	F50:O2:M10	F10:O2:M0	F0:O2:M0	F10:O0:M0	F0:O0:M0	F10:O0:M10	F0:O0:M10	F10:O2:M10	F0:O2:M10
Fish Meal (anchovetta)	50.00	50.00	10.00	0.00	10.00	0.00	10.00	0.00	10.00	0.00
Gluten	5.00	5.00	10.00	13.00	10.00	13.00	10.00	13.00	10.00	13.00
Wheat Flour	41.23	31.23	34.33	31.53	34.33	31.53	24.33	21.53	24.33	21.53
Lecithin	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Fish Oil	2.30	2.30	2.50	2.50	-	-	-	-	2.50	2.50
Microbial bioactive	-	10.00	-	-	-	-	10.00	10.00	10.00	10.00
Linseed Oil	-	-	-	-	2.50	2.50	2.50	2.50	-	-
Lupin kernel meal	-	-	15.00	20.00	15.00	20.00	15.00	20.00	15.00	20.00
Poultry Meal	-	-	26.00	30.00	26.00	30.00	26.00	30.00	26.00	30.00
DL-Methionine	-	-	0.50	1.00	0.50	1.00	0.50	1.00	0.50	1.00
L-Lysine	-	-	0.20	0.50	0.20	0.50	0.20	0.50	0.20	0.50
Carophyll Pink	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Cholesterol	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Banox E	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Vitamin C	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Vitamin Premix	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
<i>Composition (% DM unless otherwise stated)</i>										
Dry matter (% as is)	95.21	94.75	94.79	94.84	93.69	95.37	95.8	95.06	96.1	95.98
Protein	47.26	45.05	47.61	47.45	45.76	48.17	44.62	46.62	43.70	44.54
Lipid	9.87	8.42	10.04	8.58	8.36	10.56	9.52	10.16	10.19	10.38
Ash	10.18	17.47	6.88	5.88	6.92	5.77	13.82	12.60	13.78	12.93
Carbohydrate	32.69	29.06	35.47	38.09	38.96	35.51	32.03	30.63	32.33	32.15
Gross Energy (MJ/kg DM)	20.4	18.6	20.8	21.9	21.3	21.5	19.6	20.3	19.8	20.1
Total SFA (% total FA)	39.6	41.4	29.9	28.6	21.7	20.3	20.6	20.8	29.8	28.9
Total MUFA (% total FA)	15.8	15.6	35.0	37.2	35.4	37.7	36.2	38.8	35.3	37.8
Total PUFA (% total FA)	16.2	14.3	25.0	27.1	39.7	41.6	40.1	40.4	24.7	26.1
Total LC-PUFA (% total FA)	28.4	28.6	10.0	7.1	3.1	0.3	3.1	0.0	10.2	7.3
Total n-3 (% total FA)	30.2	29.9	12.0	9.2	17.7	15.4	18.4	15.2	12.2	9.3
Total n-6(% total FA)	14.5	13.0	23.1	25.0	25.1	26.6	24.8	25.3	22.7	24.1

<sup>a</sup> Fish (Peruvian anchovetta) meal, Poultry offal meal and Fish (Peruvian anchovetta) oil : Skretting Australia, Hobart, TAS, Australia. <sup>b</sup> Wheat gluten and flour : Manildra, Auburn, NSW, Australia. <sup>c</sup> Microbial bioactive: Novacq™, CSIRO, Cleveland, QLD, Australia, PCT Patent AU 2008201886. <sup>d</sup>Lupin (*L. angustifolius*) kernel meal: Coorow Seed Cleaners, Coorow, WA, Australia. <sup>e</sup> Cholesterol : MP Bio, Aurora, OH, USA. <sup>f</sup> Banox-E™ : BEC Feed Solutions, Carole Park, QLD, Australia. <sup>g</sup> Astaxanthin (10%) as Carophyll Pink™ and Stay C™: DSM, Wagga Wagga, NSW, Australia. <sup>h</sup> Vitamin and mineral premix : Rabar, Beaudesert, QLD, Australia; Total FA: total fatty acids.

Table 4 Shrimp growth and feed utilisation parameters over the 42 day period of Experiment 1

	F45:M0	F45:M5	F45:M10	F20:M0	F15:M0	F10:M0	F5:M0	F0:M0	F5:M10	F0:M10	Pooled SEM
Initial weight (g/shrimp)	8.23	8.32	8.17	8.20	8.17	8.15	8.07	8.22	8.16	8.08	0.015
Weight (g/shrimp)	13.68 <sup>b</sup>	14.36 <sup>ab</sup>	14.62 <sup>a</sup>	13.38 <sup>b</sup>	13.14 <sup>bc</sup>	12.43 <sup>c</sup>	12.65 <sup>c</sup>	12.68 <sup>c</sup>	14.59 <sup>a</sup>	14.59 <sup>a</sup>	0.145
Gain (g/shrimp)	5.45 <sup>b</sup>	6.04 <sup>ab</sup>	6.45 <sup>a</sup>	5.18 <sup>b</sup>	4.97 <sup>bc</sup>	4.28 <sup>c</sup>	4.58 <sup>c</sup>	4.46 <sup>c</sup>	6.42 <sup>a</sup>	6.51 <sup>a</sup>	0.142
Gain rate (g/wk)	0.91 <sup>b</sup>	1.01 <sup>ab</sup>	1.08 <sup>a</sup>	0.86 <sup>b</sup>	0.83 <sup>bc</sup>	0.71 <sup>c</sup>	0.76 <sup>c</sup>	0.74 <sup>c</sup>	1.07 <sup>a</sup>	1.09 <sup>a</sup>	0.024
Feed fed (g/shrimp)	20.4 <sup>c</sup>	25.6 <sup>b</sup>	28.1 <sup>ab</sup>	31.8 <sup>a</sup>	28.0 <sup>ab</sup>	27.2 <sup>ab</sup>	26.0 <sup>b</sup>	24.0 <sup>bc</sup>	33.8 <sup>a</sup>	31.7 <sup>a</sup>	0.388
FCR (fed/gain)	3.92 <sup>a</sup>	4.26 <sup>a</sup>	4.44 <sup>ab</sup>	6.23 <sup>cd</sup>	5.72 <sup>c</sup>	6.42 <sup>d</sup>	5.69 <sup>c</sup>	5.46 <sup>bc</sup>	5.27 <sup>bc</sup>	4.87 <sup>b</sup>	0.107
Survival (%)	76.0 <sup>b</sup>	100.0 <sup>a</sup>	92.0 <sup>a</sup>	92.0 <sup>a</sup>	92.0 <sup>a</sup>	96.0 <sup>a</sup>	100.0 <sup>a</sup>	84.0 <sup>ab</sup>	100.0 <sup>a</sup>	96.0 <sup>a</sup>	6.59

FCR : Feed conversion ratio. Different superscripts within rows indicate significant differences (P<0.05). An absence of superscripts implies that there were no significant differences (P>0.05).

Table 5. Shrimp growth and feed utilisation parameters over the 42 day period of clear-water study in Experiment 2

	F50:O2:M0	F50:O2:M10	F10:O2:M0	F0:O2:M0	F10:O0:M0	F0:O0:M0	F10:O0:M10	F0:O0:M10	F10:O2:M10	F0:O2:M10	Pooled SEM
Initial weight (g/shrimp)	4.36	4.25	4.41	4.38	4.36	4.37	4.35	4.33	4.36	4.33	0.01
Weight (g/shrimp)	9.57 <sup>b</sup>	12.05 <sup>d</sup>	8.39 <sup>a</sup>	8.16 <sup>a</sup>	8.53 <sup>ab</sup>	7.35 <sup>a</sup>	12.17 <sup>d</sup>	10.51 <sup>c</sup>	12.79 <sup>d</sup>	12.07 <sup>d</sup>	0.28
Gain (g/shrimp)	5.21 <sup>b</sup>	7.80 <sup>d</sup>	3.98 <sup>a</sup>	3.78 <sup>a</sup>	4.18 <sup>ab</sup>	2.98 <sup>a</sup>	7.82 <sup>d</sup>	6.18 <sup>c</sup>	8.43 <sup>d</sup>	7.73 <sup>d</sup>	0.29
Growth rate (g/shrimp/wk)	0.87 <sup>b</sup>	1.30 <sup>d</sup>	0.66 <sup>a</sup>	0.63 <sup>a</sup>	0.70 <sup>ab</sup>	0.50 <sup>a</sup>	1.30 <sup>d</sup>	1.03 <sup>c</sup>	1.41 <sup>d</sup>	1.29 <sup>d</sup>	0.05
Feed fed (g/shrimp)	16.72 <sup>a</sup>	19.89 <sup>b</sup>	17.51 <sup>a</sup>	19.43 <sup>b</sup>	19.08 <sup>b</sup>	21.26 <sup>c</sup>	29.24 <sup>d</sup>	19.19 <sup>b</sup>	28.25 <sup>d</sup>	28.46 <sup>d</sup>	0.68
FCR (fed/gain)	3.21 <sup>b</sup>	2.55 <sup>a</sup>	4.40 <sup>c</sup>	5.14 <sup>d</sup>	4.57 <sup>c</sup>	7.12 <sup>e</sup>	3.74 <sup>a</sup>	3.11 <sup>ab</sup>	3.35 <sup>b</sup>	3.68 <sup>b</sup>	0.19
Survival (%)	93.3 <sup>bc</sup>	100.0 <sup>c</sup>	87.5 <sup>b</sup>	83.3 <sup>ab</sup>	91.7 <sup>bc</sup>	70.8 <sup>a</sup>	91.7 <sup>bc</sup>	91.7 <sup>bc</sup>	95.8 <sup>c</sup>	91.7 <sup>bc</sup>	1.13

Different superscripts within rows indicate significant differences (P<0.05). An absence of superscripts implies that there were no significant differences (P>0.05).

Table 6. Shrimp growth and feed utilisation parameters over the 63 day period of the green-water study in Experiment 3

	F50:O2:M0	F50:O2:M10	F10:O2:M0	F0:O2:M0	F10:O0:M0	F0:O0:M0	F10:O0:M10	F0:O0:M10	F10:O2:M10	F0:O2:M10	Pooled SEM
Initial weight (g/shrimp)	3.84	3.94	3.86	3.94	3.88	3.82	3.90	3.86	3.95	3.88	0.01
Weight (g/shrimp)	9.02 <sup>a</sup>	11.67 <sup>b</sup>	9.67 <sup>a</sup>	9.32 <sup>a</sup>	10.33 <sup>ab</sup>	9.78 <sup>a</sup>	12.24 <sup>bc</sup>	11.08 <sup>b</sup>	13.20 <sup>cd</sup>	13.40 <sup>d</sup>	0.23
Gain (g/shrimp)	5.18 <sup>a</sup>	7.73 <sup>b</sup>	5.81 <sup>a</sup>	5.38 <sup>a</sup>	6.45 <sup>ab</sup>	5.96 <sup>a</sup>	8.34 <sup>bc</sup>	7.23 <sup>b</sup>	9.25 <sup>cd</sup>	9.52 <sup>d</sup>	0.22
Growth rate (g/shrimp/wk)	0.58 <sup>a</sup>	0.86 <sup>b</sup>	0.65 <sup>a</sup>	0.60 <sup>a</sup>	0.72 <sup>ab</sup>	0.66 <sup>a</sup>	0.93 <sup>bc</sup>	0.80 <sup>b</sup>	1.03 <sup>cd</sup>	1.06 <sup>d</sup>	0.02
Feed fed (g/shrimp)	24.44 <sup>a</sup>	29.49 <sup>b</sup>	24.22 <sup>a</sup>	26.79 <sup>ab</sup>	27.83 <sup>ab</sup>	27.67 <sup>ab</sup>	32.20 <sup>bc</sup>	27.89 <sup>ab</sup>	34.24 <sup>c</sup>	30.37 <sup>bc</sup>	0.45
FCR (fed/gain)	4.74 <sup>bc</sup>	4.16 <sup>b</sup>	4.25 <sup>b</sup>	4.97 <sup>c</sup>	4.45 <sup>bc</sup>	4.70 <sup>bc</sup>	3.86 <sup>ab</sup>	3.89 <sup>ab</sup>	3.68 <sup>a</sup>	3.18 <sup>a</sup>	0.08
Survival (%)	96.7	97.8	82.2	96.7	94.4	96.7	98.9	98.9	94.4	94.4	0.68

Different superscripts within rows indicate significant differences (P<0.05). An absence of superscripts implies that there were no significant differences (P>0.05).



Table 7. Composition of shrimp from the clear-water study in Experiment 2

	F50:O2:M0	F50:O2:M10	F10:O2:M0	F0:O2:M0	F10:O0:M0	F0:O0:M0	F10:O0:M10	F0:O0:M10	F10:O2:M10	F0:O2:M10	Pooled SEM
Dry matter (%)	25.5	24.9	25.4	25.0	25.2	25.1	25.9	24.7	26.0	26.1	0.1
Protein (%)	20.3	19.7	19.0	21.6	19.4	22.0	21.4	21.3	21.5	22.7	0.2
Lipid (%)	2.1 <sup>ab</sup>	1.8 <sup>ab</sup>	2.1 <sup>ab</sup>	2.0 <sup>ab</sup>	1.9 <sup>ab</sup>	1.6 <sup>a</sup>	1.7 <sup>a</sup>	1.7 <sup>a</sup>	2.0 <sup>ab</sup>	2.2 <sup>b</sup>	0.0
Ash (%)	3.1 <sup>a</sup>	3.0 <sup>a</sup>	3.2 <sup>ab</sup>	2.9 <sup>a</sup>	3.1 <sup>a</sup>	3.1 <sup>a</sup>	3.3 <sup>ab</sup>	3.4 <sup>b</sup>	3.7 <sup>b</sup>	3.2 <sup>ab</sup>	0.0
Gross energy (kJ/g)	5.0	4.8	5.0	5.0	4.8	4.9	4.9	4.8	5.0	5.0	0.0
C14:0	1.7 <sup>c</sup>	1.6 <sup>c</sup>	1.1 <sup>bc</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.2 <sup>a</sup>	0.1 <sup>a</sup>	0.3 <sup>a</sup>	0.8 <sup>b</sup>	1.1 <sup>bc</sup>	0.1
C16:0	24.9 <sup>b</sup>	24.9 <sup>b</sup>	22.1 <sup>ab</sup>	22.4 <sup>ab</sup>	20.6 <sup>ab</sup>	19.9 <sup>a</sup>	19.1 <sup>a</sup>	19.1 <sup>a</sup>	22.9 <sup>ab</sup>	23.2 <sup>ab</sup>	1.0
C18:0	9.2 <sup>bc</sup>	8.7 <sup>b</sup>	6.5 <sup>b</sup>	2.3 <sup>a</sup>	9.6 <sup>c</sup>	10.0 <sup>c</sup>	9.0 <sup>bc</sup>	8.9 <sup>b</sup>	4.7 <sup>ab</sup>	0.0 <sup>a</sup>	0.6
Total SFA	36.9 <sup>c</sup>	36.3 <sup>bc</sup>	30.2 <sup>b</sup>	25.9 <sup>a</sup>	31.9 <sup>b</sup>	31.4 <sup>b</sup>	28.3 <sup>ab</sup>	28.8 <sup>ab</sup>	28.7 <sup>ab</sup>	24.8 <sup>a</sup>	1.4
C14:1	0.0	0.0	0.1	0.1	0.2	0.0	0.0	0.1	0.0	0.0	0.0
C16:1	2.3 <sup>c</sup>	2.1 <sup>c</sup>	1.9 <sup>c</sup>	1.6 <sup>bc</sup>	1.0 <sup>b</sup>	0.5 <sup>a</sup>	1.1 <sup>b</sup>	1.0 <sup>b</sup>	1.7 <sup>bc</sup>	1.6 <sup>bc</sup>	0.1
C18:1	16.0 <sup>a</sup>	19.2 <sup>a</sup>	30.1 <sup>b</sup>	36.0 <sup>cd</sup>	28.5 <sup>b</sup>	27.7 <sup>b</sup>	30.2 <sup>bc</sup>	30.4 <sup>bc</sup>	32.8 <sup>c</sup>	42.8 <sup>d</sup>	1.6
C20:1	0.5 <sup>b</sup>	0.8 <sup>c</sup>	0.2 <sup>ab</sup>	0.0 <sup>a</sup>	0.4 <sup>b</sup>	0.2 <sup>ba</sup>	0.0 <sup>a</sup>	0.2 <sup>ab</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0
Total MUFA	20.1 <sup>a</sup>	23.1 <sup>a</sup>	33.3 <sup>bc</sup>	39.0 <sup>c</sup>	31.8 <sup>b</sup>	29.2 <sup>b</sup>	32.1 <sup>b</sup>	32.3 <sup>b</sup>	35.7 <sup>bc</sup>	44.9 <sup>d</sup>	1.7
C18:2n-6	12.0 <sup>a</sup>	12.0 <sup>a</sup>	16.3 <sup>ab</sup>	17.4 <sup>ab</sup>	18.4 <sup>b</sup>	22.1 <sup>b</sup>	21.8 <sup>b</sup>	21.4 <sup>b</sup>	15.7 <sup>ab</sup>	16.3 <sup>ab</sup>	0.9
C18:3n-3	0.2 <sup>a</sup>	0.3 <sup>a</sup>	0.2 <sup>a</sup>	0.0 <sup>a</sup>	4.1 <sup>b</sup>	4.7 <sup>b</sup>	6.7 <sup>c</sup>	5.9 <sup>bc</sup>	0.0 <sup>a</sup>	0.8 <sup>a</sup>	0.4
Total PUFA	12.2 <sup>a</sup>	12.4 <sup>a</sup>	16.6 <sup>ab</sup>	17.4 <sup>ab</sup>	22.7 <sup>b</sup>	26.8 <sup>b</sup>	28.6 <sup>b</sup>	27.2 <sup>b</sup>	15.7 <sup>ab</sup>	17.1 <sup>ab</sup>	1.2
C20:4n-6	2.5 <sup>b</sup>	2.3 <sup>b</sup>	2.2 <sup>ab</sup>	2.3 <sup>b</sup>	1.4 <sup>a</sup>	3.1 <sup>c</sup>	2.4 <sup>b</sup>	2.3 <sup>b</sup>	1.7 <sup>a</sup>	1.7 <sup>a</sup>	0.1
C20:5n-3	12.7 <sup>c</sup>	12.1 <sup>c</sup>	8.5 <sup>b</sup>	8.3 <sup>b</sup>	5.3 <sup>ab</sup>	4.3 <sup>a</sup>	3.9 <sup>a</sup>	4.3 <sup>a</sup>	9.4 <sup>b</sup>	6.5 <sup>ab</sup>	0.5
C22:5n-3	1.2 <sup>b</sup>	1.1 <sup>b</sup>	0.3 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.1
C22:6n-3	14.4 <sup>c</sup>	12.9 <sup>c</sup>	7.6 <sup>ab</sup>	6.4 <sup>ab</sup>	5.4 <sup>a</sup>	4.6 <sup>a</sup>	4.5 <sup>a</sup>	4.8 <sup>a</sup>	8.8 <sup>b</sup>	5.0 <sup>a</sup>	0.6
Total LC-PUFA	30.8 <sup>c</sup>	28.4 <sup>c</sup>	19.9 <sup>b</sup>	17.7 <sup>b</sup>	13.5 <sup>ab</sup>	12.6 <sup>a</sup>	11.0 <sup>a</sup>	11.7 <sup>a</sup>	19.9 <sup>b</sup>	13.1 <sup>ab</sup>	1.2
Total n-6	14.5 <sup>a</sup>	14.3 <sup>a</sup>	18.7 <sup>ab</sup>	19.7 <sup>b</sup>	20.8 <sup>b</sup>	25.9 <sup>c</sup>	24.5 <sup>c</sup>	23.9 <sup>bc</sup>	17.4 <sup>ab</sup>	18.0 <sup>ab</sup>	1.0
Total n-3	28.5 <sup>c</sup>	26.5 <sup>c</sup>	17.8 <sup>b</sup>	15.4 <sup>ab</sup>	15.5 <sup>ab</sup>	13.6 <sup>a</sup>	15.1 <sup>ab</sup>	15.0 <sup>a</sup>	18.3 <sup>b</sup>	12.2 <sup>a</sup>	1.1
n-3 : n-6	1.96 <sup>c</sup>	1.85 <sup>c</sup>	0.95 <sup>b</sup>	0.78 <sup>ab</sup>	0.75 <sup>ab</sup>	0.52 <sup>a</sup>	0.62 <sup>a</sup>	0.63 <sup>a</sup>	1.05 <sup>b</sup>	0.68 <sup>a</sup>	0.1

Different superscripts within rows indicate significant differences (P<0.05). An absence of superscripts implies that there were no significant differences (P>0.05).

Table 8. Whole body composition of shrimp from the green-water study in Experiment 3

	F50:O2:M0	F50:O2:M10	F10:O2:M0	F0:O2:M0	F10:O0:M0	F0:O0:M0	F10:O0:M10	F0:O0:M10	F10:O2:M10	F0:O2:M10	Pooled SEM
Dry matter (%)	22.2 <sup>a</sup>	23.9 <sup>ab</sup>	25.1 <sup>ab</sup>	25.0 <sup>ab</sup>	24.2 <sup>ab</sup>	24.7 <sup>ab</sup>	26.6 <sup>b</sup>	25.4 <sup>b</sup>	26.2 <sup>b</sup>	25.7 <sup>b</sup>	0.35
Protein (%)	16.4 <sup>a</sup>	18.8 <sup>ab</sup>	18.1 <sup>ab</sup>	19.1 <sup>ab</sup>	17.2 <sup>ab</sup>	17.9 <sup>ab</sup>	20.6 <sup>b</sup>	19.5 <sup>ab</sup>	19.4 <sup>ab</sup>	19.9 <sup>ab</sup>	0.30
Lipid (%)	1.0 <sup>a</sup>	1.1 <sup>a</sup>	1.5 <sup>b</sup>	1.3 <sup>ab</sup>	1.4 <sup>ab</sup>	1.3 <sup>ab</sup>	1.6 <sup>b</sup>	1.3 <sup>ab</sup>	1.5 <sup>b</sup>	1.4 <sup>ab</sup>	0.06
Ash (%)	3.3 <sup>ab</sup>	3.6 <sup>ab</sup>	3.4 <sup>ab</sup>	3.6 <sup>ab</sup>	3.1 <sup>a</sup>	3.6 <sup>ab</sup>	3.6 <sup>ab</sup>	3.9 <sup>ab</sup>	4.0 <sup>b</sup>	3.7 <sup>ab</sup>	0.09
Gross energy (kJ/g)	3.8 <sup>a</sup>	4.5 <sup>ab</sup>	4.5 <sup>ab</sup>	4.6 <sup>ab</sup>	4.6 <sup>ab</sup>	4.5 <sup>ab</sup>	4.8 <sup>ab</sup>	4.4 <sup>ab</sup>	5.0 <sup>b</sup>	4.7 <sup>ab</sup>	0.07
C14:0	0.4 <sup>ab</sup>	0.5 <sup>b</sup>	0.4 <sup>ab</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.3 <sup>ab</sup>	0.3 <sup>ab</sup>	0.02
C16:0	12.0 <sup>ab</sup>	13.0 <sup>b</sup>	12.2 <sup>b</sup>	11.3 <sup>ab</sup>	10.9 <sup>ab</sup>	10.4 <sup>ab</sup>	10.2 <sup>a</sup>	9.8 <sup>a</sup>	11.6 <sup>ab</sup>	11.3 <sup>ab</sup>	0.19
C18:0	6.6 <sup>b</sup>	5.9 <sup>ab</sup>	5.6 <sup>ab</sup>	6.0 <sup>ab</sup>	5.3 <sup>a</sup>	5.4 <sup>a</sup>	5.4 <sup>a</sup>	5.7 <sup>ab</sup>	5.7 <sup>ab</sup>	5.8 <sup>ab</sup>	0.09
Total SFA	22.8 <sup>a</sup>	23.0 <sup>a</sup>	21.3	20.9	19.3 <sup>a</sup>	19.4 <sup>a</sup>	18.8 <sup>a</sup>	19.2 <sup>a</sup>	20.8 <sup>a</sup>	20.3 <sup>a</sup>	0.47
C14:1	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.01
C16:1	0.5 <sup>b</sup>	0.4 <sup>ab</sup>	0.2 <sup>a</sup>	0.3 <sup>ab</sup>	0.2 <sup>a</sup>	0.1 <sup>a</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.02
C18:1	9.8 <sup>a</sup>	10.6 <sup>ab</sup>	13.6 <sup>b</sup>	13.0 <sup>ab</sup>	13.5 <sup>b</sup>	13.4 <sup>ab</sup>	13.8 <sup>b</sup>	13.5 <sup>b</sup>	13.7 <sup>b</sup>	13.6 <sup>b</sup>	0.35
C20:1	0.7 <sup>ab</sup>	0.8 <sup>b</sup>	0.5 <sup>ab</sup>	0.5 <sup>ab</sup>	0.4 <sup>a</sup>	0.5 <sup>ab</sup>	0.4 <sup>a</sup>	0.3 <sup>a</sup>	0.5 <sup>ab</sup>	0.4 <sup>a</sup>	0.04
Total MUFA	12.7 <sup>a</sup>	13.6 <sup>ab</sup>	15.6 <sup>ab</sup>	15.2 <sup>ab</sup>	15.7 <sup>ab</sup>	15.9 <sup>b</sup>	16.1 <sup>b</sup>	16.1 <sup>b</sup>	16.0 <sup>b</sup>	15.6 <sup>ab</sup>	0.51
C18:2n-6	6.0 <sup>a</sup>	6.0 <sup>a</sup>	8.3 <sup>ab</sup>	8.5 <sup>ab</sup>	10.0 <sup>ab</sup>	10.8 <sup>b</sup>	10.1 <sup>ab</sup>	10.7 <sup>b</sup>	8.0 <sup>ab</sup>	8.6 <sup>ab</sup>	0.32
C18:3n-3	0.2 <sup>a</sup>	0.3 <sup>a</sup>	0.4 <sup>a</sup>	0.5 <sup>a</sup>	2.5 <sup>b</sup>	2.5 <sup>b</sup>	3.0 <sup>b</sup>	2.7 <sup>b</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.22
Total PUFA	6.2 <sup>a</sup>	6.3 <sup>a</sup>	8.7 <sup>ab</sup>	9.0 <sup>ab</sup>	12.5 <sup>b</sup>	13.3 <sup>b</sup>	13.1 <sup>b</sup>	13.5 <sup>b</sup>	8.4 <sup>ab</sup>	9.1 <sup>ab</sup>	0.54
C20:4n-6	10.1 <sup>ab</sup>	8.8 <sup>a</sup>	8.7 <sup>a</sup>	9.5 <sup>ab</sup>	11.1 <sup>ab</sup>	13.1 <sup>b</sup>	11.3 <sup>ab</sup>	13.1 <sup>b</sup>	8.1 <sup>a</sup>	9.1 <sup>a</sup>	0.33
C20:5n-3	18.0 <sup>b</sup>	15.8 <sup>ab</sup>	18.2 <sup>b</sup>	20.3 <sup>b</sup>	14.2 <sup>a</sup>	15.7 <sup>ab</sup>	13.0 <sup>a</sup>	15.1 <sup>a</sup>	18.2 <sup>b</sup>	19.7 <sup>b</sup>	0.45
C22:5n-3	1.6	1.8	2.1	2.1	1.6	1.9	1.7	2.0	2.1	2.2	0.05
C22:6n-3	24.7 <sup>bc</sup>	27.0 <sup>c</sup>	22.2 <sup>b</sup>	19.5 <sup>ab</sup>	21.2 <sup>b</sup>	15.9 <sup>a</sup>	21.4 <sup>b</sup>	16.2 <sup>a</sup>	23.1 <sup>b</sup>	20.8 <sup>ab</sup>	0.62
Total LC-PUFA	56.4	55.7	53.0	53.0	51.0	49.6	50.4	49.3	53.2	53.2	1.59
Total n-6	17.9 <sup>ab</sup>	16.8 <sup>a</sup>	18.5 <sup>ab</sup>	19.5 <sup>ab</sup>	23.2 <sup>ab</sup>	26.2 <sup>b</sup>	23.4 <sup>b</sup>	25.9 <sup>b</sup>	17.6 <sup>a</sup>	19.2 <sup>ab</sup>	0.72
Total n-3	44.8 <sup>a</sup>	45.2 <sup>a</sup>	43.1 <sup>a</sup>	42.5 <sup>a</sup>	40.3 <sup>a</sup>	36.7 <sup>a</sup>	40.1 <sup>a</sup>	36.9 <sup>a</sup>	44.0 <sup>a</sup>	43.2 <sup>a</sup>	1.41
n-3 : n-6	2.5 <sup>b</sup>	2.7 <sup>b</sup>	2.3 <sup>ab</sup>	2.2 <sup>ab</sup>	1.7 <sup>ab</sup>	1.4 <sup>a</sup>	1.7 <sup>ab</sup>	1.4 <sup>a</sup>	2.5 <sup>b</sup>	2.3 <sup>ab</sup>	0.24

. Different superscripts within rows indicate significant differences (P<0.05). An absence of superscripts implies that there were no significant differences (P>0.05).

Figure 1. Effect of fishmeal reduction in formulation on growth rate of shrimp in experiment 1. Shown are the linear regression through the means  $\pm$  SEM of the data.

