

1 **The response to, and recovery from maximum strength and power training in**
2 **elite track and field athletes**

3 Glyn Howatson,^{1,2} Raphael Brandon,³ & Angus M. Hunter⁴

4
5

6 ¹Department of Sport, Exercise and Rehabilitation, Northumbria University,
7 Newcastle upon Tyne, UK

8 ²Water Research Group, School of Environmental Sciences and Development,
9 Northwest University, South Africa

10 ³National Cricket Performance Centre, Loughborough, UK

11 ⁴School of Sport, University of Stirling, Stirling, UK.

12

13 **Copyright Human Kinetics. This version is as accepted for publication in**
14 ***International Journal of Sports Physiology and Performance*, 11 (3), pp. 356-362.**

15

16 **Corresponding author details:**

17 Glyn Howatson

18 Department of Sport, Exercise and Rehabilitation

19 Northumbria University

20 Newcastle-upon-Tyne

21 UK

22 Tel: +44(0)191 227 3573

23 Fax: +44(0)191 227 3500

24 glyn.howatson@northumbria.ac.uk

25

26 **Preferred Running Head:** Elite athletes: strength, power and recovery

27 **Word Count: 3866**

28 **Abstract**

29 There is a great deal of research on the responses to resistance training; however,
30 information on the responses to strength and power training conducted by elite
31 strength and power athletes is sparse. **Purpose:** To establish the acute and 24 hour
32 neuromuscular and kinematic responses to Olympic-style barbell strength and power
33 exercise in elite athletes. **Methods:** Ten elite track and field athletes completed a
34 series of 3 back squat exercises each consisted of 4 x 5 repetitions. These were done
35 as either strength or power sessions on separate days. Surface electromyography
36 (sEMG), bar velocity and knee angle was monitored throughout these exercises and
37 maximal voluntary contraction (MVC), jump height, central activation ratio (CAR)
38 and lactate were measured pre, post and 24 hours thereafter. **Results:** Repetition
39 duration, impulse and total work were greater ($p < 0.01$) during strength sessions, with
40 mean power being greater ($p < 0.01$) following the power sessions. Lactate increased
41 ($p < 0.01$) following strength but not power sessions. sEMG increased ($p < 0.01$) across
42 sets for both sessions, with the strength session increasing at a faster rate ($p < 0.01$) and
43 with greater activation ($p < 0.01$) by the end of the final set. MVC declined ($p < 0.01$)
44 following the strength and not the power session, which remained suppressed
45 ($p < 0.05$) 24 hours later; whereas CAR and jump height remained unchanged.
46 **Conclusion:** A greater neuromuscular and metabolic demand following the strength
47 and not power session is evident in elite athletes, which impaired maximal force
48 production up to 24 hours. This is an important consideration for planning concurrent
49 athletic training.

50 **Introduction**

51 Elite strength and power athletes use very specific resistance exercises to develop the
52 physical attributes of maximum strength and maximum power. Sessions comprising
53 high intensity (> 80% maximum load) and low repetitions (two to six) are often
54 performed to develop maximum strength.¹ Adaptations to maximum strength training
55 involve increased muscle fibre cross sectional area² and increased neural drive.³
56 Conversely, lower load exercises performed at higher velocities are performed to
57 develop power.⁴ Power-type training also improves neural drive, particularly motor
58 unit activation,⁵ and increases the ability to generate force during higher velocity,
59 dynamic movements.⁶ Consequently, the adaptations following resistance exercise
60 occur in both central and peripheral areas of the neuromuscular system and are largely
61 specific to the training performed.

62 Fatigue can be globally defined as an exercise-induced decline in the ability to
63 generate maximal voluntary muscle force⁷ and is associated with reductions in central
64 activation and neural drive, which are thought to provide (at least in part) the
65 necessary stimulus for adaptations to strength training⁸. In addition, increased surface
66 electromyographic (sEMG) amplitude during resistance exercise is indicative of
67 greater motor unit recruitment and therefore provides the required stimulus for an
68 adaptive response.^{9,10} Interestingly, the neuromuscular responses to strength and
69 power training have been examined in recreational athletes,^{8,10} but very little
70 information regarding elite athletes exist. Previous work has studied neuromuscular
71 fatigue and recovery following very high intensity (20 x 1RM) and high volume (10 x
72 10RM) resistance exercise sessions^{8,11} and found decreases in MVC for males and
73 females immediately following the sessions, with incomplete recovery 24 h post-
74 session.

75 A better understanding of the neuromuscular consequences following maximum
76 strength and power resistance exercise might better inform the training plan in order
77 to optimise adaptation, particularly in elite athletes. Additionally, the degree and
78 nature of fatigue will likely determine the recovery time required, influencing the type
79 of physical or technical training that is suitable following, or in conjunction with
80 resistance exercise. For example, knowledge of neuromuscular function 24 h
81 following maximum strength and power type resistance exercise will help coaches
82 plan day-to-day sessions, given the multiple types of training that can occur across the
83 cycle.

84 In the present study we had a rare opportunity to recruit elite athletes and expose them
85 to the 'typical' training stimulus of Olympic-style barbell exercises that are regularly
86 employed by elite track and field athletes (>10 sets)^{12,13} when targeting the
87 development of maximum strength and power. Therefore, the primary purpose of this
88 study was to examine the acute neuromuscular and kinematic responses to maximum
89 strength and power type resistance exercise and the subsequent 24 h recovery. The
90 second aim was to examine male and female responses within this elite group, which
91 might help inform whether the responses differ between sexes.

92

93 **Methods**

94 **Subjects**

95 Following institutional ethical approval, 10 performance programme athletes (Table
96 1) were recruited from UK Athletics Olympic Performance Centre, Lee Valley,
97 London and health-screened before providing written informed consent. All

98 volunteers were international standard sprinters or horizontal jumpers who regularly
99 partook in barbell strength training.

100 A schematic of the experimental design is presented in Figure 1. The trials were
101 completed following the competitive season when no sport-specific training was
102 occurring. Following familiarisation athletes performed a maximum strength or power
103 session in a randomised cross-over design within a seven day period. Each visit was
104 preceded with at least one rest day. Females were assessed during the luteal phase of
105 the menstrual cycle to limit hormonal variation on performance.

106 After arriving at the testing centre in a fasted state, blood lactate measures were taken
107 (Lactate Pro, ARK Corp, Japan) and consumed a standardised breakfast. The training
108 commenced with 10 minute warm up at 100 W on a cycle ergometer (Keiser M3,
109 Keiser Corp, USA). Subjects performed the pre-session neuromuscular (NM) tests,
110 comprising isometric knee extension force assessment (MVC), central activation ratio
111 assessment (CAR) and a vertical jump test (CMJ). The maximum strength or power
112 session was then performed; whole body barbell squat, split squat and press exercises.
113 These exercises were selected as commonly used exercises employed by UK strength
114 and conditioning coaches in delivering maximum strength and power programmes to
115 elite athletes.

116 During each session, surface electromyography (EMG), barbell displacement and
117 knee flexion (determined with electrogoniometry) were recorded. Based on a prior
118 pilot investigation, blood lactate was collected 4 minutes following the completion of
119 the final set to determine peak post-exercise lactate concentration. Ten minutes
120 following the session, CMJ, MVC and CAR tests were repeated to assess the
121 influence of the session on muscle function.¹⁴ On completion of each session subjects
122 provided a session RPE rating, using the Borg scale. To examine recovery following

123 the maximum strength and power sessions, subjects returned to the testing centre the
124 following day where MVC, CAR and CMJ assessments were performed following the
125 aforementioned warm up procedure.

126 Subjects attended familiarisation not more than seven days before the initial trial.
127 This included full instruction and practice of the MVC, CAR and CMJ assessments.
128 In addition, barbell loads were determined for the maximum strength session of squat,
129 split squat and push press. For each exercise, a series of incrementally loaded sets of
130 five repetitions were performed, starting at a self-selected 'moderate' load, separated
131 by three minutes rest between sets. At the end of each set, the intensity was rated
132 (RPE), using the Borg scale (6 to 20). The load corresponding to an active muscle
133 RPE = 16 or 17 (very hard) enabled the subjects' exercise to be matched for relative
134 intensity.¹⁵ Whilst percentage of repetition maximum loads are often used, the use of
135 RPE enables the determination of a load that is repeatable across all sets within the
136 session and akin to training methods used by UK elite track and field athletes.¹⁶

137 Immediately prior to the warm up subjects were fitted with an electrogoniometer
138 (TDA-100, Biopac Systems Inc., USA) attached to the lateral aspect of the left knee
139 to determine the beginning and end of the concentric phase of the movement and
140 synched with other instruments (such as EMG and the potentiometer) to determine the
141 kinematics and the relevant epoch could be identified across sessions. Barbell
142 displacement was measured using a potentiometer (Celesco PT5A, USA) attached to
143 the barbell to estimate power during the lifting phase.¹⁷ For the squat, speed squat,
144 split squat and split squat jump repetition, the mean power was calculated from the
145 whole concentric phase. For push and power press, the power calculation was limited
146 to the period where the knee angle was decreasing and displacement was increasing.
147 Power was calculated offline, where, force (load) = system mass \times (acceleration +

148 9.812), then, power = force (load) × velocity. This was used to compare changes in
149 power within sets during each session.

150 The duration of the combined lowering and lifting movement were used to define
151 repetition duration of each exercise. From repetition duration and the derived force
152 values, impulse was calculated as the integral of force over time. In addition, total
153 work was obtained as the integral of power. Mean set values for concentric mean
154 power, repetition duration, impulse and total work were determined from the average
155 of the five repetitions. Total work performed during the entire maximum strength and
156 power sessions were also compared; all calculations were computed off line
157 (AcqKnowledge® 3.8.1, Biopac Systems Inc., USA).

158 Surface EMG (sEMG) was continually monitored throughout the strength and power
159 sessions. The appropriate area was shaved, abraded and cleaned; 10-mm-diameter
160 electrodes (PNS Dual Element Electrode; Vermed, Vermont, USA), with 10-mm
161 inter-electrode distance were attached to the right *vastus lateralis* with the ground
162 electrode attached to the patella.¹⁸ The EMG data were sampled at 2000 Hz and
163 filtered using 1 Hz - 500 Hz band pass filter. The root-mean-squared (RMS)
164 amplitude was processed from the raw EMG amplitude using a 100 ms, overlapping
165 window. RMS amplitude values were normalised to the value obtained from
166 repetition one within each set.

167 The subjects performed the knee extension MVC force and CAR test as one combined
168 assessment, using an isokinetic dynamometer (Kin Com, Chattanooga, USA).
169 Subjects were positioned according to the manufacturer's recommendations with 70°
170 of knee flexion from full extension. Following three warm up contractions of
171 increasing intensity, subjects were instructed to produce three, 7 s 'ramp' contractions
172 (whereby maximum force was reached within 4 s) with 60 s rest between test

173 contractions. Visual feedback, and strong verbal encouragement was provided
174 throughout. The trial resulting in greatest voluntary force was used for data analysis
175 and was processed as the mean value from a 200 ms window centred upon the peak
176 force value.

177 During one randomly chosen MVC, and without warning, central activation ratio
178 (CAR) was determined by percutaneous stimulation (StimISOC, Biopac Systems Inc,
179 USA) of the femoral nerve with 250 ms, 100 Hz tetanic pulse train,¹⁹ the intensity of
180 which was determined during the familiarisation session; the optimum position was
181 marked to ensure consistent placement on subsequent visits. The CAR was
182 determined from the peak force prior to stimulation and the peak force during the
183 stimulation;²⁰ from this, $CAR = (MVC \text{ force} / \text{superimposed stimulated force}) \times 100$.

184 Three maximal counter movement jumps (CMJ) were then performed with a 30 s
185 pause between each. Subjects held a wooden stick across the shoulders during the
186 jump to remove extraneous use of the arms. The stick also enabled the potentiometer
187 (Celesco PT5A, USA) to directly measure jump height. The peak CMJ height from
188 the three trials was used for data analysis.

189 Following the warm up and pre-session assessments, two sets of squat were
190 performed at a self-determined 'moderate' intensity in order to provide an exercise-
191 specific warm up prior to the sessions. A series of three exercises Using Olympic
192 barbells, each exercise consisting of four sets of five repetitions, with three minutes
193 rest between sets were completed, which accurately reflected elite training sessions
194 for strength and power athletes on the Team GB Olympic track and field programme.
195 Constant feedback was given to the athletes regarding range of movement, timing and
196 speed during both sessions.

197 During the maximum strength session, the squat, split squat and push press were
198 performed, in that order, using the pre-determined loads. The squat was performed
199 with the bar resting across the shoulders, feet shoulder width apart and squatting
200 down until the hips lowered to below knee and then standing back up during the
201 concentric phase. The split squat also involved squatting and raising, with the barbell
202 resting upon the shoulders; however, the right foot was forward with the left foot
203 back. The movement involved squatting down, flexing at the hip and knee of the
204 front leg and the knee of the back leg, whilst keeping the trunk upright. The push
205 press was performed with feet shoulder width apart and holding the barbell in the
206 hands across the front of the shoulders. The movement comprised a small squat down
207 followed by synchronously pressing the bar over the head whilst standing back up.
208 Subjects were instructed to perform the concentric phase of all movements over two
209 seconds, which was controlled by a metronome.

210 During the power session the speed squat, split squat jump and power press were
211 performed with 30% of the barbell load used in the maximum strength session.^{5,21}

212 During the speed squats, subjects were instructed to perform the eccentric and
213 concentric repetition cycle as fast as possible, with a minimal jump in order to
214 maximise repetition speed. Subjects performed the split squat jumps and power press
215 with maximum acceleration in the concentric phase, following a controlled lowering
216 phase.

217 All data are presented as mean \pm SD. Differences between sessions for MVC, CAR,
218 and CMJ were examined using a two factor (session, 2 \times time, 3) repeated measures
219 ANOVA, with one less level for lactate. Differences in sEMG between and within
220 session a three factor (session, 2 \times set, 4 \times rep, 5) ANOVA was used. A further three
221 factor ANOVA (session, 2 \times exercise, 3 \times set, 4) was used to determine differences

222 in power, impulse, repetition duration and total work. Where necessary, effects were
223 followed by Tukey's *post-hoc* tests. Given the gender differences, we explored post-
224 session changes in MVC between male and female athletes using an independent
225 samples t-test. In addition, regression analysis assessed the relationship between the
226 post-session relative MVC and squat load, and also the relationship between the post-
227 session relative MVC and the system mass (Barbell Load + (0.88 x body mass))
228 load used during the power sessions, expressed in relation to the maximum strength
229 load. All data were performed on statistical software (Minitab v.15, USA);
230 significance was accepted at $\alpha = 0.05$. Where appropriate, 95% lower and upper
231 confidence intervals (CI) and Cohen's *d* effect sizes (ES) calculated by: Cohen's $d =$
232 $Mean_1 - Mean_2 / SD_{pooled}$, where $SD_{pooled} = \sqrt{[(SD_1^2 + SD_2^2) / 2]}$. ES were then
233 interpreted as $\leq 0.2 =$ trivial, $0.2-0.5 =$ small, $0.5-0.8 =$ moderate, $\geq 0.8 =$ large. Where
234 significant and non significant main effects are described the mean ES and CI,
235 between conditions, across all time points are presented.

236

237 **Results**

238 Significant interaction between the exercises and sessions were found for repetition
239 duration ($F = 18.13$, $p < 0.001$) impulse ($F = 97.47$, $p < 0.001$), total work ($F = 8.38$, p
240 $= 0.004$) and mean power ($F = 77.37$, $p < 0.001$) – Table 2. *Post hoc* tests showed
241 impulse and repetition duration were greater and power was less ($p < 0.01$; impulse
242 speed squat ES: 3.6, CI: 2.06 to 4.82; split squat ES: 4.4, CI: 2.62 to 5.76; press push
243 ES: 2.3, CI: 1.13 to 3.38) during all three exercises in the maximum strength session
244 compared to the equivalent power session. *Post hoc* tests between equivalent
245 exercises showed that only squat exercise had greater total work than the speed squat.
246 However, the total work performed during the entire maximum strength session was

247 significantly greater ($F = 3.65$, $p = 0.008$; ES: 1.34, CI: 0.32 to 2.29) than the power
248 session.

249 Lactate (Figure 2.) showed a session and a session by time interaction effect ($F = 57.56$,
250 $p < 0.001$). Lactate values post- maximum strength session were higher than baseline
251 (6.86 ± 2.2 versus 0.94 ± 0.2 mmol.L⁻¹; ES: 3.8, CI: 2.2 to 5.06), whilst post-power
252 session lactate was relatively unchanged (0.89 ± 0.2 versus 1.2 ± 0.3 mmol.L⁻¹; ES: 1.2,
253 CI: -2.11 to -0.22). Post-session RPE was higher ($t = 11.92$, $p = 0.012$; ES: 2.8, CI: 1.46
254 to 3.87) following the strength (16.5 ± 1.8) versus the power session (11.2 ± 2.0).

255 Repetition sEMG (Figure 3.) increased within sets for both sessions ($F = 18.76$, $p <$
256 0.001 ; ES: 0.28, CI: 0.035 to 0.36). For example, during set four of the maximum
257 strength session, sEMG increased (relative to repetition one of each set) to $116.5 \pm$
258 14.3% , $125.8 \pm 15.6\%$ and $125.8 \pm 15.6\%$ for squat, split squat and push press,
259 respectively. During set four of the power session RMS increased to $121.1 \pm 18.5\%$,
260 $102.0 \pm 13.1\%$, and $112.7 \pm 16.2\%$ for speed squat, split squat jump and power press,
261 respectively. There were session by set interaction effects ($F = 4.78$, $p = 0.029$); *post-*
262 *hoc* tests revealed repetitions four and five were higher to repetition one ($p < 0.01$; mean
263 ES: 0.26, mean CI: 0.0255 to 0.3472) during all sets of maximum strength session,
264 whereas repetitions four and five were only different during set one of the power session.

265 There were no differences in pre-session values between maximum strength and
266 power session on any variable showing that athletes were in a similar physical
267 condition between sessions (MVC- ES: 0.03, CI: -0.92 to 0.83; CAR-ES: 0.34, CI:
268 -1.21 to 0.55; CMJ – ES: 0.19, CI: -0.69 to 1.07) (Table 3). There was a significant
269 effect of the session on MVC ($F = 9.37$, $p = 0.014$) and across time ($F = 7.83$, $p =$
270 0.004). *Post-hoc* analysis revealed that following the strength session MVC was lower
271 than pre strength MVC ($p < 0.01$; ES: 0.4, CI: -0.49 to 1.28) with no significant

272 decline (ES: 0.17, CI: -0.71 to 1.04) demonstrated following the power session.
273 Importantly, MVC was still depressed by a small amount 24 h following strength
274 session ($p < 0.05$; ES: 0.23, CI: -0.66 to 1.10), whereas the restoration of MVC at 24
275 h post-power session was largely resolved. There were no main effects or interactions
276 for CAR (ES: 0.24 CI: -1.11 to 0.65) or CMJ height (ES: 0.13 CI: -0.75 to 1.00).

277 The relative change in MVC for male ($n = 6$) and female ($n = 4$) subjects, expressed
278 as a percentage of pre-session values, was $89.9 \pm 9.3\%$ versus $86.9 \pm 5.8\%$ post the
279 maximum strength session and $98.6 \pm 5.9\%$ versus $86.4 \pm 7.5\%$ post the power
280 session, respectively. T-test revealed the female subjects suffered significantly
281 greater decrement, albeit by a trivial amount, in MVC post-power session compared
282 to the males ($t = 2.88$, $p = 0.02$; ES: 1.8, CI: -0.23 to -0.00979).

283 There was a significant relationship ($r^2 = 0.705$, $p < 0.01$) between the athletes' strength
284 during the squat exercise (determined as the system mass (bar mass + body mass)
285 divided by body mass) and relative change in MVC (Figure 4A). In addition, there was
286 a significant relationship ($r^2 = 0.744$, $p < 0.001$) between the change in post-power
287 session MVC (Figure 4B) and the relative load used during the power session in
288 comparison to maximum strength session.

289

290 Discussion

291 This study investigated the consequences of strength and power sessions in elite track
292 and field athletes. These data show increased neuromuscular activity throughout both
293 training sessions, but there is an acute and prolonged (24 h post-session) reduction in
294 function following the maximum strength training results, but not power.

295 The important findings were reduced MVC immediately following strength but not
296 power sessions, whilst there were no changes in CAR or CMJ height. This is most
297 readily explained by greater total work during strength vs. power session,
298 accompanied by greater post-session lactate, suggesting greater metabolic challenge.
299 This difference in decline following maximum strength and power concur with our
300 previous results¹⁶ and from those studies using machine-based exercise sessions with
301 non-elite exercisers.²¹ The reduction in MVC with no change in CAR suggests
302 peripheral rather than central fatigue mechanisms were the dominant cause of MVC
303 decline.²² This observation disagrees with previous work,⁸ based upon sEMG
304 changes, that nervous system fatigue occurred.²¹ However, other research using
305 similar methods to the present study found no evidence of central fatigue following
306 three sets of elbow flexion resistance exercise.²³ Consequently, comparing with these
307 data on non-elite subjects might be somewhat futile given the obvious differences in
308 training status; nonetheless it seems that structured resistance exercise, designed for
309 maximum strength adaptation, result primarily in peripheral fatigue that is not evident
310 following sessions designed to enhance maximum power.

311 Although previous findings are somewhat contradictory, the sport-specific training
312 response in the current investigation has hitherto, not been reported for elite athletes.
313 Muscle function assessments were conducted 10-minutes following completion of the
314 final set, rather than immediately following the final repetition where ischemia or
315 muscle pH changes could influence action potential propagation and contractile
316 function, thus influencing outcome measures.²⁴ The choice of assessment timing
317 could influence CAR measurement, as central fatigue recovers quickly post-
318 exercise.²⁵ Therefore, it is conceivable that central fatigue was present immediately
319 after training, but was resolved before the 10-minute post-exercise assessment.

320 Nonetheless, it was surprising that high intensity resistance exercise did not result in
321 central fatigue given the neuromuscular system is heavily implicated in adaptation to
322 maximum strength and power training.^{3,26} It is also conceivable that central fatigue
323 *per se* is not necessary to induce an adaptive response and we speculate that the
324 ability to recruit the target areas of the neuromuscular system during the session is
325 arguably the most important element of resistance exercise in elite athletes.

326 During both the maximum strength and power sessions, RMS increased within the
327 sets, with no concomitant change in mean power. This indicates greater recruitment
328 and/or firing rates, possibly of larger non-fatigued motor units. Somewhat intuitively,
329 RMS increased more during strength than the power sessions, suggesting greater
330 neuromuscular activation to maintain repetition performance, compared to lower load
331 higher velocity repetitions.^{25,27} The peripheral fatigue indicated by decreased MVC,
332 could be attributed to localised muscle damage, although in trained athletes the
333 repeated bout effect will limit the damage response.²⁸ Nonetheless, reporting of
334 muscle soreness at 24 h might have provided indirect evidence of muscle damage.
335 Alternatively, the accumulation of metabolites (evidenced by modest elevations in
336 blood lactate) affected the release and re-uptake of Ca^{2+} in the sarcoplasmic reticulum
337 and thereby impaired excitation-contraction coupling.²⁹ In either case, greater
338 peripheral fatigue following maximum strength-type training provides a larger
339 stimulus for muscle protein synthesis,³⁰ although both high and low load training has
340 been shown to increase skeletal muscle hypertrophy in trained men.³¹

341 MVC was still depressed by ~6% below pre-session force following the maximum
342 strength session which has important implications for subsequent exercise
343 prescription and training programme design considerations for elite athletes. Previous
344 research on non-elite populations^{8,11,21} showed similar, but nonetheless larger effects;

345 however the load, used in these studies would not be used in optimal elite strength
346 training programmes. In addition, muscle function changes in post-session relative
347 MVC of male vs. female are somewhat limited by low numbers, but are still insightful
348 given the elite nature of these athletes. Although, all of them showed reduced MVC
349 post strength session (11-12%), only females reduced MVC post-power session by a
350 similar amount, whereas the males maintained MVC. However, previous findings,
351 using non-elite subjects showed similar reductions in MVC for both genders post
352 power sessions²¹ and that when females are matched for strength, there were no
353 difference in fatigue to men.³² Therefore, as we did not match strength it is possible
354 that individual strength accounted for the difference in NM fatigue. Furthermore, we
355 showed a strong relationship between strength and the relative change in MVC
356 following the power ($r = 0.84$), but not strength session. This is likely to be from
357 varied relative loading level used between subjects during the power session.
358 Furthermore, the system mass load lifted during the power session (relative to the
359 loaded lifted during the maximum strength session) was inversely related to the
360 degree of change in MVC post power session (Figure 4B). Consequently, it is likely
361 that MVC force reduction differences of male vs. female is weaker, lighter subjects
362 were working 'relatively' harder during the power session than stronger, heavier
363 athletes. Definitive gender differences are not possible to glean from these data in
364 elite athletes, but it does highlight the importance of training intensities in a 'system
365 mass' term because of the practical issues in exercise prescription. Setting load levels
366 for power sessions as percentages of system mass loads might help ensure individuals
367 train at a similar relative intensity.

368

369 **Practical applications and Conclusion**

370 In summary, these data provide new information of the fatigue and recovery
371 following resistance exercise sessions designed to improve maximum strength and
372 power in elite track and field athletes. The findings show that 12 sets of maximum
373 strength resistance exercise results in reduced force generating capacity that take more
374 than 24 h to be resolved, whereas force is largely unchanged following power
375 sessions. This is likely from higher intensity and time under tension during the
376 maximum strength session (impulse) and total work done. The study provides
377 valuable information for athletes, coaches and practitioners when planning the
378 training programme to understand the consequences of engaging elite athletes in
379 strength and power resistance exercise. Specifically, in the day following maximum
380 strength training there is likely to be an impairment of maximum strength; therefore
381 practitioners should be mindful of appropriate programming particularly where
382 subsequent maximal or perimaximal efforts might be required.

383

384 **Acknowledgements**

385 Gratitude to extended to UK Sport's Graduate Innovations Programme for funding the
386 research project.

- 388 1. Crewther B, Cronin J, Keogh J. Possible stimuli for strength and power
389 adaptation: acute mechanical responses. *Sports Med.* 2005; 35:967-989.
- 390 2. Andersen LL, Andersen JL, Magnusson SP, Suetta C, Madsen JL, Christensen
391 LR, Aagaard P. Changes in the human muscle force-velocity relationship in
392 response to resistance training and subsequent detraining. *J Appl Physiol.*
393 2005;99: 87-94.
- 394 3. Aagaard P, Simonsen E, Andersen JL, Magnusson SP, Dyhre-Poulsen P.
395 Increased rate of force development and neural drive of human skeletal muscle
396 following resistance training. *J Appl Physiol.* 2002;93:1318-1326.
- 397 4. McBride JM, Triplett-McBride T, Davie A, Newton RU. The effect of heavy- vs.
398 light-load jump squats on the development of strength, power, and speed. *J*
399 *Strength Cond Res.* 2002;16:75-82.
- 400 5. Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour
401 contribute to the increase in contraction speed after dynamic training in humans.
402 *J Physiol.* 1998;513: 295-305.
- 403 6. Cormie P, McGuigan MR, Newton RU. Adaptations in athletic performance after
404 ballistic power versus strength training. *Med Sci Sports Exerc.* 2010;42:1582-
405 1598.
- 406 7. Gandevia SC. Central factors in human muscle fatigue. *Physiol Rev.* 2001;81:
407 1725-1789.
- 408 8. Hakkinen K. Neuromuscular fatigue in males and females during strenuous heavy
409 resistance loading. *Electromyogr Clin Neurophysiol.* 1994; 34: 205-214.
- 410 9. Pincivero DM, Gandhi V, Timmons M, Coelho AJ. Quadriceps femoris
411 electromyogram during concentric, isometric and eccentric phase of fatiguing
412 dynamic knee extensions. *J Biomech.* 2006;39:246-54.
- 413 10. Ahtiainen J, Hakkinen K. Strength Athletes are capable to produce greater muscle
414 activation and neural fatigue during high-intensity resistance exercise than
415 nonathletes. *J Strength Cond Res.* 2009;23:1129-1134.
- 416 11. Hakkinen K. Neuromuscular fatigue and recovery in male and female athletes
417 during heavy resistance exercise. *Int J Sports Med.* 1993;14:53-59.
- 418 12. Andersen JL, Aagaard P. Myosin heavy chain IIX overshoot in human skeletal
419 muscle. *Muscle Nerve.* 2000;23: 1095-1104.
- 420 13. Campos GE, Luecke TJ, Wendeln HK, Toma K, Hagerman FC, Murray
421 TF, Ragg KE, Ratamess NA, Kraemer WJ, Staron RS. Muscular adaptations in
422 response to three different resistance-training regimens: specificity of repetition
423 maximum training zones. *Eur J Appl Physiol.* 2002;88:50-60.
- 424 14. Chiu LZ, Fry AC, Schilling BK, Johnson EJ, Weiss LW. Neuromuscular fatigue
425 and potentiation following two successive high intensity resistance exercise
426 sessions. *Eur J Appl Physiol.* 2004;92:385-392.
- 427 15. Gearhart RE, Goss FL, Lagally KM, Jakicic JM, Gallagher J, Robertson RJ.
428 Standardized scaling procedures for rating perceived exertion during resistance
429 exercise. *J Strength Cond Res.* 2001;15:320-325.
- 430 16. Brandon R, Howatson G, Hunter AM. Neuromuscular response differences to
431 power vs strength back squat exercise in elite athletes. *Scand J Med Sci Sports.*
432 (Epub ahead of press - 10.1111/sms.12289)
- 433 17. Cormie P, Deane R, McBride JM. Methodological concerns for determining
434 power output in the jump squat. *J Strength Cond Res.* 2007; 21:424-430.

- 435 18. Hermens HJ, Freriks B, Disselhorst-Klug C, Rau. (1999). Development of
436 recommendations for SEMG sensors and sensor placement procedures. *J*
437 *Electromyogr Kinesiol.* 2000; 10:361-74.
- 438 19. Lattier G, Millet GY, Martin A, Martin V. Fatigue and recovery after high-
439 intensity exercise part I: neuromuscular fatigue. *Int J Sports Med.* 2004;25:450-
440 456.
- 441 20. Nybo L, Nielsen B. Hyperthermia and central fatigue during prolonged exercise
442 in humans. *J Appl Physiol.* 2001;91:1055-1060.
- 443 21. Linnamo V, Hakkinen K, Komi PV. Neuromuscular fatigue and recovery in
444 maximal compared to explosive strength loading. *Eur J Appl Physiol Occup*
445 *Physiol.* 1998;77:176-181.
- 446 22. Kent-Braun, J. (1999). Central and peripheral contributions to muscle fatigue in
447 humans during sustained maximal effort. *Eur J Appl Physiol.* 1999;80:57-63.
- 448 23. Tran QT, Docherty D, Behm D. The effects of varying time under tension and
449 volume load on acute neuromuscular responses. *Eur J Appl Physiol.*
450 2006;98:402-410.
- 451 24. Fitts RH. Cellular mechanisms of muscle fatigue. *Physiol Rev.* 1994;74:49-94.
- 452 25. Sogaard K, Gandevia S, Todd G, Petersen N, Taylor JT. The effect of sustained
453 low-intensity contractions on supraspinal fatigue in human elbow flexor muscles.
454 *J Physiol.* 2006;572:511-523.
- 455 26. Aagaard P, Simonsen E, Andersen JL, Magnusson SP, Halkjaer-Kristensen J,
456 Dyhre-Poulsen P. Neural inhibition during maximal eccentric and concentric
457 quadriceps contraction: effects of resistance training. *J Appl Physiol.*
458 2000;89:2249-2257.
- 459 27. Moritani T, Muro M, Nagata A. Intramuscular and surface electromyogram
460 changes during muscle fatigue. *J Appl Physiol.* 1986; 60:1179-1185.
- 461 28. Howatson and van Someren (2008). Prevention and treatment of exercise-induce
462 muscle damage. *Sports Med.* 2008;38:483-503.
- 463 29. Hill C, Thompson M, Ruell P, Thom J, White M. Sarcoplasmic reticulum
464 function and muscle contractile character following fatiguing exercise in humans.
465 *J Physiol.* 2001;531:871-878.
- 466 30. Schoenfeld, B. J. (2010). The Mechanisms of Muscle Hypertrophy and Their
467 Application to Resistance Training. *J Strength Cond Res.* 2010;24:2857-2872.
- 468 31. Schoenfeld BJ, Peterson MD, Ogborn D, Contreras B, Sonmez GT. Effects of
469 Low- Versus High-Load Resistance Training on Muscle Strength and
470 Hypertrophy in Well-Trained Men. *J Strength Cond Res.* (Epub of ahead:
471 10.1519/JSC.0000000000000958)
- 472 32. Hunter SK, Critchlow A, Shin IS, Enoka RM. Fatigability of the elbow flexor
473 muscles for a sustained submaximal contraction is similar in men and women
474 matched for strength. *J Appl Physiol.* 2004;96:195-202.

475 **Figure Legends:**

476

477 **Figure 1.** Timed summary of the procedures assessing maximum strength and power
478 sessions.

479

480 **Figure 2.** Pre- and post-session lactate during maximum strength and power sessions.
481 Values given as mean \pm SD, n = 10. ** Significant time difference for lactate, $p < 0.01$
482 and \$ significant interaction effect, $p < 0.01$.

483

484 **Figure 3.** Normalised RMS amplitude within sets of maximum strength and power
485 exercises. Mean values given relative to repetition one of each set, n = 10. *
486 Significant difference between repetitions, $p < 0.001$, \$ significant interaction effect
487 between set x repetition and exercise x repetition, $p < 0.05$.

488

489 *NB: Split squat set 1 and press set 1 were sets 5 & 9 of the sessions, respectively.*

490

491

492 **Figure 4.** Relationships between relative change in MVC post power session and
493 load level. A) Relative change in MVC versus relative squat load expressed as
494 bodyweights (BW), where post MVC = $0.413 + 0.225 \times \text{SM load}$. ($r^2 = 0.705$, $p < 0.01$).
495 Jagged line shows 95% confidence intervals. B) Relative change in MVC versus load
496 lifted during power session relative to maximum strength session (%), where post MVC =
497 $1.88 - 1.58 \times \text{relative load}$ ($r^2 = 0.744$, $p < 0.001$). Jagged line shows 95% confidence
498 intervals.

499 Table 1. Subjects' physical characteristics; Values are given as mean \pm SD.
500

	Age (years)	Body mass (kg)	100m best time (s)	Squat 1RM (kg)	MVC force (N)
Male, n = 6	28 \pm 2	81.2 \pm 12.2	10.44 \pm 0.37	190.0 \pm 38.0	1092.5 \pm 245.1
Female, n = 4	26 \pm 5	60.0 \pm 3.7	11.73 \pm 0.34	107.5 \pm 12.0	821.0 \pm 102.8

501

502 Table 2. Repetition duration, impulse, mean power and total work data during squat,
 503 split squat and press during maximum strength and power sessions. Values are given
 504 as mean \pm SD. Significant session x exercise interaction effects $p < 0.01$ were found for
 505 all variables with * significant difference between exercises within the sessions
 506 shown, $p < 0.01$. ** Significantly different between strength and power session,
 507 $p < 0.001$.

508
 509
 510

	Repetition Duration (s) **	Impulse (N.s) **	Mean Power (W)**	Total work (J)
Maximum Strength (n=10)				
Squat (S)	3.4 \pm 0.28	5676 \pm 1854	528 \pm 245	1791 \pm 756*
Split Squat (SS)	3.3 \pm 0.3	4578 \pm 1175	340 \pm 130	1089 \pm 370
Press (Pr)	1.9 \pm 0.7*	2072 \pm 806*	988 \pm 389*	1074 \pm 334
Maximum Power (n=10)				
Squat (S)	0.8 \pm 0.2	934 \pm 228	1234 \pm 385*	1004 \pm 344
Split Squat (SS)	0.8 \pm 0.2	887 \pm 206	1760 \pm 582*	1119 \pm 422
Press (Pr)	0.6 \pm 0.2	692 \pm 194	3297 \pm 1298*	1049 \pm 368


511

512 Table 3. Maximum voluntary contraction (MVC), central activation ratio (CAR), and
 513 counter movement jump (CMJ) height at pre, post and 24 h post strength and power
 514 sessions. Values are given as mean \pm SD, n = 10. **Significant difference (p < 0.01)
 515 to pre strength session MVC and post power session MVC; * Significant different to
 516 pre-strength MVC (p < 0.05).
 517

		Strength	Power
MVC (N)	pre	975.5 \pm 246.7	983.9 \pm 237.8
	post	871.9 \pm 255.2**	937.6 \pm 298.7
	24 h	920.5 \pm 226.2*	953.3 \pm 233.8
CAR (%)	pre	92.6 \pm 4.4	94.2 \pm 4.9
	post	93.5 \pm 3.0	95.4 \pm 3.9
	24 h	92.7 \pm 4.7	93.2 \pm 4.2
CMJ Height (cm)	pre	49.1 \pm 9.8	47.1 \pm 10.5
	post	47.8 \pm 10.4	47.4 \pm 11.1
	24 h	48.6 \pm 8.9	48.7 \pm 8.8


518

519 **Figure 1.**




520

521 **Figure 2.**
522





523

524 **Figure 3.**



525
526

527 **Figure 4.**



528