

1 Plasticity and consistency of behavioural responses to predation risk in laboratory  
2 environments

3

4 Martina Heynen<sup>1,2,3</sup>(✉), Jost Borchering<sup>3</sup>, Nils Bunnefeld<sup>4</sup> and Carin Magnhagen<sup>5</sup>

5

6 <sup>1</sup>Department of Chemistry

7 Umeå University, Umeå, Sweden

8

9 <sup>2</sup>Department of Ecology and Environmental Sciences

10 Umeå University, Umeå, Sweden

11 E-mail: martina\_heyne@web.de

12

13 <sup>3</sup>Institute of Zoology of the University of Cologne

14 Department of General Ecology & Limnology

15 Ecological Field Station Grietherbusch, Rees, Germany

16 E-mail: Jost.Borchering@Uni-Koeln.de

17

18 <sup>4</sup>Biological and Environmental Sciences,

19 School of Natural Sciences, University of Stirling, Stirling, FK9 4LA, UK

20 E-mail: nils.bunnefeld@stir.ac.uk

21

22 <sup>5</sup>Department of Wildlife, Fish, and Environmental Studies

23 Swedish University of Agricultural Sciences, Umeå, Sweden

24 E-mail: Carin.Magnhagen@slu.se

25

26

27 Corresponding author: Martina Heynen, Department of Chemistry and Department of

28 Ecology and Environmental Sciences, Umeå University, Umeå, Sweden, e-mail:

29 [martina\\_heyne@web.de](mailto:martina_heyne@web.de)

30

31 **ABSTRACT**

32 The individual animal is currently a major focus of behavioural research and an increasing  
33 number of studies raise the question of how between-individual behavioural consistency and  
34 behavioural plasticity interact. Applying the reaction norm concept on groups, our study  
35 addresses both of these aspects in one framework and within an animal's natural social  
36 environment. Risk-taking behaviour in one-year-old perch (*Perca fluviatilis*) was assayed in  
37 aquarium experiments before and after the fish were subjected to the presence or absence of a  
38 piscivorous predator for three weeks. To analyse the inter-individual behavioural variation  
39 across the repeated measurements, we dissected the behavioural change across the predator  
40 treatment into individual constant and plastic components using hierarchical mixed effects  
41 models. During the predator treatment juvenile perch increased in boldness and decreased in  
42 vigilance, the magnitude of these behavioural changes was influenced by group composition.  
43 However, the behavioural changes were not influenced by the presence of a predator,  
44 indicating the difficulties in generating realistic long-term predation pressure in the  
45 laboratory. Individuals differed in the relative increase in boldness across the predator  
46 treatment and, thus, varied in the shape of their reaction norms. In accordance, the best linear  
47 unbiased predictors (BLUP), extracted from the random effects of separate linear mixed  
48 effects models for the data before and after the predator treatment were only weakly  
49 correlated. Hence, between-individual variation seems to change under laboratory conditions  
50 and therewith not necessarily represents the initially present "natural" variation, giving  
51 important implications for the conduction and interpretation of behavioural experiments.

52  
53 Keywords: reaction norm, *Perca fluviatilis*, phenotypic plasticity, boldness, personality,  
54 behavioural consistency  
55 Running title: Behavioural plasticity and consistency in the laboratory

56 **INTRODUCTION**

57 Behaviour is considered to be one of the most plastic phenotypic traits (Price et al. 2003) and  
58 many studies have documented the potential plasticity of behavioural traits in a variety of  
59 animals [mammals (Hayes & Jenkins 1997), cephalopods (Sinn et al. 2007), insects (Agrawal  
60 2001), birds (Cresswell & Quinn 2005; Miller et al. 2006), fish (Coleman & Wilson 1998)].  
61 Since a shift in focus to the individual, an increasing amount of studies have reported  
62 consistent behavioural differences within animal populations (reviewed in Sih et al. 2004;  
63 Bell 2007), but also between individual plasticity *per se* (Dingemanse et al. 2010). The  
64 observed consistent behavioural differences across time or situations have been ascribed to an  
65 animal's underlying distinct personality, also termed coping style, temperament or  
66 behavioural syndrome (Koolhaas et al. 1999; Sih et al. 2004; Dingemanse & Reale 2005).  
67 These, initially contradictory, findings raise the question of how inter-individual variability  
68 and intra-individual stability interact (Nussey et al. 2007; Dingemanse et al. 2010).  
69 Studies on fish have shown that behavioural stability along the shy-bold axis varies between  
70 species, but also within a species, when studied with regard to behavioural traits or situations.  
71 Wilson et al. (1993) found that pumpkinseed sunfish exhibited constant behavioural  
72 differences in the wild, and that these differences disappeared after the fish were held in the  
73 laboratory for some time. In sticklebacks, high predation risk altered the degree of aggression  
74 a stickleback displays towards conspecifics, and the observed overall decrease in boldness  
75 was due to selective predation, as the individual's degree of boldness did not change (Bell &  
76 Sih 2007).  
77 Clearly, behavioural differences may have important fitness consequences (Dingemanse et al.  
78 2004; Smith & Blumstein 2008). However, to assess the selective value of a trait, there is a  
79 need to understand how variable it is, both between and within individuals (Boake 1989;  
80 Hayes & Jenkins 1997; Dingemanse et al. 2010). One approach simultaneously addressing

81 variability and consistency is the reaction norm concept (Via et al. 1995; Nussey et a. 2007),  
82 where an individual's behaviour is tested repeatedly along an environmental gradient. The  
83 behavioural differences along the environmental gradient represent the plastic ability at the  
84 population level. Deconstructed to the individual level, the presence of an interaction between  
85 individual reaction and environment (I x E) suggests that individuals differ in their plastic  
86 ability (Nussey et al. 2007; Briffa et al. 2008; Martin & Reale 2008; Dingemanse et al. 2010).  
87 In juvenile Eurasian perch, *Perca fluviatilis* L., relative differences in boldness towards a  
88 predator have been shown to be consistent between individuals across different social  
89 contexts (Magnhagen & Bunnefeld 2009). Perch also display a high degree of behavioural  
90 plasticity, changing their behaviour with status of nourishment (Borcherding & Magnhagen  
91 2008), consumed prey (Heynen et al. 2010), social background (Magnhagen & Staffan 2005)  
92 or the experienced intensity of predation pressure (Magnhagen & Borcherding 2008;  
93 Hellström & Magnhagen 2011; Magnhagen et al. 2012).

94 The aim of the present study was to investigate whether the inter-individual behavioural  
95 variation within a group of perch changes with the adaptation to a novel situation.  
96 Furthermore, we wanted to test whether and how such changes, if any, are influenced by  
97 predation risk. Boldness towards a predator was measured in groups of juvenile perch, before  
98 and after participating in a three week tank treatment in absence or presence of an adult  
99 piscivorous perch. To analyse the consistency of the inter-individual variation across the  
100 repeated measurements, we compared the behavioural ranking within the same groups before  
101 and after the predator treatment and the variation in the groups that experienced a predator  
102 during the three weeks in the tank with those without predation risk. Using mixed effects  
103 models to analyse our data, we were able to dissect the obtained behavioural variation across  
104 the phenotypic response, at the group level, into individual constant and plastic components.  
105

106 **MATERIAL AND METHODS**

107 In June 2009 one-year-old juvenile perch (body length,  $X \pm SD$ ;  $90.3 \pm 6.4$  mm, weight  $7.4 \pm$   
108  $1.7$  g,  $N = 96$ ) were caught with a beach seine in Lake Ängersjön, close to the city of Umeå,  
109 Sweden ( $63^\circ 47'N$ ;  $20^\circ 17'E$ ). The fish were transported to Umeå Marine Research Centre  
110 (UMF, 45 km south of Umeå). In the 100-litre transport vessel, a pump run by a car battery was  
111 constantly recirculating and oxygenating the water. The fish were stocked to a circular tank (60  
112 cm high,  $0.471 \text{ m}^2$ ) with continuously running water (13L:11D, 14-15 °C) to acclimate to  
113 indoor conditions (5 or 10 days). They were fed daily with pre-frozen red chironomid larvae  
114 (6% of total body mass). The predators, older perch (body length,  $X \pm SD$ ;  $200.8 \pm 13.5$  mm,  
115  $N = 15$ ), were caught in a small stream near the laboratory and fed with earthworms daily.  
116 Fulton's condition factor was used as measure of the physical condition of the fish  
117 ( $K=10^5M*TL^{-3}$ , where  $M$ =weight in g and  $TL$ =total length in mm; Bagenal & Tesch 1978).  
118 After the experiments the tested fish were killed with an overdose of MS222. The predator  
119 perch were released into their natural habitat.

120

121 *Experimental design*

122 Before being handled all fish were sedated with MS222. The juvenile perch were marked with  
123 individual colour codes, generated with 4 different colours and carefully applied with a needle  
124 and tattoo colours (Tattoo-Flame©) on the upper and/or lower caudal fin. Subsequently, fish  
125 were weighed, measured and randomly assigned to groups of four ( $N = 24$  groups). Each  
126 group participated in two sets of behavioural experiments, the first one directly before and the  
127 second one directly after being exposed to a three week predator treatment in tanks (Table 1).  
128 After the first set of behavioural experiments were conducted, 4 groups were added to each of  
129 the 6 tanks used for the predator treatment ( $N = 24$  groups). Those six tanks (60 cm high,  
130  $0.471 \text{ m}^2$ , 50% cover with artificial vegetation) were circular and had continuously running

131 water. After three weeks all fish were collected from the tanks, sedated, weighed, measured  
132 and the behaviour of the same groups was re-assayed (second set of behavioural experiments),  
133 in the same way as during the first set of behavioural experiments. Due to the limited capacity  
134 of experimental aquaria, the study was conducted in two successive experimental blocks,  
135 starting five days apart. 12 groups (48 individuals) were tested in each block (see Table 1).

136

#### 137 *Predator treatment*

138 In the experimental block 1 two tanks were stocked with a predator and one tank was predator  
139 free, while the opposite was done for block 2, resulting in 3 predator and 3 non-predator tank  
140 treatments. In each experimental block juvenile perch were allowed to acclimate to the tank  
141 for one day before the predators were added. To generate a real impression of danger, but to  
142 minimize the consumption of prey individuals the predator size ratio was chosen close to the  
143 maximal border of ingestability (literature data: 0.45, Claessen et al. 2000; our data:  $X \pm SD$ ;  
144  $0.39 \pm 0.03$  mm, range 0.34 – 0.45) and the predator was only allowed to swim free in the  
145 tank for 6 days out of the 21 days of the experiment. After 6 days the predator was transferred  
146 into a transparent perforated plastic box (41 x 26 x 29 cm) within the tank, so that visual and  
147 olfactory predator cues were still present. The tanks without predators were treated the same,  
148 a net was swept through the tank and the box was opened and closed. During the tank  
149 treatment juvenile fish were fed with pre-frozen chironomid larvae daily (6% of total body  
150 weight) and the predators with earthworms every second day.

151

#### 152 *Experimental aquaria*

153 The experimental aquaria were 170 l (95 x 41 x 44 cm) and had continuously running water  
154 (17 °C; light regime 13L:11D). One-third of each aquarium was separated with a plastic net  
155 (mesh size 5 mm) and used for the predator, the rest for the group of juvenile perch. To

156 prevent the fish habituating to the predator during the behavioural assays an opaque plastic  
157 screen was placed next to the net. The water inlet was placed in the section with the small  
158 perch and the outlet in the predator section to minimize olfactory cues between observations.  
159 The bottom of the aquaria was covered with gravel. Artificial vegetation was provided in the  
160 predator space and in the half of the space for the perch group that was furthest away from the  
161 predator.

162

### 163 *Behavioural experiments*

164 Both sets of behavioural experiments (before and after the predator tank treatment) consisted  
165 each of three repeated observations. Prior to each set of three behavioural observations the  
166 small perch were acclimatized to the aquarium for 3 days and fed daily with red chironomid  
167 larvae in the open area. The fish were then observed three times, twice on the first day with a  
168 break of three hours between experiments and once on the second day. Before each  
169 observation the juvenile perch were enclosed by the opaque screen in the half of their section  
170 that also contained the vegetation. Chironomid larvae (approx. 75 larvae, corresponding to 3%  
171 of the total fish weight) were poured into the open space produced between the net and the  
172 opaque screen and allowed to sink to the bottom. The observation started by lifting the opaque  
173 screen, making the large perch visible to the juvenile perch through the net. Each aquarium  
174 was observed for 10 min. The observer recorded (in real time) four different activities for  
175 each individual fish: occurrence in the vegetation, occurrence in the open, feeding and  
176 predator inspection. Thereby, feeding was defined as being oriented towards the bottom and  
177 attacking the food and predator inspection as being within two fish lengths distance of the net  
178 and being orientated exactly towards the predator. The activities were entered into a computer  
179 program, which record one behavioural unit every second. The recorded behavioural data  
180 were used to calculate 7 behavioural variables: (1) time spent in the open area, (2) total time

181 spent feeding, (3) time to start feeding, (4) duration of the first feeding bout, (5) activity  
182 (number of changes between open area and vegetation), (6) time until first change of habitat,  
183 and (7) time spent with predator inspection. After each observation the opaque screen was put  
184 back next to the net. Each group was always tested in the same aquaria and in presence of the  
185 same predator. The predators used during the behavioural experiments were different from  
186 those used during the predator treatment in the tanks.

187

### 188 *Statistical analyses*

189 One group of four fish was excluded from the analyses, as two of its members were lost  
190 during the tank treatment (non-predator tank treatment), leaving a total of 23 groups and 92  
191 individuals that were used for the analyses.

192 All 7 calculated behavioural variables were entered into a principal component analyses  
193 (PCA). By using a PCA we are able to use all data, and gain information about the  
194 relationship of the different variables to each other. The resulting scores illustrate an  
195 individual's personality type rather than an isolated behaviour and retain the structure and  
196 variation present in the recorded data. We retained all extracted principal components with  
197 eigenvalues larger than 1 for further analyses.

198 To control for a possible ceiling effect due to the limit of the behavioural assays (10 min), we  
199 conducted a second PCA including an additional imaginary individual, which was given  
200 maximal values in each variable.

201 To analyse the effect of the predator tank treatment on the extracted principal components a  
202 linear mixed effect model approach was set-up. To avoid pseudoreplication in the analysis a  
203 nested design was created. Within individual repetition (behavioural observation 1-3) was  
204 added as random effect at the innermost level. Between-individuals within group was added  
205 as the next level, between-groups in one tank was added at the following level and between-

206 tanks was added as random effect at the outer level. Treatment (before and after the predator  
207 tank treatment), experimental block (block 1 and 2) and their interaction were included as  
208 fixed effects with two factor levels each. Hence, the models for PC1 and PC2 were fit with the  
209 predictors of predator tank treatment, experimental block and their interaction and the random  
210 intercepts of within-individual repetition, individual, group and tank ID.

211 The effect of predator presence during the predator tank treatment was analysed incorporating  
212 only the data obtained after the predator tank treatment. Keeping the previously described  
213 structure of the random effects (within individual (repetition)/between individuals/between  
214 groups/between tanks), we included predator (predator or no predator during the treatment),  
215 experimental block (block 1 and 2) and their interaction as fixed effects.

216 The equivalent mixed-effect model was run using the behavioural data from before the tank  
217 treatment, including only experimental block (block 1 and 2) as fixed effect. The most  
218 parsimonious models for the separate data from before and after the tank treatment were  
219 derived by testing the fixed effects using Wald statistics (Pinheiro & Bates 2000).

220 To obtain an individual score for the relative ranking of an individual within its group we  
221 extracted the best linear unbiased predictors (BLUPs) of the random effect of `between  
222 individuals` from the two most parsimonious models for PC1 for before and after the tank  
223 treatment (Magnhagen & Bunnefeld 2009). Using a linear model, BLUPs from before were  
224 correlated with the BLUPs from after, to analyse the relative individual behavioural  
225 consistency across the predator tank treatment (before and after). Additionally, we analysed  
226 the relative individual behavioural consistency across the predator tank treatment (before and  
227 after) using the average (observation 1-3) of the seven calculated behavioral variables and  
228 Kendall correlations, as the data was not normally distributed.

229 To analyse the magnitude of the behavioural change, the difference between the behaviour  
230 before and the behaviour after the predator tank treatment was calculated (PC1 after – PC1

231 before). The difference between the behaviour before and the behaviour after the predator  
232 tank treatment was used as response variable to run a linear mixed effect model. In this model  
233 we used the same nested hierarchy for the random effects as stated above (within individual  
234 (repetition)/between individuals/between groups/between tanks). In addition to the categorical  
235 fixed effects experimental block and predator, the mean change of the accompanying group  
236 members (company change) and the change in condition factor during the predator tank  
237 treatment (condition factor after – condition factor before) as continuous fixed effects.  
238 The free software R for statistical computing (R Development Core Team 2009) was used for  
239 all analyses. The PCA (*prcomp*) and the correlation (*lm*) were calculated with the standard  
240 *stats* library. The library *nlme* v.3.1.-90 was used to run the mixed effect models.

241

#### 242 *Ethical note*

243 No prey individual was consumed or harmed (no bite marks or injuries) during the  
244 experiments. The experiments in this study comply with the guidelines of the Association for  
245 the Study of Animal Behaviour, and were approved by the Local Ethics Committee of the  
246 Swedish National Board for Laboratory Animals (CFN, license no A94-06).

247

## 248 **RESULTS**

249 The PCA on the 7 behavioural variables produced two principle components with eigenvalues  
250  $> 1$  (PC1 and PC2), explaining together 72.5% of the variation (Figure 1). Positive scores on  
251 PC1 indicated more time in the open, more time feeding, a longer duration of the first feeding  
252 bout, a lower latency to leave the vegetation and lower latency to start feeding, which would  
253 signify a fish with a high degree of boldness. Positive scores on PC2 indicated more time  
254 spent with predator inspection and a lower duration of the first feeding bout, specifying  
255 vigilance. Including an imaginary individual with maximal values in all variables in a second

256 PCA showed that the boldest observed individual in our study showed 71.9% of the possible  
257 maximum boldness, indicating that our results are not biased by any ceiling effect.

258

### 259 *The effect of the predator treatment*

260 The presence of a predator during the tank treatment did not have any effect on PC1  
261 (boldness) and PC2 (vigilance) (Table 2), neither did experimental block, nor their interaction  
262 show a significant effect. However, after the predator treatment, perch were significantly  
263 bolder and less vigilant than before treatment (Table 3, Figure 2). For PC1 the predator  
264 treatment additionally showed a significant interaction with the experimental block (Table 3).  
265 Before the tank treatment PC1 scores were slightly lower for the first block than for the  
266 second, while these relations were absent after the predator treatment.

267

### 268 *Magnitude of behavioural change*

269 For PC1, the most parsimonious models using the separate data from before and after the  
270 predator treatment were the models without any fixed factors, which were used to extract the  
271 BLUPs. The BLUPs for individual perch before the predator treatment were correlated with  
272 the BLUPs from after the predator treatment ( $t_{90} = 2.3$ ,  $P = 0.02$ ,  $r^2 = 0.05$ ), however, the data  
273 points are scattered (Figure 3), and the low  $r^2$ -value indicates that the correlation is weak and  
274 95% of the variation remains unexplained. Similarly, out of the seven calculated behavioral  
275 variables, the behavior before and after the predator treatment was only significantly and  
276 weakly correlated for two behavioral variables (Total time spend in the Open:  $p = 0.001$ ;  
277 Kendall's tau coefficient 0.235; Time until first change:  $p < 0.001$ ; Kendall's tau coefficient  
278 0.289).

279 During the predator treatment juvenile fish slightly decreased in condition factor ( $t_{177.78} =$   
280 6.49,  $P = 0.001$ ; condition factor,  $X \pm SD$ .; before  $0.99 \pm 0.06$ , after  $0.93 \pm 0.08$ ). This change

281 in condition factor did not affect the magnitude of behavioural change (PC1 after the tank  
282 treatment – PC1 before the predator tank treatment), nor did predator presence or the  
283 experimental block (Table 4). An individual's magnitude of change was significantly  
284 influenced by the mean difference between the scores before and after predator treatment of  
285 the accompanying group members (Table 4), in a way that the individual magnitude of change  
286 increased with increasing company change (Figure 4).

287

## 288 **DISCUSSION**

289 We could show that the behavioural reaction towards a predator in groups of juvenile perch  
290 can be expressed by two distinct behavioural axes. The behavioural measures that load on the  
291 first (PC1) axis (e.g. time spent in the open or time to start feeding), comply with those used  
292 in other studies on fish to investigate differences in boldness (Snickars et al. 2004;  
293 Magnhagen & Borcharding 2008). Further, the variation in PC2 scores mainly reflects  
294 whether the fish performed predator inspection or not, generally interpreted as vigilance  
295 (Pitcher 1992).

296 Comparing the scores of the same groups of juvenile perch, before and after they participated  
297 in the three week predator tank treatment, showed that juvenile perch increased in boldness  
298 (PC1) and decrease in vigilance (PC2) across the repeated measurements. Similar results were  
299 previously interpreted as habituation effects to a novel environment, represented by the  
300 laboratory conditions (Milot et al. 2009), but were also observed in response to decreasing  
301 predation risk (Goldenberg et al. 2014). Surprisingly, we found no behavioural difference  
302 between the fish that experienced a predator during the treatment and those without predator  
303 cues, although perch generally seem to adapt their behaviour to the experienced level of  
304 predation pressure (Magnhagen & Borcharding 2008; Magnhagen et al. 2012; Goldenberg et  
305 al. 2014). Laboratory studies have shown that juvenile fish rely on predator cues to optimize

306 the trade-off between foraging and anti-predator reaction, where juveniles responded strongest  
307 to the connections of olfactory and visual cues (Mikheev et al. 2006; Martin et al. 2010).

308 When predator and prey belong to the same species the diet of the predator (Mirza & Chivers  
309 2001), but also the relative size of prey and predator may be important factors influencing the  
310 behavioural reaction (Lundvall et al. 1999). The predators in our study were fed earthworms,  
311 but the prey/predator size ratio in the tanks was below the maximal ratio for ingestibility  
312 (Claessen et al. 2000) and juveniles responded to quick movements of the equally large  
313 predators in the aquaria. However, fish are capable of learning (Braithwaite & Salvanes 2008)  
314 and to habituate to initially threatening cues (Meliska & Meliska 1976), suggesting that, after  
315 a habituation period the predator, confined to a box, might not have been considered a real  
316 danger. Those results are indicative for a general problem and highlight the difficulty in  
317 generating naturalistic, but harmless scenarios of predation risk, to study the effects of long-  
318 term predation pressure in laboratory environments.

319 Analysing the behavioural consistency, we found a significant correlation of the BLUPs from  
320 before and after the predator tank treatment, this correlation explained 5% of the variation ( $r^2$   
321 = 0.05). This indicates that the consistency of inter-individual behavioural differences within  
322 a group of perch was rather low in this study. Individuals differed in the relative increase in  
323 boldness across the predator tank treatment and, thus, varied in the shape of their reaction  
324 norms. The analysis of the individual magnitude of behavioural change showed that the  
325 individuals in our study were not influenced by changes in condition, which is a known factor  
326 to alter boldness in juvenile fish (Vehanen 2003; Borcharding & Magnhagen 2008). Instead,  
327 there was a relationship between the individual and the accompanying group members, as  
328 individual magnitude of behavioural and the magnitude of behavioural change of the  
329 accompanying group members were positively correlated. In many social species the  
330 members of a group have been found to influence each other (Krause & Ruxton 2002). The

331 behavioural conformity in a group facilitated by these social mechanisms has been suggested  
332 to further confuse an attacking predator (Zheng et al. 2005; Tosh et al. 2009). Thus, our result  
333 suggests that the individuals within a group influenced each other in the magnitude of their  
334 behavioural change, probably reflecting social constraints and increasing behavioural  
335 conformity (Laubu et al. 2016).

336 Comparing the influence of holding conditions on the outcome and comparability of  
337 behavioural experiments in birds, Miller et al. (2006) could show that the holding conditions  
338 (e.g. presented food or structure within the cage) may substantially alter the obtained results  
339 of behavioural experiments. Our results suggest that also the holding time might be a crucial  
340 factor affecting behaviour, as the between-individual variation seems to change along the  
341 temporal gradient and therewith not necessarily represents the initially present “natural”  
342 behavioural variation. These results emphasize the practical implications of the reaction norm  
343 concept and the benefits of measuring behavioural variation repeatedly across an influential  
344 gradient.

345

346 In conclusion, we found juvenile perch to increase in boldness across the repeated  
347 measurements, phenotypically adapting their behaviour to the predator tank treatment, but  
348 individuals differed in the shape of their reaction norms. The magnitude of behavioural  
349 change was influenced by group composition. However, there was no behavioural difference  
350 between the fish that experienced a predator during the tank treatment and those without  
351 predator cues, indicating the general difficulties in generating realistic long-term predatory  
352 threat in laboratory environments. Furthermore, the between-individual variation seems to  
353 change along the temporal gradient and therewith not necessarily represents the initially  
354 present “natural” variation, emphasizing the importance of repeated behavioural

355 measurements and highlighting that initial holding/ laboratory acclimatisation time needs to  
356 be chosen carefully.

357

358 *Acknowledgments*

359 We thank Martin Vestman for assistance in the field, Markus Volpers who programmed the  
360 computer software to record behaviour and Lynsey McInnes for comments on the manuscript.

361 The study was financially supported by German Research Foundation to JB (BO 1507/6-2).

362

363

## References

364

365

366 Agrawal, A.A. (2001). Ecology - phenotypic plasticity in the interactions and evolution of  
367 species. *Science*. **294**, 321-326.

368 Bagenal, T.B. & Tesch, F.W. (1978). Age and growth. In: *Methods for assessment of fish*  
369 *production in fresh waters* (Ed. by T. B. Bagenal), pp. 101-136. Oxford, London,  
370 Edinburgh, Melbourne: Blackwell Scientific Pub.

371 Bell, A.M. (2007). Future directions in behavioural syndromes research. *Proc. R. Soc. Lond*  
372 *B*. **274**, 755-761.

373 Bell, A.M. & Sih, A. (2007). Exposure to predation generates personality in threespined  
374 sticklebacks (*Gasterosteus aculeatus*). *Ecol. Lett.* **10**, 828-834.

375 Boake, C.R.B. (1989). Repeatability - its role in evolutionary studies of mating-behavior.  
376 *Evol. Ecol.* **3**, 173-182.

377 Borcharding, J. & Magnhagen, C. (2008). Food abundance affects both morphology and  
378 behaviour of juvenile perch. *Ecol. Freshw. Fish.* **17**, 207-218.

379 Braithwaite, V.A. & Salvanes, A.G.V. (2008). Cognition: learning and memory. In: *Fish*  
380 *Behaviour* (Ed. by C. Magnhagen, V.A. Braithwaite, E. Forsgren & B.G. Kapoor), pp. 33-  
381 61. Enfield, NH, USA. Science Publishers.

382 Briffa, M., Rundle, S.D. & Fryer, A. (2008). Comparing the strength of behavioural plasticity  
383 and consistency across situations: animal personalities in the hermit crab *Pagurus*  
384 *bernhardus*. *Proc. R. Soc. Lond B*. **275**, 1305-1311.

385 Claessen, D., De Roos, A.M. & Persson, L. (2000). Dwarfs and giants: cannibalism and  
386 competition in size-structured populations. *Amer. Nat.* **155**, 219-237.

387 Coleman, K. & Wilson, D.S. (1998). Shyness and boldness in pumpkinseed sunfish:  
388 individual differences are context-specific. *Anim. Behav.* **56**, 927-936.

389 Cresswell, W. & Quinn, J.L. (2005). Personality, anti-predation behaviour and behavioural  
390 plasticity in the chaffinch *Fringilla coelebs*. *Behaviour*. **142**, 1377-1402.

391 Dingemanse, N.J., Both, C., Drent, P.J. & Tinbergen, J.M. (2004). Fitness consequences of  
392 avian personalities in a fluctuating environment. *Proc. R. Soc. B*. **271**, 847-852.

393 Dingemanse, N.J., Kazem, A.J.N., Reale, D. & Wright, J. (2010). Behavioural reaction  
394 norms: animal personality meets individual plasticity. *Trends Ecol. Evol.* **25**, 81-89.

395 Dingemanse, N.J. & Reale, D. (2005). Natural selection and animal personality. *Behaviour*.  
396 **142**, 1159-1184.

397 Goldenberg, U.S., Borcharding, J. & Heynen, M. 2014. Balancing the response to predation –  
398 The effects of shoal size, predation risk and habituation on behavior of juvenile perch.  
399 *Behav Ecol Sociobiol.* **68**, 989-998

400 Hayes, J.P. & Jenkins, S.H. (1997). Individual variation in mammals. *J Mammal.* **78**, 274-  
401 293.

402 Hellström, G. & Magnhagen, C. (2011). The influence of experience on risk taking. Results  
403 from a common-garden experiment on populations of Eurasian perch. *Behav. Ecol.*  
404 *Sociobiol.* **65**, 1917–1926.

405 Heynen, M., Heermann, L. & Borcharding, J. (2010). Does the consumption of divergent  
406 resources influence risk taking behaviour in juvenile perch (*Perca fluviatilis* L.)? *Ecol.*  
407 *Freshw. Fish.* **19**, 163-169.

408 Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster,  
409 H., De Jong, I.C., Ruis, M.A.W. & Blokhuis, H.J. (1999). Coping styles in animals: current  
410 status in behavior and stress-physiology. *Neurosci. Biobehav. Rev.* **23**, 925-935.

411 Krause, J. & Ruxton, G.D. (2002). *Living in groups*. Oxford: Oxford University Press.

412 Laubu, C., Dechaume-Moncharmont, F.-X., Motreuil, S. & Schweitzer, C. (2016).  
413 Mismatched partners that achieve postpairing behavioral similarity improve their  
414 reproductive success. *Sci Adv.* **2**(3), e1501013

415 Lundvall, D., Svanbäck, R., Persson, L. & Byström, P. (1999). Size-dependent predation in  
416 piscivores: interactions between predator foraging and prey avoidance abilities. *Can. J.*  
417 *Fish Aquat. Sci.* **56**, 1285-1292.

418 Magnhagen, C. & Borcharding, J. (2008). Risk-taking behaviour in foraging perch: does  
419 predation pressure influence age-specific boldness? *Anim. Behav.* **75**, 509-517.

420 Magnhagen, C. & Bunnefeld, N. (2009). Express your personality or go along with the group:  
421 what determines the behaviour of shoaling perch? *Proc. R. Soc. Lond B.* **276**, 3369-3375.

422 Magnhagen, C. & Staffan, F. (2005). Is boldness affected by group composition in young-of-  
423 the-year perch (*Perca fluviatilis*)? *Behav. Ecol. Sociobiol.* **57**, 295-303.

424 Magnhagen, C., Hellström, G., Borcharding, J. & Heynen, M. (2012). Boldness in two perch  
425 populations – long-term differences and the effect of predation pressure. *J. Anim. Ecol.* **81**,  
426 1311

427 Martin, C.W., Fodrie, F.J., Heck, K.L. & Mattila, J. (2010). Differential habitat use and  
428 antipredator response of juvenile roach (*Rutilus rutilus*) to olfactory and visual cues from  
429 multiple predators. *Oecologia.* **162**, 893-902.

430 Martin, J.G.A. & Reale, D. (2008). Temperament, risk assessment and habituation to novelty  
431 in eastern chipmunks, *Tamias striatus*. *Anim. Behav.* **75**, 309-318.

432 Meliska, J.A. & Meliska, C.J. (1976). Effects of habituation on threat display and dominance  
433 establishment in Siamese fighting fish, *Betta Splendens*. *Anim. Learn. Behav.* **4**, 167-171.

434 Mikheev, V.N., Wanzenbock, J. & Pasternak, A.F. (2006). Effects of predator-induced visual  
435 and olfactory cues on 0+perch (*Perca fluviatilis* L.) foraging behaviour. *Ecol. Freshw.*  
436 *Fish.* **15**, 111-117.

437 Miller, K.A., Garner, J.P. & Mench, J.A. (2006). Is fearfulness a trait that can be measured  
438 with behavioural tests? A validation of four fear tests for Japanese quail. *Anim. Behav.* **71**,  
439 1323-1334.

440 Millot, S., Bégout, M.-L. & Chatain, B. (2009). Risk-taking behaviour variation over time in  
441 sea bass *Dicentrarchus labrax*: effects of day-night alternation, fish phenotypic  
442 characteristics and selection for growth. *J. Fish Biol.* **75**, 1733-1749

443 Mirza, R.S. & Chivers, D.P. (2001). Do juvenile yellow perch use diet cues to assess the level  
444 of threat posed by intraspecific predators? *Behaviour.* **138**, 1249-1258.

445 Nussey, D.H., Wilson, A.J. & Brommer, J.E. (2007). The evolutionary ecology of individual  
446 phenotypic plasticity in wild populations. *J. Evol. Biol.* **20**, 831-844.

447 Pinheiro, J. & Bates, D. (2000). *Mixed-effect models in S and S-Plus*. New York, NY:  
448 Springer-Verlag

449 Pitcher, T. (1992). Who dares wins: the function and evolution of predator inspection  
450 behaviour in shoaling fish. *Neth. J. Zool.* **42**, 371-391

451 Price, T.D., Qvarnstrom, A. & Irwin, D.E. (2003). The role of phenotypic plasticity in driving  
452 genetic evolution. *Proc. R. Soc. B.* **270**, 1433-1440.

453 R Development Core Team 2009 R: a language and environment for statistical computing.  
454 Vienna, Austria: R Foundation for Statistical Computing. (ISBN 3-900051-07-0).  
455 <http://www.R-project.org>.

456 Sih, A., Bell, A. & Johnson, J.C. (2004). Behavioral syndromes: an ecological and  
457 evolutionary overview. *Trends Ecol. Evol.* **19**, 372-378.

458 Sinn, D.L., Gosling, S.D. & Moltschaniwskyj, N.A. (2007). Development of shy/bold  
459 behaviour in squid: context-specific phenotypes associated with developmental plasticity.  
460 *Anim. Behav.* **75**, 433-442.

461 Snickars, M., Sandstrom, A. & Mattila, J. (2004). Antipredator behaviour of 0+ year *Perca*  
462 *fluviatilis*: effect of vegetation density and turbidity. *J Fish Biol.* **65**, 1604-1613.

463 Smith, B.R. & Blumstein, D.T. (2008). Fitness consequences of personality: a meta-analysis.  
464 *Behav. Ecol.* **19**, 448-455.

465 Tosh, C., Krause, J. & Ruxton, G. (2009). Basic features, conjunctive searches, and the  
466 confusion effect in predator-prey interactions. *Behav. Ecol. Sociobiol.* **63**, 473-475.

467 Vehanen, T. (2003). Adaptive flexibility in the behaviour of juvenile Atlantic salmon: short-  
468 term responses to food availability and threat from predation. *J. Fish Biol.* **63**, 1034-1045.

469 Via, S., Gomulkiewicz, R., Dejong, G., Scheiner, S.M., Schlichting, C.D. & Vantienderen,  
470 P.H. (1995). Adaptive phenotypic plasticity - consensus and controversy. *Trends Ecol.*  
471 *Evol.* **10**, 212-217.

472 Wilson, D.S., Coleman, K., Clark, A.B. & Biederman, L. 1993. Shy-bold continuum in  
473 pumpkinseed sunfish (*Lepomis gibbosus*): an ecological study of a psychological trait. *J.*  
474 *Comp. Physiol.* **107**, 250-260.

475 Zheng, M., Kashimori, Y., Hoshino, O., Fujita, K. & Kambara, T. (2005). Behavior pattern  
476 (innate action) of individuals in fish schools generating efficient collective evasion from  
477 predation. *J. Theor. Biol.* **235**, 153-167.

478 Tables

479 Table 1: Experimental design

480

	Block 1	Block 2
Day 1-5	acclimatization in tank	acclimatization in tank
Day 6-8	acclimatization to aquaria	acclimatization in tank
Day 9-10	behavioural experiments “before”	acclimatization in tank
Day 11-14	predator treatment	acclimatization to aquaria
Day 15-16	predator treatment	behavioural experiments “before”
Day 16-31	predator treatment	predator treatment
Day 32-34	acclimatization to aquaria	predator treatment
Day 35-36	behavioural experiments “after”	predator treatment
Day 38-40		acclimatization to aquaria
Day 41-42		behavioural experiments “after”

481

482 Table 2: Wald statistic for the fixed effects for PC1 and PC2 before and after the predator tank  
 483 treatment, tested separately with mixed effect models

484

	<u>PC1 - boldness</u>			<u>PC2 - vigilance</u>		
	F	df,df <sub>den</sub>	P	F	df,df <sub>den</sub>	P
<hr/> Before						
Exp block	1.42	1,4	0.29	0.74	1,4	0.43
<hr/> After						
Predator	1.18	1,2	0.39	0.24	1,2	0.67
Exp block	1.89	1,2	0.30	4.47	1,2	0.16
Predator x exp block	0.03	1,2	0.86	5.04	1,2	0.15

485

486

487 Table 3: Wald statistic for the fixed effects Treatment (before and after) experimental block  
 488 and their interaction for PC1 and PC2 tested with a mixed effect model

489

	<u>PC1 - boldness</u>			<u>PC2 - vigilance</u>		
	F	df,df <sub>den</sub>	P	F	df,df <sub>den</sub>	P
Treatment	<i>1160</i>	<i>1,270</i>	<i>0.001</i>	<i>8.87</i>	<i>1,270</i>	<i>0.003</i>
Exp block	0.37	1,4	0.57	4.85	1,4	0.09
Treatment x exp block	<i>21.69</i>	<i>1,270</i>	<i>0.001</i>	2.22	1,270	0.13

490

491

492 Table 4: Wald statistic for the fixed effects company change (mean magnitude of behavioural  
493 change of the accompanying group members across the predator tank treatment), predator  
494 (predator presence or absence during the treatment), condition factor change and experimental  
495 block on the individual magnitude of change (before after difference in boldness score PC1  
496 across the predator tank treatment)

497

	F	df,df <sub>den</sub>	P
Company change	<i>43.28</i>	<i>1,179</i>	<i>0.001</i>
Predator	0.77	1,3	0.81
Condition factor change	0.14	1,68	0.52
Exp block	3.50	1,3	0.25

498

499

500 Figure legends

501

502 Figure 1: Biplot of the first two principal components, PC1 (Eigenvalue 4.02) and PC2  
503 (Eigenvalue 1.06), extracted from a Principal Component Analysis (PCA) over the seven  
504 different measured behavioral variables [(1) time spent in the open area, (2) total time spent  
505 feeding, (3) time to start feeding, (4) duration of the first feeding bout, (5) activity (number of  
506 changes between open area and vegetation), (6) time to first change of habitat, and (7) time  
507 spent with predator inspection].

508

509 Figure 2: Mean boldness (PC1) and vigilance score (PC2) before and after the predator tank  
510 treatment, in absence or presence of a predator during the treatment


511

512 Figure 3: Correlation of the best linear unbiased predictors (BLUPs) from before the predator  
513 tank treatment with the BLUPs after the predator tank treatment, extracted on individual level  
514 from the most parsimonious mixed effect models for before and after the predator tank  
515 treatment, respectively

516

517 Figure 4: Mean individual increase in boldness (After – before difference PC1) plotted against  
518 the mean company difference (mean increase in boldness of the accompanying group  
519 members).

Without predator



With predator

