

1 **Effect of number of sprints in a SIT session on change in VO₂max: a meta-analysis**

2

3 Niels BJ Vollaard^{1*}, Richard S Metcalfe², Sean Williams¹

4

5 ¹ *Department for Health, University of Bath, Bath, UK*

6 ² *School of Sport, Ulster University, Northern Ireland, UK*

7

8 *** Corresponding Author**

9 Dr Niels BJ Vollaard

10 Department for Health

11 University of Bath

12 Bath, BA2 7AY

13 UK

14 Phone: +44 1225 384649

15 Email: n.vollaard@stir.ac.uk

16

17 This is a non-final version of an article published in final form in *Medicine & Science in Sports*

18 & *Exercise* published by American College of Sports Medicine. Available at:

19 <https://doi.org/10.1249/MSS.0000000000001204>

20

21 **ABSTRACT**

22

23 **Purpose:** Recent meta-analyses indicate that sprint interval training (SIT) improves
24 cardiorespiratory fitness ($\dot{V}O_{2\max}$), but the effects of various training parameters on the
25 magnitude of the improvement remain unknown. The present meta-analysis examined the
26 modifying effect of the number of sprint repetitions in a SIT session on improvements in
27 $\dot{V}O_{2\max}$.

28 **Methods:** The databases PubMed and Web of Science were searched for original studies
29 that have examined pre- and post-training $\dot{V}O_{2\max}$ in adults following ≥ 2 weeks of training
30 consisting of repeated (≥ 2) Wingate-type cycle sprints, published up to 1 May 2016. Articles
31 were excluded if they were not in English, involved patients, athletes, or participants with a
32 mean baseline $\dot{V}O_{2\max}$ of $>55 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ or a mean age <18 years, and if a SIT trial was
33 combined with another intervention or used intervals shorter than 10 s. A total of 38 SIT trials
34 from 34 studies were included in the meta-analysis. Probabilistic magnitude-based inferences
35 were made to interpret the outcome of the analysis.

36 **Results:** The meta-analysis revealed a likely large effect of a typical SIT intervention on
37 $\dot{V}O_{2\max}$ (mean \pm 90 CL %: $7.8\% \pm 4.0\%$) with a possibly small modifying effect of the
38 maximum number of sprint repetitions in a training session ($-1.2 \pm 0.8\%$ decrease per 2
39 additional sprint repetitions). Apart from possibly small effects of baseline $\dot{V}O_{2\max}$ and age,
40 all other modifying effects were unclear or trivial.

41 **Conclusion:** We conclude that the improvement in $\dot{V}O_{2\max}$ with SIT is not attenuated with
42 fewer sprint repetitions, and possibly even enhanced. This means that SIT protocols can be
43 made more time-efficient, which may help SIT to be developed into a viable strategy to impact
44 public health.

45

46 **Key words:** systematic review; cardiorespiratory fitness; aerobic capacity; sprint interval
47 training

48 **1 INTRODUCTION**

49

50 The global increase in prevalence of noncommunicable diseases over the past decades (34)
51 can be attributed, at least in part, to the low levels of physical activity undertaken by the
52 majority of the general population (16). In light of this, a key aim of public health organisations
53 is to increase population physical activity levels (20). Of the health markers that can be
54 improved by physical activity, maximal aerobic capacity ($\dot{V}O_2\text{max}$) is consistently shown to be
55 the strongest prognostic marker for future cardiovascular health and premature death in cross-
56 sectional studies (38, 56). Furthermore, longitudinal studies demonstrate that improvements
57 in $\dot{V}O_2\text{max}$ are associated with substantial reductions in all-cause and cardiovascular mortality
58 during follow-up (9, 43).

59 Over the past two decades, relatively high volumes of moderate-intensity aerobic exercise
60 (total time commitment ≥ 150 min per week) have consistently been recommended for
61 improving health markers (20). However, uptake of and adherence to these recommendations
62 remains low in the general population (25), with lack of time identified as one of the main
63 perceived barriers to becoming and remaining physically active (39, 41, 71). Therefore, the
64 seminal finding by Burgomaster *et al.* (12) that a training protocol consisting of repeated brief
65 'all-out' cycle sprints (i.e. Wingate sprints) is associated with aerobic adaptations, has led to
66 substantial interest in the use of (sub)maximal high-intensity interval training (HIIT) and
67 supramaximal sprint interval training (SIT) as time-efficient alternative/adjunct exercise
68 strategies for improving $\dot{V}O_2\text{max}$ (21). The most commonly studied SIT protocol consists of 4-
69 7 repeated 30-s Wingate sprints, thus resulting in less than 4 minutes of high-intensity exercise
70 per session (75). Over the past few years, several meta-analyses have reported the efficacy
71 of SIT in increasing $\dot{V}O_2\text{max}$ (24, 53, 65, 75). These have concluded that in healthy individuals,
72 SIT improves $\dot{V}O_2\text{max}$ to a similar (24) or greater extent (53) than traditional aerobic training,
73 with greater benefits for individuals with lower pre-training $\dot{V}O_2\text{max}$ (53, 75).

74 Although these findings provide strong support for the effectiveness of SIT in improving
75 $\dot{V}O_{2\max}$, surprisingly few efforts have been made to identify 'optimal' SIT protocols, e.g.
76 protocols which will either provide the greatest increase in $\dot{V}O_{2\max}$, or a set increase with the
77 lowest total training volume or time commitment. Weston *et al.* (75) reported a likely small
78 effect of increasing the intervention duration and a possibly moderate effect of increasing the
79 work-to-rest ratio, but no studies have meta-analysed or directly investigated the potential
80 effects of the number of sprint repetitions in a SIT session. This parameter is particularly
81 important as it has a large influence on the total duration of a training session, as well as the
82 level of fatigue (44) and affective responses (19) experienced by the participant, thus
83 influencing the likelihood of individuals taking up and adhering to a specific SIT intervention
84 (26). As the main aim of investigating SIT protocols is generally to identify a time-efficient
85 alternative to aerobic exercise, there is a need to identify the effect of this training parameter
86 on the associated increase in $\dot{V}O_{2\max}$. Recent evidence suggests that the positive effects of
87 SIT on $\dot{V}O_{2\max}$ can be attained with fewer sprints (22, 23, 35, 50), and therefore the aim of
88 the present study was to perform a meta-analysis to provide estimates of the modifying effect
89 of the number of sprint repetitions in SIT protocols on the increase in $\dot{V}O_{2\max}$ in untrained
90 adult participants following training.

91 **2 METHODS**

92

93 **2.1 Literature Search Criteria and Study Selection**

94 This study was undertaken in accordance with the Preferred Reporting Items for Systematic
95 Reviews and Meta-Analyses (PRISMA) statement guidelines (54). We aimed to identify all
96 studies that have examined pre- and post-training $\dot{V}O_2\text{max}$ following a period of at least 2
97 weeks of training consisting of repeated (≥ 2) 'all-out' Wingate cycle sprints or modifications
98 thereof (e.g. studies using 10-s, 15-s, or 20-s 'all-out' sprints instead of 30-s Wingate sprints).
99 For this purpose, the electronic databases PubMed and Web of Science were searched for
100 relevant available records up to 1 May 2016, using the 28 possible combinations of the
101 independent variable search terms 'Wingate', 'all-out', 'sprint', and 'interval training', and the
102 dependent variable search terms 'fitness', 'aerobic capacity', 'aerobic power', ' $\dot{V}O_2\text{max}$ ',
103 ' $\dot{V}O_2\text{peak}$ ', 'oxygen uptake', and 'oxygen consumption'. Relevant studies cited in recent meta-
104 analyses were also used (24, 53, 65, 75), as well as our own recent work (52). The following
105 articles were excluded: 1) review articles / commentaries, 2) articles not written in English, 3)
106 studies concerning patients, athletes, or participants with a mean baseline $\dot{V}O_2\text{max}$ of >55
107 $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ or a mean age <18 years, 4) animal studies, 5) study-trials in which SIT was
108 combined with another intervention; and 6) SIT studies using non-cycling exercise, intervals
109 shorter than 10 s, or intervals that were not 'all-out'. Two authors (NBJV and RSM)
110 independently conducted the literature search and data extraction, and any discrepancies
111 were resolved by consensus. The reviewers were not blinded to manuscript journals or
112 authors. After removal of duplicate records, the titles and abstracts of all identified articles
113 were screened for records that were clearly not relevant. These articles were omitted before
114 assessing the full-text versions of the remaining articles for eligibility to be included in the
115 meta-analysis. If more than one article reported data for the same experiment, duplicate data
116 for these participants were only included once. The final dataset included the results of 38
117 trials from 34 studies (**Figure 1**).

118

119 2.2 Data Extraction

120 Full papers were assessed for mean absolute pre- and post-training $\dot{V}O_2\text{max}$. Absolute
121 $\dot{V}O_2\text{max}$ ($\text{L}\cdot\text{min}^{-1}$) was used rather than relative $\dot{V}O_2\text{max}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) as this provides an
122 estimate of true changes in the ability to take up and use oxygen, independent of possible
123 concomitant changes in body mass. Relative $\dot{V}O_2\text{max}$ was used for the five studies for which
124 absolute $\dot{V}O_2\text{max}$ data were not available (8, 42, 48, 57, 68). Any data for $\dot{V}O_2\text{max}$ obtained
125 at intermediate time-points during the intervention were excluded. The corresponding authors
126 of papers without the required data were contacted by email; authors from 23 studies were
127 contacted (1, 2, 5, 6, 10-13, 22, 23, 27, 28, 31, 32, 35, 40, 49, 57, 62-64, 68, 70, 77) and we
128 received raw data from 17 studies (5, 10-13, 22, 23, 27, 28, 31, 40, 49, 57, 62-64, 77). Graph
129 digitizer software (Digitizelt, Braunschweig, Germany) was used to obtain the data from one
130 study for which absolute pre- and post-training $\dot{V}O_2\text{max}$ data were only available in a figure
131 (70). The effect of training was expressed as a percentage change-score. Percentage effects
132 of SIT on $\dot{V}O_2\text{max}$ were converted to factors ($= 1 + \text{effect} / 100$), log transformed for the
133 analysis, and then back transformed to percentages. Effects were weighted using percentage
134 standard errors derived from exact p-values, or from estimated errors of measurement as
135 recommended by Weston *et al.* (75). Under the assumption that studies with similar test
136 protocols and subject characteristics would have similar typical errors of measurement, the
137 typical errors from these studies were averaged (via the weighted mean variance) and
138 assigned to the studies that did not report an exact p value (1, 2, 6, 35, 46, 68, 70). The SE
139 was then calculated via the relationship between typical error and SE (72). Finally, data for
140 the following potential moderators were extracted for each study: participant characteristics
141 (sex, age, body mass index (BMI), baseline $\dot{V}O_2\text{max}$), training parameters (intervention
142 duration, total number of training sessions, maximal number of sprint repetitions per training
143 session, sprint duration, sprint/recovery ratio, sprint resistance), and study-type (controlled /
144 uncontrolled; dummy variable). For trials with a no-exercise control group, the effect entered

145 into the meta-analysis was intervention minus control. Data for aerobic exercise comparator
146 groups were not included in the meta-analysis.

147

148 **2.3 Statistical Analysis**

149 To evaluate the extent of publication bias, a funnel plot of model residuals versus their
150 corresponding standard errors was inspected for evidence of asymmetrical scatter (75). This
151 approach takes into account any heterogeneity explained by the meta-regression, which is not
152 accounted for in standard funnel plots of observed effects vs. their standard errors. No
153 evidence of asymmetrical scatter was apparent (**Figure 2**).

154 A mixed effects meta-regression model was conducted using the 'metafor' package in R
155 (version 3.2.4, R Foundation for Statistical Computing, Vienna, Austria) (73). The overall effect
156 of SIT on $\dot{V}O_2\text{max}$ was evaluated using the mean values of the covariates. The modifying
157 effects of covariates were evaluated as the difference between levels (e.g. male/female) for
158 nominal variables. For numeric variables, effects were evaluated as the change in $\dot{V}O_2\text{max}$
159 associated with a two standard deviation (SD) change in the predictor (i.e. a typically low vs.
160 a typically high value (33)), or a practically relevant value (e.g. three additional SIT sessions
161 would typically constitute an additional week of training). The random effects in the model
162 specified a between-study SD, representing the typical difference in the true value of the effect
163 in different study settings, plus a within-study random effect to account for within-study
164 repeated measurements (a control treatment and/or more than one training treatment) (75).
165 The SD was doubled before interpreting its magnitude with the scale used to interpret fixed
166 effects (66), for the same reason that the magnitude of the effect of a linear covariate is
167 evaluated with two SD of the covariate (33). We performed a sensitivity analysis to determine
168 whether the inference relating to the modifying effect of maximum number of sprints was
169 substantially altered when two potentially influential studies (with 12 and 15 maximum sprints,
170 respectively (32, 64)) were removed from the analysis.

171 We used magnitude-based inferences to provide an interpretation of the real-world relevance
172 of the outcomes. Uncertainty in effect estimates was expressed as $\pm 90\%$ confidence limits,
173 and as the likelihood that the true value was beneficial, trivial, or harmful in relation to threshold
174 values for benefit (improved fitness) and harm (reduced fitness) (33). The overall effect of SIT
175 on $\dot{V}O_2\text{max}$ was interpreted as a clinical outcome, whereby an effect was deemed unclear if
176 the chance that the true value was beneficial was $>25\%$, with odds of benefit relative to odds
177 of harm (odds ratio) of <66 . Modifying effects were evaluated mechanistically and deemed
178 unclear if the likelihood that the true value was beneficial *and* harmful were both $>5\%$.
179 Otherwise, the effect was deemed clear, and was qualified with a probabilistic term using the
180 following scale: $<0.5\%$, most unlikely; $0.5\text{-}5\%$, very unlikely; $5\text{-}25\%$, unlikely; $25\text{-}75\%$,
181 possible; $75\text{-}95\%$, likely; $95\text{-}99.5\%$, very likely; $>99.5\%$, most likely. As robust anchors for the
182 smallest worthwhile clinical and practical effects relating to $\dot{V}O_2\text{max}$ were not available,
183 standardised effect thresholds of 0.2, 0.6 and 1.2 SD were adopted for small, moderate and
184 large effects, respectively (75). Here, the SD related to the average between-subject variances
185 for baseline $\dot{V}O_2\text{max}$; these corresponded to magnitude thresholds of 1.0%, 2.9% and 5.8%.

186 **3 RESULTS**

187

188 Data available for the 34 studies and 38 trials included in the meta-analysis are shown in **Table**
189 **1** and **Figure 3**. The meta-analysis indicated an overall likely large effect of an 'average' SIT
190 protocol on $\dot{V}O_2\text{max}$ (mean \pm 90 CL % effect on the increase in $\dot{V}O_2\text{max}$: $7.8 \pm 4.0\%$; **Table**
191 **2**). A possibly small effect was evident for the modifying effect of the maximum number of
192 sprint repetitions in a training session ($-1.2 \pm 0.8\%$ decrease per 2 additional sprint repetitions;
193 **Figure 4a**). The percentage chances that the modifying effect was negative, trivial or positive
194 were calculated to be 62.7%, 37.3% and 0.0% respectively. There were possibly small effects
195 of baseline $\dot{V}O_2\text{max}$ ($-1.5 \pm 1.9\%$ decrease per $10 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ higher baseline $\dot{V}O_2\text{max}$;
196 **Figure 4b**) and age ($-1.1 \pm 1.2\%$ decrease per 7 y increase; **Figure 4c**). All other modifying
197 effects (intervention duration, number of sessions, sprint duration, recovery time, sprint
198 resistance, BMI, sex, and study type) were unclear or trivial (**Table 2**). Unexplained variance
199 between studies was $2.2 \pm 0.8\%$ (likely moderate). The inference relating to the effect of
200 maximum number of sprint repetitions was not altered when the two studies with the highest
201 number of sprint repetitions (32, 64) were removed from the analysis ($-1.0 \pm 1.1\%$; possibly
202 small decrease; chances that the modifying effect was negative, trivial or positive of 51.6%,
203 48.2% and 0.0% respectively).

204 4 DISCUSSION

205

206 The main aim of the present meta-analysis was to examine the modifying effect of the number
207 of sprint repetitions in a SIT session on the increase in $\dot{V}O_2\text{max}$ following training. Using data
208 from 34 training studies and 418 participants we demonstrate that the improvement in $\dot{V}O_2\text{max}$
209 with SIT is not attenuated with fewer sprint repetitions, and possibly even enhanced.
210 Considering the low physical activity levels in the general population (25), and the fact that
211 lack of time is consistently identified as one of the main perceived barriers to becoming and
212 remaining physically active (39, 41, 71), this finding has implications for the design of practical
213 SIT interventions for improving general health. SIT protocols have the potential to be the most
214 time-efficient interventions that are associated with improvements in key health markers, but
215 due to the need for recovery intervals following sprints, this potential can only truly be achieved
216 if the number of sprint repetitions is low. Therefore, our observation that reducing the number
217 of sprint repetitions will not attenuate the increase in $\dot{V}O_2\text{max}$ associated with SIT, and in fact
218 may possibly improve the effect, is an important novel finding.

219 Based predominantly on the results of studies investigating the dose-response relationship
220 between regular *aerobic* exercise and improvements in health markers, it has generally been
221 accepted that at a given exercise intensity a greater volume of exercise training (in terms of
222 training duration and frequency) is associated with greater improvements in $\dot{V}O_2\text{max}$ (20). For
223 example, in a clinical trial comparing low or high-intensity aerobic training protocols with
224 matched energy expenditure (Studies of a Targeted Risk Reduction Intervention through
225 Defined Exercise (STRRIDE I)) the magnitude of change in $\dot{V}O_2\text{max}$ was greater in the group
226 exercising at a higher intensity (15). Although the volume of exercise used in HIIT and SIT
227 protocols is generally reduced compared to aerobic exercise programmes (11, 47, 63), the
228 principle of a dose-response relationship has not been challenged in these studies directly; it
229 is the interaction between training volume and intensity that is used to justify the lower volume.
230 Thus, HIIT and SIT studies investigating the effects of protocols with a lower intensity or a

231 shorter sprint duration tend to increase the number of sprint repetitions (45, 69). Apart from
232 two studies that demonstrated that reducing sprint duration from 30 s to either 15 s (77) or 10
233 s (31) does not attenuate the improvement in $\dot{V}O_{2\max}$ with SIT, there have been no HIIT or
234 SIT studies that have specifically investigated the dose-response relationship between the
235 volume of high-intensity exercise and health outcomes. Our meta-analysis provides the first
236 evidence that at 'all-out' supramaximal exercise intensities the generally accepted positive
237 association between volume of training and magnitude of training adaptations does not hold
238 true. Thus, research into the health benefits of SIT should increase the focus on protocols with
239 fewer sprints.

240 Due to the relatively low number of studies examining the effects of SIT protocols with fewer
241 than six sprint repetitions, the present meta-analysis was not powerful enough to make
242 conclusions on the optimal number of all-out sprint repetitions. Only two studies have
243 investigated the effects of a SIT protocol incorporating just two sprints (50, 52). As one of
244 these used the largest sample size of all the studies included in the review ($n=34$ (52)), the
245 mean 10% increase in $\dot{V}O_{2\max}$ observed with this protocol (termed reduced-exertion high-
246 intensity interval training, REHIT) appears to be robust. The greatest improvement in absolute
247 $\dot{V}O_{2\max}$ (17%) was reported by Gibala's group (22), who modified the original REHIT protocol
248 to include a third sprint. However, the total duration of this intervention was 12 weeks, whereas
249 at an intermediate measurement-point after 6 weeks the increase in $\dot{V}O_{2\max}$ was 12%, very
250 similar to the 10% and 14% improvements observed with the original REHIT protocol (50, 52).
251 Although future studies should determine whether the magnitude of the response for $\dot{V}O_{2\max}$
252 is different between SIT protocols incorporating 2-4 sprints, the data presented in the present
253 manuscript suggest that this difference will be small. If this is indeed the case, then a number
254 of considerations support the use of the smallest number of sprints, i.e. the two sprints used
255 in the REHIT protocol. Firstly, including a warm-up, recovery, and cool-down, this protocol has
256 the potential to be the most time-efficient protocol. Furthermore, a drawback of the use of SIT
257 as a public health intervention is the potential for high associated perceived exertion and

258 negative affective responses (8, 21). In this light it is important to point out that the number of
259 sprint repetitions has been shown to negatively affect both of these parameters (19, 44).
260 Therefore, effective SIT protocols with fewer sprint repetitions will likely offer the best chance
261 of sedentary target populations taking up and adhering to a SIT intervention for improving
262 health (18). With this in mind, the available evidence suggests that two sprints can be
263 recommended as effective at improving the important health marker of $\dot{V}O_{2max}$. It could be
264 argued that considering the apparent linear association between the number of sprint
265 repetitions and improvement in $\dot{V}O_{2max}$ (**Figure 4a**), a single sprint could be expected to
266 produce similar improvements with a lower time-commitment. However, we have recently
267 performed the first study to investigate the effects of a single supramaximal sprint on $\dot{V}O_{2max}$,
268 and observed no significant increase compared to a no-exercise control condition in response
269 to 4 weeks of training with a sample size of $n=15$ (67). Further studies are required to confirm
270 whether supramaximal sprints only improve $\dot{V}O_{2max}$ if they are repeated. Furthermore, in light
271 of the fact that the majority of studies that have studied the effects of SIT protocols
272 incorporating 2 or 3 sprint repetitions have used 20-s sprints rather than the more commonly
273 used 30-s sprints (22, 23, 50, 52), further studies are required to investigate the shortest sprint
274 duration that can be used without attenuating the adaptations to $\dot{V}O_{2max}$.

275 Our present analysis does not provide an explanation for the possibly negative effect of
276 reducing the maximal number of sprint repetitions on improvements in $\dot{V}O_{2max}$, but a
277 discussion of possible mechanisms is warranted. The main limiting factor of $\dot{V}O_{2max}$ is
278 generally assumed to be maximal cardiac output, possibly through increased blood volume
279 (7, 55). To date no studies have examined the effect of SIT on blood volume, but there is
280 evidence in favour (17, 74) and against (36) increases in blood volume in response to HIIT.
281 Similarly, there is evidence in favour (3) and against (47) increased maximal cardiac output
282 with SIT, with the latter finding suggesting that the adaptations to SIT for $\dot{V}O_{2max}$ may be
283 peripheral in origin. Indeed, several authors have proposed that improvements in $\dot{V}O_{2max}$ with
284 SIT are caused by improved skeletal muscle oxygen extraction due to increased mitochondrial

285 density (22, 36, 57, 65, 77). Although it remains unclear whether the improvement in $\dot{V}O_{2\max}$
286 with SIT is due to central or peripheral adaptations, we propose that both increased blood
287 volume and increased mitochondrial density could plausibly be explained by the rapid
288 glycogen depletion associated with supramaximal exercise (51). Firstly, maximal rates of
289 glycogenolysis in the initial 15 seconds of a supramaximal sprint (58) are associated with the
290 accumulation of metabolic derivatives, resulting in a hypertonic intramyocellular environment,
291 influx of water, and a transient ~15-20% drop in plasma volume within a timespan of just a few
292 minutes (51). This severe disturbance of circulatory homeostasis could be a stimulus for the
293 body to increase blood volume in response to repeated SIT sessions. Secondly,
294 glycogenolysis is associated with the release and activation of glycogen-bound 5' AMP-
295 activated protein kinase (AMPK) (59), which through downstream signalling pathways
296 involving peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC1 α , a
297 proposed master regulator of aerobic adaptations), could be a mechanism leading to
298 increased mitochondrial density (30). Glycogen breakdown during repeated supramaximal
299 sprints has been shown to be completely attenuated by the time of the third sprint (58), and it
300 is therefore plausible, for both of these speculated mechanisms, that performing just two
301 repeated supramaximal sprints is sufficient to 'saturate' (i.e. maximally activate) the adaptive
302 response. In other words, if either increased blood volume or mitochondrial density underpins
303 the changes in $\dot{V}O_{2\max}$ with SIT, and if rapid glycogen breakdown regulates those
304 adaptations, then no additional improvements would be expected if more than 2-3 sprints are
305 performed in a training session.

306 Apart from this hypothesis it is also possible that increasing the number of sprint repetitions
307 will result in 'pacing' strategies that affect the 'all-out' nature of the sprints (e.g. reduction of
308 peak and mean power output in initial sprints), or that accumulated fatigue may reduce the
309 effectiveness of later sprints. Furthermore, the fact that increasing the number of sprint
310 repetitions does not enhance the improvement in $\dot{V}O_{2\max}$ with SIT provides strong evidence
311 against a role for the magnitude of the acute effects of supramaximal sprints on oxygen

312 transfer, energy turnover, or total energy use, as part of the stimulus for improving $\dot{V}O_2\text{max}$
313 with SIT, because for each of these factors the stimulus should be greater with more sprint
314 repetitions.

315 A number of limitations to the present meta-analysis should be noted. Firstly, in order to be of
316 use as a practical intervention for preventing and/or treating inactivity-related chronic disease,
317 SIT interventions should also be effective at improving for example insulin sensitivity and
318 glycaemic control, blood pressure, blood lipid profile, and body composition. Therefore, one
319 limitation is that only $\dot{V}O_2\text{max}$ was included as an outcome measure in the present analysis.
320 Whereas insufficient data for a meta-analysis is available for the effects of SIT on blood
321 pressure (14, 23, 76), blood lipid profile (4, 76), and body composition (69, 76), the effect of
322 SIT on insulin sensitivity and glycaemic control has received more attention (4, 22, 23, 50, 52,
323 60, 76). However, the methods used to assess the effects of SIT on these parameters have
324 varied, with different studies using oral glucose tolerance tests (4, 50, 52, 76), intravenous
325 glucose tolerance tests (22), euglycemic hyperinsulinemic clamps (60), or continuous glucose
326 monitoring (23). This means that a meta-analysis of the effects of the number of sprint
327 repetitions in a SIT protocol on insulin sensitivity and glycaemic control is also currently not
328 feasible. Nonetheless, the improvements in insulin sensitivity and glycaemic control observed
329 to date with SIT protocols incorporating two (50) or three sprints (22, 23) are encouraging.

330 Secondly, due to the number of available SIT studies the power of our meta-analysis is
331 insufficient to conclude with certainty that the modifying effect of the number of sprint
332 repetitions is negative; i.e. it remains possible that in reality performing more sprints will result
333 in the same improvements in $\dot{V}O_2\text{max}$ (a chance of approximately 1 in 3). However, this is not
334 of major importance to the significance of our findings: even 'no effect' of the number of sprint
335 repetitions would lead to the logical conclusion that performing SIT protocols with more than
336 2 or 3 sprints is unnecessary for improving $\dot{V}O_2\text{max}$ in sedentary individuals. Based on the
337 present analysis, the chance that in reality the effect of performing more sprints is positive was
338 calculated as 0.0%.

339 A final limitation of our meta-analysis is that only SIT interventions using all-out intensities
340 were included. Optimising time-efficient interventions aimed at improving general health
341 requires consideration of various parameters, and exercise intensity is undoubtedly one of the
342 key parameters affecting the effectiveness of HIIT and SIT protocols. However, due to the
343 large range of intensities used in SIT and HIIT protocols (~80%-350% of $\dot{V}O_2\text{max}$) we felt it
344 was important to attempt to 'control' for this variable in the present analysis by including only
345 studies that used 'all-out' cycling exercise. Nonetheless, there is a clear need for studies
346 examining the effect of the number of sprint repetitions at lower exercise intensities, e.g. in
347 HIIT studies.

348 In conclusion, in the present meta-analysis we demonstrate that SIT is possibly more effective
349 at improving $\dot{V}O_2\text{max}$ if fewer sprint repetitions are performed in a training session.
350 Considering the proclaimed aim of SIT to provide a time-efficient alternative / adjunct to high-
351 volume moderate-intensity aerobic exercise, this finding has important implications for the
352 design of practical SIT interventions. We put forward that SIT research should move away
353 from further characterising the commonly used 4-7 x 30-s Wingate protocol, and towards
354 establishing acceptable and effective protocols. This will require more studies to examine the
355 modifying effects of a range of training parameters (including number of sprint repetitions,
356 sprint duration, sprint intensity, and training frequency) on adaptations to key health markers,
357 as well as exercise enjoyment and acceptability, perceived exertion, and the potential to cause
358 negative affective responses.

359 **Acknowledgements**

360 We would like to thank Professor Will Hopkins for helpful advice during the preparation of the
361 manuscript. No sources of funding were used to assist in the preparation of this review. The
362 authors have no conflicts of interest to declare. The results of the study are presented clearly,
363 honestly, and without fabrication, falsification, or inappropriate data manipulation. The results
364 of the present study do not constitute endorsement by ACSM.

References

1. Allemeier CA, Fry AC, Johnson P, Hikida RS, Hagerman FC, Staron RS. Effects of sprint cycle training on human skeletal muscle. *Journal of applied physiology*. 1994;77(5):2385-90.
2. Astorino TA, Allen RP, Roberson DW et al. Adaptations to high-intensity training are independent of gender. *European journal of applied physiology*. 2011;111(7):1279-86.
3. Astorino TA, Edmunds RM, Clark A et al. High-Intensity Interval Training Increases Cardiac Output and VO₂max. *Medicine and science in sports and exercise*. 2016.
4. Babraj JA, Vollaard NB, Keast C, Guppy FM, Cottrell G, Timmons JA. Extremely short duration high intensity interval training substantially improves insulin action in young healthy males. *BMC endocrine disorders*. 2009;9:3.
5. Bailey SJ, Wilkerson DP, Dimenna FJ, Jones AM. Influence of repeated sprint training on pulmonary O₂ uptake and muscle deoxygenation kinetics in humans. *Journal of applied physiology*. 2009;106(6):1875-87.
6. Barnett C, Carey M, Proietto J, Cerin E, Febbraio MA, Jenkins D. Muscle metabolism during sprint exercise in man: influence of sprint training. *Journal of science and medicine in sport / Sports Medicine Australia*. 2004;7(3):314-22.
7. Bassett DR, Jr., Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Medicine and science in sports and exercise*. 2000;32(1):70-84.
8. Bayati M, Farzad B, Gharakhanlou R, Agha-Alinejad H. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble 'all-out' sprint interval training. *Journal of sports science & medicine*. 2011;10(3):571-6.
9. Blair SN, Kohl HW, 3rd, Barlow CE, Paffenbarger RS, Jr., Gibbons LW, Macera CA. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *Jama*. 1995;273(14):1093-8.
10. Burgomaster KA, Heigenhauser GJ, Gibala MJ. Effect of short-term sprint interval training on human skeletal muscle carbohydrate metabolism during exercise and time-trial performance. *Journal of applied physiology*. 2006;100(6):2041-7.
11. Burgomaster KA, Howarth KR, Phillips SM et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *The Journal of physiology*. 2008;586(1):151-60.
12. Burgomaster KA, Hughes SC, Heigenhauser GJ, Bradwell SN, Gibala MJ. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *Journal of applied physiology*. 2005;98(6):1985-90.
13. Cochran AJ, Percival ME, Tricarico S et al. Intermittent and continuous high-intensity exercise training induce similar acute but different chronic muscle adaptations. *Experimental physiology*. 2014;99(5):782-91.
14. Cocks M, Shaw CS, Shepherd SO et al. Sprint interval and endurance training are equally effective in increasing muscle microvascular density and eNOS content in sedentary males. *The Journal of physiology*. 2013;591(3):641-56.
15. Duscha BD, Slentz CA, Johnson JL et al. Effects of exercise training amount and intensity on peak oxygen consumption in middle-age men and women at risk for cardiovascular disease. *Chest*. 2005;128(4):2788-93.
16. Ekelund U, Ward HA, Norat T et al. Physical activity and all-cause mortality across levels of overall and abdominal adiposity in European men and women: the European Prospective Investigation into Cancer and Nutrition Study (EPIC). *The American journal of clinical nutrition*. 2015;101(3):613-21.
17. Esfandiari S, Sasson Z, Goodman JM. Short-term high-intensity interval and continuous moderate-intensity training improve maximal aerobic power and diastolic filling during exercise. *European journal of applied physiology*. 2014;114(2):331-43.
18. Foster C, Farland CV, Guidotti F et al. The Effects of High Intensity Interval Training vs Steady State Training on Aerobic and Anaerobic Capacity. *Journal of sports science & medicine*. 2015;14(4):747-55.

19. Frazao DT, de Farias Junior LF, Dantas TC et al. Feeling of Pleasure to High-Intensity Interval Exercise Is Dependent of the Number of Work Bouts and Physical Activity Status. *PLoS one*. 2016;11(3):e0152752.
20. Garber CE, Blissmer B, Deschenes MR et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Medicine and science in sports and exercise*. 2011;43(7):1334-59.
21. Gillen JB, Gibala MJ. Is high-intensity interval training a time-efficient exercise strategy to improve health and fitness? *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2014;39(3):409-12.
22. Gillen JB, Martin BJ, MacInnis MJ, Skelly LE, Tarnopolsky MA, Gibala MJ. Twelve Weeks of Sprint Interval Training Improves Indices of Cardiometabolic Health Similar to Traditional Endurance Training despite a Five-Fold Lower Exercise Volume and Time Commitment. *PLoS one*. 2016;11(4):e0154075.
23. Gillen JB, Percival ME, Skelly LE et al. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PLoS one*. 2014;9(11):e111489.
24. Gist NH, Fedewa MV, Dishman RK, Cureton KJ. Sprint interval training effects on aerobic capacity: a systematic review and meta-analysis. *Sports medicine*. 2014;44(2):269-79.
25. Hallal PC, Andersen LB, Bull FC et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet*. 2012;380(9838):247-57.
26. Hardcastle SJ, Ray H, Beale L, Hagger MS. Why sprint interval training is inappropriate for a largely sedentary population. *Frontiers in psychology*. 2014;5:1505.
27. Harmer AR, McKenna MJ, Sutton JR et al. Skeletal muscle metabolic and ionic adaptations during intense exercise following sprint training in humans. *Journal of applied physiology*. 2000;89(5):1793-803.
28. Harmer AR, Ruell PA, McKenna MJ et al. Effects of sprint training on extrarenal potassium regulation with intense exercise in Type 1 diabetes. *Journal of applied physiology*. 2006;100(1):26-34.
29. Harris E, Rakobowchuk M, Birch KM. Sprint interval and sprint continuous training increases circulating CD34+ cells and cardio-respiratory fitness in young healthy women. *PLoS one*. 2014;9(9):e108720.
30. Hawley JA. Molecular responses to strength and endurance training: are they incompatible? *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2009;34(3):355-61.
31. Hazell TJ, Macpherson RE, Gravelle BM, Lemon PW. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *European journal of applied physiology*. 2010;110(1):153-60.
32. Hellsten-Westing Y, Balsom PD, Norman B, Sjodin B. The effect of high-intensity training on purine metabolism in man. *Acta physiologica Scandinavica*. 1993;149(4):405-12.
33. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Medicine and science in sports and exercise*. 2009;41(1):3-13.
34. Hunter DJ, Reddy KS. Noncommunicable diseases. *The New England journal of medicine*. 2013;369(14):1336-43.
35. Ijichi T, Hasegawa Y, Morishima T, Kurihara T, Hamaoka T, Goto K. Effect of sprint training: training once daily versus twice every second day. *European journal of sport science*. 2015;15(2):143-50.
36. Jacobs RA, Fluck D, Bonne TC et al. Improvements in exercise performance with high-intensity interval training coincide with an increase in skeletal muscle mitochondrial content and function. *Journal of applied physiology*. 2013;115(6):785-93.
37. Katz A, Sharp RL, King DS, Costill DL, Fink WJ. Effect of high intensity interval training on 2,3-diphosphoglycerate at rest and after maximal exercise. *European journal of applied physiology and occupational physiology*. 1984;52(3):331-5.
38. Keteyian SJ, Brawner CA, Savage PD et al. Peak aerobic capacity predicts prognosis in patients with coronary heart disease. *American heart journal*. 2008;156(2):292-300.

39. Kimm SY, Glynn NW, McMahon RP, Voorhees CC, Striegel-Moore RH, Daniels SR. Self-perceived barriers to activity participation among sedentary adolescent girls. *Medicine and science in sports and exercise*. 2006;38(3):534-40.
40. Kiviniemi AM, Tulppo MP, Eskelinen JJ et al. Autonomic Function Predicts Fitness Response to Short-Term High-Intensity Interval Training. *International journal of sports medicine*. 2015;36(11):915-21.
41. Korhakangas EE, Alahuhta MA, Laitinen JH. Barriers to regular exercise among adults at high risk or diagnosed with type 2 diabetes: a systematic review. *Health promotion international*. 2009;24(4):416-27.
42. Larsen RG, Befroy DE, Kent-Braun JA. High-intensity interval training increases in vivo oxidative capacity with no effect on P(i)-->ATP rate in resting human muscle. *American journal of physiology. Regulatory, integrative and comparative physiology*. 2013;304(5):R333-42.
43. Lee DC, Sui X, Artero EG et al. Long-term effects of changes in cardiorespiratory fitness and body mass index on all-cause and cardiovascular disease mortality in men: the Aerobics Center Longitudinal Study. *Circulation*. 2011;124(23):2483-90.
44. Little JP, Gillen JB, Percival ME et al. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *Journal of applied physiology*. 2011;111(6):1554-60.
45. Little JP, Safdar A, Wilkin GP, Tarnopolsky MA, Gibala MJ. A practical model of low-volume high-intensity interval training induces mitochondrial biogenesis in human skeletal muscle: potential mechanisms. *The Journal of physiology*. 2010;588(Pt 6):1011-22.
46. MacDougall JD, Hicks AL, MacDonald JR, McKelvie RS, Green HJ, Smith KM. Muscle performance and enzymatic adaptations to sprint interval training. *Journal of applied physiology*. 1998;84(6):2138-42.
47. Macpherson RE, Hazell TJ, Olver TD, Paterson DH, Lemon PW. Run sprint interval training improves aerobic performance but not maximal cardiac output. *Medicine and science in sports and exercise*. 2011;43(1):115-22.
48. McGarr GW, Hartley GL, Cheung SS. Neither short-term sprint nor endurance training enhances thermal response to exercise in a hot environment. *Journal of occupational and environmental hygiene*. 2014;11(1):47-53.
49. McKenna MJ, Heigenhauser GJ, McKelvie RS, Obminski G, MacDougall JD, Jones NL. Enhanced pulmonary and active skeletal muscle gas exchange during intense exercise after sprint training in men. *The Journal of physiology*. 1997;501 (Pt 3):703-16.
50. Metcalfe RS, Babraj JA, Fawkner SG, Vollaard NB. Towards the minimal amount of exercise for improving metabolic health: beneficial effects of reduced-exertion high-intensity interval training. *European journal of applied physiology*. 2012;112(7):2767-75.
51. Metcalfe RS, Koumanov F, Ruffino JS et al. Physiological and molecular responses to an acute bout of reduced-exertion high-intensity interval training (REHIT). *European journal of applied physiology*. 2015;115(11):2321-34.
52. Metcalfe RS, Tardif N, Thompson D, Vollaard NBJ. Health benefits of reduced-exertion high-intensity interval training (REHIT) are not different between sedentary men and women. *Applied Physiology, Nutrition, and Metabolism*. 2016.
53. Milanovic Z, Sporis G, Weston M. Effectiveness of High-Intensity Interval Training (HIT) and Continuous Endurance Training for VO2max Improvements: A Systematic Review and Meta-Analysis of Controlled Trials. *Sports medicine*. 2015;45(10):1469-81.
54. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Bmj*. 2009;339:b2535.
55. Montero D, Diaz-Canestro C, Lundby C. Endurance Training and V O2max: Role of Maximal Cardiac Output and Oxygen Extraction. *Medicine and science in sports and exercise*. 2015;47(10):2024-33.
56. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *The New England journal of medicine*. 2002;346(11):793-801.
57. Nalcakan GR. The Effects of Sprint Interval vs. Continuous Endurance Training on Physiological And Metabolic Adaptations in Young Healthy Adults. *Journal of human kinetics*. 2014;44:97-109.

58. Parolin ML, Chesley A, Matsos MP, Spriet LL, Jones NL, Heigenhauser GJ. Regulation of skeletal muscle glycogen phosphorylase and PDH during maximal intermittent exercise. *The American journal of physiology*. 1999;277(5 Pt 1):E890-900.
59. Philp A, Hargreaves M, Baar K. More than a store: regulatory roles for glycogen in skeletal muscle adaptation to exercise. *Am J Physiol Endocrinol Metab*. 2012;302(11):E1343-51.
60. Richards JC, Johnson TK, Kuzma JN et al. Short-term sprint interval training increases insulin sensitivity in healthy adults but does not affect the thermogenic response to beta-adrenergic stimulation. *The Journal of physiology*. 2010;588(Pt 15):2961-72.
61. Richardson AJ, Gibson OR. Simulated hypoxia does not further improve aerobic capacity during sprint interval training. *The Journal of sports medicine and physical fitness*. 2015;55(10):1099-106.
62. Scalzo RL, Peltonen GL, Binns SE et al. Greater muscle protein synthesis and mitochondrial biogenesis in males compared with females during sprint interval training. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2014;28(6):2705-14.
63. Shepherd SO, Cocks M, Tipton KD et al. Sprint interval and traditional endurance training increase net intramuscular triglyceride breakdown and expression of perilipin 2 and 5. *The Journal of physiology*. 2013;591(3):657-75.
64. Skleryk JR, Karagounis LG, Hawley JA, Sharman MJ, Laursen PB, Watson G. Two weeks of reduced-volume sprint interval or traditional exercise training does not improve metabolic functioning in sedentary obese men. *Diabetes, obesity & metabolism*. 2013;15(12):1146-53.
65. Sloth M, Sloth D, Overgaard K, Dalgas U. Effects of sprint interval training on VO₂max and aerobic exercise performance: A systematic review and meta-analysis. *Scandinavian journal of medicine & science in sports*. 2013;23(6):e341-52.
66. Smith TB, Hopkins WG. Variability and predictability of finals times of elite rowers. *Medicine and science in sports and exercise*. 2011;43(11):2155-60.
67. Songsorn P, Lambeth-Mansell A, Mair JL et al. Exercise training comprising of single 20-s cycle sprints does not provide a sufficient stimulus for improving maximal aerobic capacity in sedentary individuals. *European journal of applied physiology*. 2016;Epub: 6 June 2016.
68. Stathis CG, Febbraio MA, Carey MF, Snow RJ. Influence of sprint training on human skeletal muscle purine nucleotide metabolism. *Journal of applied physiology*. 1994;76(4):1802-9.
69. Trapp EG, Chisholm DJ, Freund J, Boutcher SH. The effects of high-intensity intermittent exercise training on fat loss and fasting insulin levels of young women. *International journal of obesity*. 2008;32(4):684-91.
70. Trilk JL, Singhal A, Bigelman KA, Cureton KJ. Effect of sprint interval training on circulatory function during exercise in sedentary, overweight/obese women. *European journal of applied physiology*. 2011;111(8):1591-7.
71. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Medicine and science in sports and exercise*. 2002;34(12):1996-2001.
72. Vandenberghe TJ, Hopkins WG. Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis. *Sports medicine*. 2011;41(9):773-92.
73. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*. 2010;36(3):1-48.
74. Warburton DE, Haykowsky MJ, Quinney HA et al. Blood volume expansion and cardiorespiratory function: effects of training modality. *Medicine and science in sports and exercise*. 2004;36(6):991-1000.
75. Weston M, Taylor KL, Batterham AM, Hopkins WG. Effects of low-volume high-intensity interval training (HIT) on fitness in adults: a meta-analysis of controlled and non-controlled trials. *Sports medicine*. 2014;44(7):1005-17.
76. Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism: clinical and experimental*. 2010;59(10):1421-8.
77. Zelt JG, Hankinson PB, Foster WS et al. Reducing the volume of sprint interval training does not diminish maximal and submaximal performance gains in healthy men. *European journal of applied physiology*. 2014;114(11):2427-36.

Figure 1: Flow diagram of the study selection process




Figure 2: Funnel plot of model residuals versus their corresponding standard errors, with 90% confidence interval region




Figure 3: Main effects of SIT on $\dot{V}O_{2\max}$




Figure 4: Modifying effects of number of sprint repetitions (A), baseline $\dot{V}O_2\text{max}$ (B), and age (C) on the effect of SIT on $\dot{V}O_2\text{max}$. Data-points represent individual trials included in the meta-analysis, and the size of the data-point is proportional to study weighting. Solid and dotted lines represent the effect of the moderator \pm 90% confidence limits respectively.




Table 1: Training effects, training protocol parameters, and participant characteristics for the studies included in the meta-analysis

Reference	Study design	SIT-group sample size (n)	Proportion of men	Mean baseline $\dot{V}O_2\text{max}$ ($\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$)	Mean age (y)	Mean BMI ($\text{kg}\cdot\text{m}^{-2}$)	Training duration (weeks)	Total training sessions	Sprint duration (s)	Recovery duration (s)	Resistance (% of BM)	Sprint repetitions		Effect on $\dot{V}O_2\text{max}$ (%)	
												Min	Max	Mean	SE
Metcalfe (50)	C	11	0.45	34.2	25.0	23.5	6	18	20	200	7.5	1	2	12.7	2.8
Metcalfe (52)	NC	34	0.50	35.0	34.1	24.6	6	18	20	200	5.0	1	2	9.6	1.5
Allemeier (1)	C	11	1.00	48.7	22.7	24.8	6	15	30	1200	7.5	3	3	13.5	2.0
Gillen (23)	NC	14	0.50	29.5	30.0	-	6	18	20	120	5.0	3	3	11.6	2.0
Ijichi (35)	C	10	1.00	47.7	20.4	21.0	4	20	30	600	5.0	3	3	13.9	1.8
Gillen (22)	C	9	1.00	32.0	27.0	27.0	12	36	20	120	5.0	3	3	17.3	3.3
Harris (29)	C	6	0.00	35.0	22.0	23.6	4	12	30	270	7.5	4	4	9.0	3.4
Bayati (8)	C	8	1.00	44.6	25.0	23.7	4	12	30	240	7.5	3	5	9.6	3.9
Barnett (6)	C	8	1.00	47.6	20.4	-	8	24	30	180	-	3	6	8.2	2.1
Burgomaster (11)	C	10	0.50	41.0	23.6	23.6	6	18	30	270	7.5	4	6	6.3	4.0
Hazell (31)	C	13	0.81	47.0	24.0	24.7	2	6	30	240	10.0	4	6	8.3	2.2
Hazell (31)	C	13	0.81	47.0	24.0	24.7	2	6	10	240	10.0	4	6	8.5	2.4
Hazell (31)	C	13	0.81	47.0	24.0	24.7	2	6	10	120	10.0	4	6	3.9	1.3
Whyte (76)	NC	10	1.00	32.8	32.1	30.3	2	6	30	270	6.5	4	6	8.4	2.6
Astorino (2)	C	20	0.55	43.6	25.0	24.1	2	6	30	300	7.5	4	6	6.3	1.5
Shepperd (63)	C	8	1.00	41.9	22.0	24.8	6	18	30	270	7.5	4	6	7.6	2.3
Larsen (42)	NC	8	1.00	25.8	27.0	26.8	2	6	30	240	7.5	4	6	9.8	3.5
Ijichi (35)	C	10	1.00	46.8	21.3	22.2	4	10	30	600	5.0	6	6	8.4	1.8
Kiviniemi (40)	C	13	1.00	34.7	48.0	25.6	2	6	30	240	7.5	4	6	4.7	1.4
McGarr (48)	C	8	0.75	47.2	25.0	-	2	8	30	240	7.5	4	6	14.2	4.5
Nalcakan (57)	C	8	1.00	40.2	21.7	25.5	7	21	30	270	7.5	4	6	7.0	1.8
Zelt (77)	C	11	1.00	48.6	23.0	25.0	4	12	30	270	7.5	4	6	5.3	2.6
Zelt (77)	C	12	1.00	43.9	22.0	26.0	4	12	15	285	7.5	4	6	7.4	2.7
Cochran (13)	C	12	1.00	50.6	22.0	25.7	6	18	30	240	7.5	4	6	10.3	2.1
Burgomaster (12)	C	8	0.75	44.6	22.0	25.6	2	6	30	240	7.5	4	7	1.4	2.0
Burgomaster (10)	C	8	1.00	48.9	21.0	23.8	2	6	30	240	7.5	4	7	5.6	2.8
Bailey (5)	C	8	0.63	42.0	21.0	23.7	2	6	30	240	7.5	3	7	7.4	2.4
Trilk (70)	C	14	0.00	21.6	30.1	35.7	4	12	30	240	5.0	4	7	11.7	1.6
Richardson (61)	C	9	0.56	40.0	21.0	23.8	2	6	30	240	7.5	4	7	11.2	2.7
Katz (37)	NC	8	1.00	51.8	24.2	-	8	32	30	240	-	8	8	7.0	2.9
Scalzo (62)	NC	21	0.52	41.5	22.5	22.4	3	9	30	240	7.5	4	8	3.7	2.1
Stathis (68)	NC	8	0.75	49.6	22.1	-	7	21	30	180	-	3	10	4.2	2.9
McKenna (49)	NC	8	1.00	47.1	20.9	23.7	7	21	30	180	7.5	4	10	13.7	3.2
MacDougall (46)	NC	12	1.00	50.8	22.7	24.0	7	21	30	180	7.5	4	10	7.5	2.4
Harmer (27)	NC	7	1.00	49.8	22.0	23.5	7	21	30	180	7.5	4	10	6.9	3.1
Harmer (28)	C	7	0.71	43.7	24.0	23.8	7	21	30	180	7.5	4	10	8.2	3.6
Skleryk (64)	C	8	1.00	29.7	40.2	32.2	2	6	10	80	5.0	8	12	-1.7	2.9
Hellsten-Westing (32)	NC	11	1.00	53.0	23.6	-	6	18	10	50	7.0	15	15	2.4	2.5

Abbreviations: BM - body mass, BMI - body mass index, C - controlled, NC - not controlled, SE - standard error, SIT - sprint interval training

Table 2 Main effect of SIT on $\dot{V}O_{2\max}$ and modifying effects

	Effect on $\dot{V}O_{2\max}$ (mean %, \pm 90% CL)	Inference
Main effect:	7.8 \pm 4.0	Likely large increase
Modifying effects:		
2 more sprint repetitions*	-1.2 \pm 0.8	Possibly small decrease
3 more training sessions*	0.7 \pm 0.4	Likely trivial change
10 s longer sprint duration*	0.6 \pm 1.3	Possibly trivial change
60 s longer recovery interval duration*	0.2 \pm 0.3	Most likely trivial change
3% of BM greater sprint resistance	1.0 \pm 2.3	Unclear
10 mL·kg ⁻¹ ·min ⁻¹ lower baseline $\dot{V}O_{2\max}$	1.5 \pm 1.9	Possibly small increase
7 years higher age	-1.1 \pm 1.2	Possibly small decrease
6.2 kg·m ⁻² higher BMI	0.8 \pm 2.7	Unclear
Female sex	-0.2 \pm 3.5	Unclear
Uncontrolled study	-0.9 \pm 2.1	Unclear

*The reference condition is an intervention using 14 SIT sessions and a maximum of 7 repeated 30-s sprints with 240 s recovery. Effects of SIT are presented as the % change compared to pre-training. *, indicates a practically relevant value was chosen to evaluate the effect magnitude; other numeric modifiers were evaluated as a 2 x SD change in the parameter. Abbreviations: BMI: body mass index, CL: confidence limits, SIT: sprint interval training, $\dot{V}O_{2\max}$: maximal aerobic capacity.*