

This is the peer reviewed version of the following article: Lazado CC, Fridman S, Sinai T, Zilberg D. First report of *Streptococcus parauberis* in a cultured freshwater ornamental fish, the ram cichlid *Mikrogeophagus ramirezi* (Myers & Harry, 1948). *Journal of Fish Diseases* 2018;41:161–164, which has been published in final form at <https://doi.org/10.1111/jfd.12676>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

1 Short communication – Revision 1 (JFD-2017-92)

2

3 **First report of *Streptococcus parauberis* in a cultured freshwater**
4 **ornamental fish, the ram cichlid *Mikrogeophagus ramirezi* (Myers &**
5 **Harry, 1948)**

6

7 Carlo C. Lazado¹, Sophie Fridman^{1,2}, Tamar Sinai¹ and Dina Zilberg^{1*}

8

9 Running title: *Streptococcus parauberis* infection in the ram cichlid

10 Accepted for publication in *Journal of Fish Diseases* published by Wiley. The published version is available at:
11 <https://doi.org/10.1111/jfd.12676>

11

12 ¹ The French Associates Institute for Agriculture and Biotechnology of Drylands, Ben-Gurion
13 University of the Negev, Midreshet Ben Gurion, Israel

14 ² Institute of Aquaculture, University of Stirling, FK9 4LA, Scotland

15

16

17 *Corresponding author

18 D. Zilberg

19 Email: dzilberg@bgu.ac.il

20 Tel: +972-8-6596818

21 Fax: +972-8-6596742

22

23

24 **Keywords:** aquaculture, fish health, histopathology, *Streptococcus parauberis*, *streptococcosis*

25 Since the first report of an outbreak of a streptococcal infection in rainbow trout
26 (*Oncorhynchus mykiss*) in Japan in 1958 (Hoshina *et al.* 1958), streptococcosis has been
27 responsible for significant mortalities resulting in considerable losses to the aquaculture
28 industry (Salati 2006; Noga 2010). Numerous species from the family Streptococcaceae have
29 been identified as etiological agents of streptococcosis in fish (Toranzo *et al.* 2005; Salati
30 2006; Noga 2010), susceptibility to which was documented in both food (Inglis *et al.* 1993)
31 and ornamental fish species (Russo *et al.* 2006). *Streptococcus parauberis* is a coccoid, non-
32 motile, alpha-hemolytic Gram-positive bacterium belonging to the *Streptococcaceae* family
33 (Nho *et al.* 2011) and has been reported as the etiological agent of streptococcosis in a few
34 fish species, including turbot (*Scophthalmus maximus*), olive flounder (*Paralichthys*
35 *olivaceus*), sea bass (*Sebastes ventricosus*) and striped bass (*Morone saxatilis*) (Domeénech *et*
36 *al.* 1996; Mata *et al.* 2004; Baeck *et al.* 2006; Park *et al.* 2009; Haines *et al.* 2013; Oguro *et al.*
37 2014). *S. parauberis* has been previously identified as the etiologic agent of bovine mastitis
38 (Bradley 2002). It was formerly known as *Streptococcus uberis* Type II until comparative
39 analysis of the sequence data of *Streptococcus uberis* Types I and II showed that both were
40 phylogenetically distinct, and the new species *Streptococcus parauberis* was proposed
41 (Williams and Collins 1990).

42 This report describes the first occurrence of septicemic disease associated with *S.*
43 *parauberis* in a cultured freshwater ornamental fish, the ram cichlid (*Mikrogeophagus*
44 *ramirezi*). This small, colorful omnivorous fish is popular among aquarists. The
45 histopathological changes associated with the infection are presented, as well as the

46 preliminary bacteriological characteristics of this first isolate of *S. parauberis* from a
47 freshwater ornamental fish.

48 Mortalities had been reported following a routine sorting procedure at a commercial
49 fish farm culturing the ram cichlid in Southern Israel in January 2014. Fish were seen to be
50 exhibiting apparent signs of sickness that included weakness, loss of equilibrium, skin redness
51 and ecchymotic hemorrhaging as well as lepidorthosis and exophthalmia (Supplementary
52 material 1). Fish were brought for examination to the Fish Health Laboratory at The Jacob
53 Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev (Midreshet Ben-
54 Gurion, Israel) monthly between January and June of 2014 and a total of around 30 fish were
55 examined. From amongst these, around 20 fish underwent a direct microscopic examination
56 of wet mounts and aseptic bacterial isolation and around 10 fish were processed for
57 histopathological analysis. For bacteriological examination, sterile swabs from the liver and
58 kidney were streaked onto tryptone soy agar (Oxoid, Hampshire, UK) and the plates were
59 incubated at 25°C for 24 h. Biochemical analyses were performed with API 20 STREP and API
60 50 CH test (API system, La Balme les Grottes, France). The isolate was sent to Hy Laboratories
61 Ltd. (Rehovot, Israel) for 16s rRNA gene sequencing and the resulting sequence was
62 subjected to comparative phylogenetic analysis. Whole fish were fixed in formalin for 48 h
63 and stored in 70% ethanol until processing by routine histological techniques.

64 Histopathological analysis revealed infiltration of macrophages, which was mostly
65 evident in liver, kidney, and muscle (Fig. 1a-e). Gram staining demonstrated the presence of
66 densely packed, Gram-positive bacteria in the infiltrating macrophages (Fig 1b, d). Focal

67 necrosis occurred in muscle fibers (Fig. 1f) and vacuolization was seen in the liver (Fig. 1c).
68 There was no evident damage to kidney tubules or stroma (Fig. 1a).

69 Gram positive cocci were isolated from symptomatic fish. On TSA plates,
70 morphological characteristics of the colony of around 1 mm in diameter included whitish-to-
71 yellowish coloration, a circular shape with a raised cross sectional elevation and a smooth
72 surface. The isolate was molecularly identified as *S. parauberis* and, from here onwards will
73 be referred to as *S. parauberis* RC. The partial 16s rRNA sequence was deposited in GenBank
74 under accession no. MF102143. Partial sequences of several *S. parauberis* isolates from
75 aquatic and terrestrial environments were retrieved from the National Center for
76 Biotechnology Information (NCBI) database to perform phylogenetic and molecular
77 evolutionary analyses in Phylogeny.fr (Dereeper *et al.* 2008). The *S. parauberis* RC was closely
78 related to other *S. parauberis* strains of aquatic origin (Fig. 2a) although it formed a separate
79 clade from the rest of the group. In the API 20 STREP test, *S. parauberis* RC was Voges-
80 Proskauer, hippuric acid and esculin positive. All other tests were negative. Furthermore, the
81 isolate was able to metabolize a number of carbohydrates including galactose, glucose,
82 fructose, mannosen-acetylglucosamine, amygdalin, arbutin, esculin ferric citrate, salicin,
83 cellobiose, maltose, lactose, saccharose, trehalose, amidon, glycogen, and gentiobiose. The
84 isolate was able to grow in a wide range of temperatures (17-33°C), though growth (OD₆₂₀)
85 was affected in temperatures lower than 21°C (Fig. 2b). It was found to thrive at various NaCl
86 concentrations (0-40 ppt) in the culture media, however, growth was negatively affected at
87 the highest salinity tested (*i.e.* 40 ppt) (Fig. 2c).

88 We evaluated the susceptibility of *S. parauberis* RC to several antibiotics including
89 *SXT*: trimethoprim/sulphamethoxazole; *T30*: oxytetracycline; *N30*: neomycin; *NOR1*:
90 norfloxacin; *FFC30*: florfenicol by the disc diffusion method. An overnight bacterial inoculum
91 (approx. 10^8 CFU ml⁻¹) was applied onto the surface of Mueller-Hinton agar plate before
92 placement of the antibiotic discs (BBL™ Sensi-Disc™, BD, NJ). *Streptococcus parauberis* RC
93 was resistant to T30 but susceptible to SXT, N30, NOR1 and FFC30. A strain of *S. parauberis*
94 from olive flounder (*Paralichthys olivaceus*) had similarly been previously identified to be
95 resistant to tetracycline (Park *et al.* 2009). Based on the results of the biogram, on-farm
96 treatment with florfenicol was applied through medicated feed. The treatment reduced the
97 mortalities, but the infection reoccurred when treatment was withdrawn. After four cycles of
98 repeated antibiotic treatments and reoccurrence of the disease, the farm started feeding the
99 fish with a diet supplemented with rosemary (*Rosmarinus officinalis*). Rosemary has been
100 previously reported to be effective against *Streptococcus iniae* and *Streptococcus agalactiae*
101 (Abutbul *et al.* 2004; Zilberg *et al.* 2010). Bacteria could not be isolated from fish during and
102 soon after the application of rosemary, but infection reoccurred once rosemary
103 supplementation was withdrawn.

104 Basic factors contributing to bacterial virulence were comparatively analyzed in our *S.*
105 *parauberis* RC isolate and the most common causative agents of streptococcosis in fish,
106 including *S. iniae* and *S. agalactiae*. Intra-community (*i.e.* biofilm, autoaggregation) and inter-
107 community interactions (*i.e.* co-aggregation) are common mechanisms of bacterial survival in
108 nature and have been identified to play a part in the virulence of pathogens, including in the
109 streptococci (Cvitkovitch *et al.* 2003; Khemaleelakul *et al.* 2006). Many aquatic bacteria are

110 capable of forming a biofilm, a dense aggregate of surface-adherent microorganisms
111 embedded in an exopolysaccharide matrix (Cvitkovitch *et al.* 2003; Branda *et al.* 2005).
112 Biofilm-forming ability was determined by a modified crystal violet assay protocol (Lazado *et*
113 *al.* 2010). *S. parauberis* RC was shown to be capable of forming biofilms under static (Fig. 2d)
114 or mobile (Fig. 2e) conditions. The biofilm forming potential of *S. parauberis* RC was similar to
115 that of *S. iniae* at both static and mobile conditions, and to *S. agalactiae* under mobile
116 conditions (Fig. 2d). Auto-aggregation allows cell-cell interactions to occur and has properties
117 similar to those of biofilms, providing protection from the host defense factors and from
118 external treatments, such as antibiotics (Aparna and Yadav 2008; Lazado *et al.* 2010). A
119 spectrophotometric-based assay was adopted to evaluate this feature (Lazado *et al.* 2011).
120 *Streptococcus parauberis* RC auto-aggregating index was calculated to be $23.4 \pm 5.68\%$ (Lazado
121 *et al.* 2011), indicating that around 23% of the individual bacteria clumped together.
122 Comparing to the other pathogenic streptococci, the capability is 19% higher than *S. iniae* but
123 43% lower than *S. agalactiae*. The ability of *S. parauberis* RC to aggregate provides insight to
124 the documented re-occurrence of infection following treatment withdrawal, *i.e.* this ability
125 may have provided protection and allowed the bacteria to survive the treatment.
126 Interestingly, *S. parauberis* RC was also capable to co-aggregating with the 2 pathogenic
127 streptococci (Rickard *et al.* 2003), with *S. iniae* $41.8 \pm 13.8\%$ and with *S. agalactiae*
128 $41.7 \pm 5.68\%$. This interaction suggests the potential for co-infection to occur.

129 We are speculating two probable causes of the presence of *S. parauberis* on the farm
130 where the bacterium was isolated. One likely scenario was that the bacteria originated from
131 incoming fish. Phylogenetic relationship of *S. parauberis* RC with other aquatic-derived

132 strains lends support to such a speculation. The farm personnel reported that there was no
133 delivery of fish to the farm for a long time period prior the outbreak, but its correlation with
134 fish handling could suggest that the bacterial infection was latent, and the resulting stress
135 may have caused an outbreak. Another likely scenario is the possibility of transmission of the
136 infection from an adjacent dairy facility.

137 The only reported fish-derived *S. parauberis* isolate (GenBank accession no.
138 JQ780604) in Israel was from a diseased broomtail wrasse (*Cheilinus lunulatus*). However,
139 this is the first report to discuss the histopathological changes associated with the infection
140 caused by an *S. parauberis* isolate from Israel in a freshwater ornamental fish. In addition,
141 some of the fundamental microbiological features characterized in *S. parauberis* RC may
142 offer insights in the subsequent study of the virulence and pathogenesis associated with this
143 pathogen.

144

145 **Acknowledgements**

146 The study was supported by research grants from the Ramat Negev Research and
147 Development, Israel, and the Central and Northern Arava Research and Development, Israel.
148 C. Lazado would like to thank the Jacob Blaustein Center for Scientific Cooperation for his
149 postdoctoral fellowship.

150

151

152

153 **References**

- 154 Abutbul S., Golan-Goldhirsh A., Barazani O. & Zilberg D. (2004) Use of *Rosmarinus officinalis*
155 as a treatment against *Streptococcus iniae* in tilapia (*Oreochromis* sp.). *Aquaculture* **238**, 97-
156 105.
- 157 Aparna M.S. & Yadav S. (2008) Biofilms: microbes and disease. *Brazilian Journal of Infectious*
158 *Diseases* **12**, 526-530.
- 159 Baeck G.W., Kim J.H., Gomez D.K. & Park S.C. (2006) Isolation and characterization of
160 *Streptococcus* sp. from diseased flounder (*Paralichthys olivaceus*) in Jeju Island. *Journal of*
161 *Veterinary Science* **7**, 53-58.
- 162 Bradley A.J. (2002) Bovine mastitis: an evolving disease. *The Veterinary Journal* **164**, 116-128.
- 163 Branda S.S., Vik Å., Friedman L. & Kolter R. (2005) Biofilms: the matrix revisited. *Trends in*
164 *Microbiology* **13**, 20-26.
- 165 Cvitkovitch D.G., Li Y.-H. & Ellen R.P. (2003) Quorum sensing and biofilm formation in
166 Streptococcal infections. *The Journal of Clinical Investigation* **112**, 1626-1632.
- 167 Dereeper A., Guignon V., Blanc G., Audic S., Buffet S., Chevenet F., Dufayard J.F., Guindon S.,
168 Lefort V., Lescot M., Claverie J.M. & Gascuel O. (2008) Phylogeny.fr: robust phylogenetic
169 analysis for the non-specialist. *Nucleic Acids Research* **36**, W465-W469.
- 170 Domeénech A., Derenaáandez-Garayzábal J.F., Pascual C., Garcia J.A., Cutuli M.T., Moreno
171 M.A., Collins M.D. & Dominguez L. (1996) Streptococcosis in cultured turbot, *Scophthalmus*
172 *maximus* (L.), associated with *Streptococcus parauberis*. *Journal of Fish Diseases* **19**, 33-38.
- 173 Haines A.N., Gauthier D.T., Nebergall E.E., Cole S.D., Nguyen K.M., Rhodes M.W. & Vogelbein
174 W.K. (2013) First report of *Streptococcus parauberis* in wild finfish from North America.
175 *Veterinary Microbiology* **166**, 270-275.
- 176 Hoshina T., Sano T. & Morimoto Y. (1958) A Streptococcus pathogenic to fish. *Journal of*
177 *Tokyo University of Fisheries* **44**, 57-68.
- 178 Inglis V., Roberts R.J. & Bromage N.R. (1993) *Chapter 12: Streptococcal infections*. NY: Halsted
179 Press, John Wiley & Sons, Inc.
- 180 Khemaleelakul S., Baumgartner J.C. & Pruksakom S. (2006) Autoaggregation and
181 coaggregation of bacteria associated with acute endodontic infections. *Journal of*
182 *Endodontics* **32**, 312-318.
- 183 Lazado C.C., Caipang C.M.A., Brinchmann M.F. & Kiron V. (2011) *In vitro* adherence of two
184 candidate probiotics from Atlantic cod and their interference with the adhesion of two
185 pathogenic bacteria. *Veterinary Microbiology* **148**, 252-259.


- 186 Lazado C.C., Caipang C.M.A., Rajan B., Brinchmann M.F. & Kiron V. (2010) Characterization of
187 GP21 and GP12: Two potential probiotic bacteria isolated from the gastrointestinal tract of
188 Atlantic cod. *Probiotics and Antimicrobial Proteins* **2**, 126-134.
- 189 Mata A.I., Gibello A., Casamayor A., Blanco M.M., Domínguez L. & Fernández-Garayzábal J.F.
190 (2004) Multiplex PCR Assay for detection of bacterial pathogens associated with warm-water
191 streptococcosis in fish. *Applied and Environmental Microbiology* **70**, 3183-3187.
- 192 Nho S.W., Hikima J.-i., Cha I.S., Park S.B., Jang H.B., del Castillo C.S., Kondo H., Hirono I., Aoki
193 T. & Jung T.S. (2011) Complete genome sequence and immunoproteomic analyses of the
194 bacterial fish pathogen *Streptococcus parauberis*. *Journal of Bacteriology* **193**, 3356-3366.
- 195 Noga E.J. (2010) *Fish disease: diagnosis and treatment* Iowa, USA: John Wiley & Sons.
- 196 Oguro K., Yamane J., Yamamoto T., Ohnishi K., Oshima S.-i. & Imajoh M. (2014) Draft genome
197 sequence of *Streptococcus parauberis* strain SK-417, isolated from diseased *Sebastes*
198 *ventricosus* in Kagoshima, Japan. *Genome Announcements* **2**.
- 199 Park Y.-K., Nho S.-W., Shin G.-W., Park S.-B., Jang H.-B., Cha I.-S., Ha M.-A., Kim Y.-R., Dalvi
200 R.S., Kang B.-J. & Jung T.-S. (2009) Antibiotic susceptibility and resistance of *Streptococcus*
201 *iniae* and *Streptococcus parauberis* isolated from olive flounder (*Paralichthys olivaceus*).
202 *Veterinary Microbiology* **136**, 76-81.
- 203 Rickard A.H., Gilbert P., High N.J., Kolenbrander P.E. & Handley P.S. (2003) Bacterial
204 coaggregation: an integral process in the development of multi-species biofilms. *Trends in*
205 *Microbiology* **11**, 94-100.
- 206 Russo R., Mitchell H. & Yanong R.P.E. (2006) Characterization of *Streptococcus iniae* isolated
207 from ornamental cyprinid fishes and development of challenge models. *Aquaculture* **256**,
208 105-110.
- 209 Salati F. (2006) *Enterococcus seriolicida* and *Streptococcus* spp. (*S. iniae*, *S. agalactiae* and *S.*
210 *dysgalactiae*). In *Fish diseases and disorders, volume 3: viral, bacterial and fungal infections*
211 eds. Woo, P.T.K. & Bruno, D.W. Oxfordshire, UK: CAB International
- 212 Toranzo A.E., Magariños B. & Romalde J.L. (2005) A review of the main bacterial fish diseases
213 in mariculture systems. *Aquaculture* **246**, 37-61.
- 214 Zilberg D., Tal A., Froyman N., Abutbul S., Dudai N. & Golan-Goldhirsh A. (2010) Dried leaves
215 of *Rosmarinus officinalis* as a treatment for streptococcosis in tilapia. *Journal of Fish Diseases*
216 **33**, 361-369.
- 217

218 **List of figure legends**

219 **Figure 1. Histopathology from *S. parauberis*-infected ram cichlid.** Infiltrating macrophages in
220 kidney tissue **(a)** containing Gram positive bacteria **(b)**. Liver appears vacuolated **(c)** with
221 focally occurring infiltrating macrophages, containing Gram positive bacteria **(d)**. Infiltrating
222 macrophages in the muscle **(e)** and focally occurring necrosis in muscle fibers **(f)**. Sections are
223 stained with H&E **(a, c, e, f)** and Gram stain **(b, c)**; m, macrophages.

224 **Figure 2. *S. parauberis* RC: Phylogeny, growth characteristics and biofilm formation. (a)**
225 Phylogram of *S. parauberis* from ram cichlid and other isolates of terrestrial and aquatic (*with*
226 *red arrowhead*) origins. The isolate with an arrowhead shaded in red and outlined in black
227 was previously isolated in Israel. Branches corresponding to partitions reproduced in less
228 than 50% bootstrap replicates are collapsed. The analysis involved 11 nucleotide sequences.
229 All positions containing gaps and missing data were eliminated. The alignment in MUSCLE
230 was curated in Glocks 0.91b to include a total of 635 positions, representing 40% of the
231 alignment. The curated alignment was used for phylogenetic analysis in PhyML and the tree
232 was rendered by TreeDyn. Culture conditions, including **(b)** temperature and **(c)** NaCl
233 concentration, affecting the growth of *S. parauberis* RC. Biofilm formation at 25°C either in
234 **(d)** static or **(e)** mobile conditions were analyzed in a microplate. For mobile conditions, the
235 plate was incubated with shaking (80 rpm). Values presented in **b, c, d** and **e** are mean ± SE of
236 observations from three independent experiments each with three replicate set-ups.
237 Column bars with different letters indicate significant difference ($P < 0.05$) as tested by one-
238 way ANOVA followed by Tukey's multiple comparison tests.






239



240

241 Figure 1.

242




243

244 Figure 2.

245

246 **Supplementary information**



247

248

249 Supplementary material 1. Gross pathology of ram cichlid (*Mikrogeophagus ramirezi*)
250 infected with *Streptococcus parauberis*.

251

252