



## ABSTRACT # 104

**DO PRIMATE CONSERVATION ACTION PLANS WORK?**

B. J. Morgan<sup>1,2,3,4</sup>, A. Dunn<sup>4,5</sup>, M. K. Gonder<sup>4,6</sup>, R. Kormos<sup>4</sup>, F. Maisels<sup>3,5</sup>, A. Nicholas<sup>4,5</sup>, J. Sunderland-Groves<sup>4,7</sup> and E. A. Williamson<sup>3,4</sup>

<sup>1</sup>Zoological Society of San Diego, 15600 San Pasqual Valley Road, Escondido, CA, 92027-7000, USA, <sup>2</sup>Ebo Forest Research Project, BP3055 Messa, Yaounde, Cameroon, <sup>3</sup>School of Natural Sciences, University of Stirling, UK, <sup>4</sup>IUCN/SSC Primate Specialist Group, <sup>5</sup>Wildlife Conservation Society, 2300 Southern Boulevard, Bronx, New York, USA, <sup>6</sup>Department of Biological Sciences, University at Albany – State University of New York, Albany, New York, USA, <sup>7</sup>Borneo Orangutan Survival Foundation, Indonesia

John Oates authored the first primate conservation Action Plan in 1986, which assessed the status of and proposed conservation actions for all mainland African primate species. A revised version of the continent-wide plan was published in 1996, but since then, action plans have generally evolved into prioritizing actions for specific species, often within defined landscapes. We will review and evaluate the content and success of conservation action plans for the nine currently recognized taxa of chimpanzees and gorillas in Africa. Since 2003, six detailed action plans and one population viability analysis have been published, covering priority actions and landscapes for seven of the nine great ape taxa in Africa. Two further action plans (for gorillas and chimpanzees in Eastern DRC and for bonobos) are in the final stages of review and may also be included in the analysis. Assessments for western chimpanzees, Cross River gorillas, western lowland gorillas and central chimpanzees have been peer reviewed, and we will consider their recommendations and the challenges of quantitatively evaluating the success of primate conservation action plans.

## ABSTRACT # 105

**MEAT, PLANT FOOD AND TOOL TRANSFER AMONG SAVANNA CHIMPANZEES: THE ACTIVE FEMALE**

J. D. Pruettz and S. Lindshield

Department of Anthropology, Iowa State University, Ames, Iowa, 50011, USA

Transferring food to mates or other unrelated social group members is uncommon in most animal species but common in humans. Hypotheses to explain provisioning behavior in nonhumans usually invoke reciprocal altruism, where the immediate or

delayed benefits of transferring food include enhancement of agonistic support and mating opportunity, or other social benefits. In chimpanzees (*Pan troglodytes*), meat transfer among nonrelatives is common, but plant food transfer is usually rare except between mother-offspring pairs. However, observations of plant food and tool transfer among savanna chimpanzees at Fongoli, Senegal contrast with previous research on chimpanzees in general and support findings that West and East African chimpanzee subspecies differ socially. In 45 cases, chimpanzees at Fongoli transferred plant foods and tools to unrelated individuals. Most observations involved males transferring resources to females, perhaps as a long-term strategy to increase males' copulation frequencies. Fongoli chimpanzees also share meat, similar to patterns seen elsewhere. Furthermore, we were able to examine meat transfer from the female perspective, since females at Fongoli hunt as often as males. A similar pattern of tolerance emerged. Males rarely monopolized carcasses, and while females transferred meat to males, they also effectively ignored males' begging behavior. We examine these relatively unusual food and tool transfer behaviors in Fongoli chimpanzees within the context of chimpanzee sociality and hominid evolution.

## ABSTRACT # 106

**REDUCTION OF REGURGITATION AND REINGESTION IN RETIRED LABORATORY CHIMPANZEES (*PAN TROGLODYTES*)**

G. Kranendonk<sup>1,2</sup>, H. Van Bolhuis<sup>1</sup>, A. Lange<sup>1</sup>, R. Huiskes<sup>1</sup>, M. Brüne<sup>3</sup> and B. Spruijt<sup>2</sup>

<sup>1</sup>AAP, Rescue Centre for Exotic Animals, P.O. Box 50.313, Almere, 1305 AH, The Netherlands,

<sup>2</sup>Behavioural Biology, Department of Biology, Utrecht University, the Netherlands, <sup>3</sup>Research Department of Cognitive Neuropsychiatry and Psychiatric Preventive Medicine, LWL-University

Hospital, Bochum University, Germany

When retired laboratory chimpanzees were relocated to a sanctuary, abnormal behaviour, mainly consisting of regurgitation and reingestion (R/R), rocking, and coprophagy, increased. This required intervention. Therefore, in April 2010, an intervention program was started to reduce abnormal behaviour in a group of six chimpanzees. A group of seven chimpanzees served as a control group. The intervention consisted of two main aspects. The first was an increase in environmental enrichment items, including an increase of low-energy/high-fiber contents in the diet. The second was psychopharmacological treatment using