



Limb Power Outputs and Cardiopulmonary Responses in Swimmers

Journal:	<i>International Journal of Sports Medicine</i>
Manuscript ID:	IJSM-02-2009-1164-tt.R2
Manuscript Type:	Training & Testing
Key word:	Swimmers, Power, Machine

SCHOLARONE™
Manuscripts

Review

1
2
3 **Reproducibility of Limb Power Outputs and Cardiopulmonary Responses to Exercise**
4 **Using a Novel Swimming Training Machine**
5
6
7
8

9
10 RUNNING TITLE: Novel Swimming Training Machine
11

12
13 **Abstract**
14

15 The purpose of this study was to determine the reproducibility of limb power output and
16 cardiopulmonary responses, to incremental whole-body exercise using a novel swimming
17 training machine. Eight swimmers with a mean age of 23.7 ± 4.6 (yrs), stature 1.77 ± 0.13 (m)
18 and body mass of 74.7 ± 2.8 (kg) gave informed consent and participated in repeat exercise
19 testing on the machine. All subjects performed two incremental exercise tests to exhaustion
20 using front crawl movements. From these tests peak oxygen consumption (VO_{2peak}), peak
21 heart rate (HR_{peak}), peak power output (W_{peak}) and individual limb power outputs were
22 determined. Results showed there were no significant differences between test 1 and 2 for any
23 variable at exhaustion, and the CV% ranged from 2.8% to 3.4%. The pooled mean values
24 were; VO_{2peak} 3.7 ± 0.65 L.min⁻¹, HR_{peak} 178.7 ± 6.6 b.min⁻¹ and W_{peak} 349.7 ± 16.5 W. The
25 mean contributions to the total power output from the legs and arms were (37.3 ± 4.1 % and
26 62.7 ± 5.1 % respectively). These results show that it is possible to measure individual limb
27 power outputs and cardiopulmonary parameters reproducibly during whole-body exercise
28 using this training machine, at a range of exercise intensities.
29
30
31
32
33
34
35
36
37
38
39

40 Key words

41 Novel machine, leg-kick, arm-pull
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Laboratory-based ergometers, such as the swim bench have been used previously to study swimmers [6, 11, 21, 23, 25-27]. This has been mostly to assess arm power [3, 6, 8, 12, 31] but also to assess cardiopulmonary responses to exercise [25, 27, 28]. More recently, Swaine [30] has reported arm and leg power in swimmers during separate laboratory-based ergometer tests which attempted to replicate the front crawl arm-stroke and leg-kick. Konstantaki et al. [14] have reported cardiopulmonary responses to simultaneous front crawl arm and leg exercise. However, there are no previous reports which have measured the individual limb power outputs at the same time as cardiopulmonary parameters during incremental whole-body simulated swimming.

In swimming itself, the relative contribution from each limb to the total power generated during front crawl is unknown. The absence of such data is largely because it has been impossible to measure the individual limb power outputs during swimming itself. This is because the force applied by the swimmer partly results in propulsion but is also partly dissipated in moving water [33]. Knowledge of the relative contributions to the total power output from each of the swimmer's limbs might enhance understanding of the principles of front crawl swimming.

Previously, it has been possible to measure the propulsive power that results from the work done by the swimmer [9] and the leg-kick was shown to contribute approximately 10% to the total front crawl swimming propulsive power. However, this study reports the resultant propulsive power during swimming and not the power output from the swimmer's limbs. Indeed, in a review by Toussaint and Beek [33] the assumption is made that since most propulsion comes from the arms in swimming, then the power output delivered by the swimmer will mainly come from the arms and trunk. However, there are no direct studies to support this. Quantification of power output from each limb during whole-body exercise on a machine might provide some insight into the contribution that each limb makes to the total power output of the swimmer during front crawl swimming.

1
2
3 The quantification of the contribution of each limb to the total power output during exercise
4 on a swimming machine requires an ergometer capable of measuring power output from each
5 limb and which allows manipulation of exercise intensity. A prototype ergometer has been
6 detailed previously [14, 29]. Such an ergometer would allow incremental testing to exhaustion
7 and it would permit the freely-chosen contribution that each limb makes, to total power
8 output, to be assessed through a range of exercise intensities. This type of measurement has
9 not been made previously. Accordingly, the purpose of this study was to assess the
10 reproducibility of the limb power outputs and cardiopulmonary responses to repeated
11 incremental exercise tests using a novel swimming training machine.
12
13
14
15
16
17
18
19

20 **Methods**

21 **Subjects**

22
23 Eight men of mean age 23.7 ± 4.6 (yrs), stature 1.77 ± 0.13 (m) and body mass 74.7 ± 2.8 (kg)
24 (mean \pm SD) performed two incremental exercise tests to exhaustion. The mean best times for
25 400m front crawl swimming within the 3 month period prior to testing were 262.7 ± 50 (s).
26 Six of the eight swimmers were right-arm dominant, and seven were 'bilateral breathers'
27 (with one swimmer breathing right-side only) during training. The participants were trained
28 swimmers who, for the 6 months prior to the testing, completed a minimum of 6 swimming
29 training sessions per week, of 1.5 hours duration. The study was conducted according to the
30 ethical standards of this journal [7] and all participants gave written informed consent prior to
31 participation and the study which was given approval by the University Ethics Committee.
32
33
34
35
36
37
38
39
40

41 **The swimming training machine**

42 Subjects performed exercise testing on a prototype novel whole-body swimming training
43 machine which was built for this study and can be seen in [Figure 1](#). Resistance to the
44 movement of each limb was created by four air-dynes (Lawler Engineering, UK) which were
45 mounted on spindles which rotated upon pay-out of pull-ropes, attached to hand-paddles or
46 foot-plates. The design of the leg-kick ergometer allowed force to be exerted in the upward
47 and downward kicking action but only during the pulling action of the arms, not during the
48
49
50
51
52
53
54
55
56
57
58
59
60

Deleted: f

1
2
3 recovery phase. On each air-dyne there was a photoelectric sensor which detected the
4 revolution of the air-dyne. The revolution rate that each air-dyne made was passed into a
5 computer where power was derived using software which contained a previously-determined
6 calibration algorithm. Subjects adopted a prone position and were instructed to simulate the
7 front crawl swimming action as closely as possible (including arm recovery), attempting to
8 achieve maximum pull and kick movements in each arm-stroke or leg-kick. Mean power
9 output for leg-kick and arm-stroke was averaged over each arm-pull or leg-kick. After
10 computation, the instantaneous mean power output for the combination of arm stroking and
11 leg-kicking was fed back to the swimmer on a visual display unit. The power output of the
12 swimmer was plotted against a 'target', so that the intensity of exercise could be manipulated.
13 This was done in a similar way to that first detailed in Swaine [25].
14
15
16
17
18
19
20
21
22

23 **Calibration of the air-dynes**

24
25 For each air-dyne the relationship between force applied and revolution rate was determined
26 by suspending known weights (0.5 to 4.5 kg) from the drive gear of the air-dynes. This
27 calibration technique has been used previously [23, 26]. An algorithm was derived for use in
28 the computer software used to feedback the power output to the swimmer.
29
30
31
32

33 **Measurement of cardiopulmonary variables**

34
35 Analysis of expired air was made using a breath-by-breath gas analysis system (Innocor,
36 Innovision, Denmark). This system uses laser diode absorption spectroscopy for oxygen and
37 photoacoustic spectroscopy for carbon dioxide analysis and a differential pressure sensor for
38 determination of ventilatory flow rates. The VO_{2peak} was defined as the highest oxygen
39 consumption value recorded. Heart rates were determined using a Polar heart rate monitor
40 (Polar AB, Finland) which gave instantaneous values at 5 s intervals.
41
42
43
44
45
46

47 **Determination of the reproducibility of peak responses to incremental exercise**

48
49 Assessment of the reproducibility of the peak responses to exercise required the subjects'
50 attendance at the laboratory on three occasions which were arranged at the same time of day.
51
52

1
2
3 The first visit was so that subjects could become accustomed to the swimming training
4 machine and the incremental exercise test procedures. On each visit the swimmers performed
5 an incremental exercise test to volitional exhaustion, however analysis of expired air was only
6 made on the second and third visits. During the incremental test the power output 'target' on
7 the visual display unit commenced at 100 W and was increased by 25 W.min⁻¹ and swimmers
8 were permitted to meet the increasing demands through a freely-chosen combination of work
9 from all four limbs at freely-chosen stroke rates. The maximal revolution rates of the air-dyne
10 devices was 15 s⁻¹ so that maximal pull velocity and stroke rates were close to those shown
11 previously to best replicate free swimming [27].
12
13
14
15
16
17
18
19

20 Analysis of data

Deleted: Data

21
22 Reproducibility was assessed by comparison of means (t) for test 1 and 2 and accompanied by
23 coefficient of variation (CV%). Differences in submaximal limb power outputs and
24 cardiopulmonary responses to the two tests were assessed using analysis of covariance
25 (ANCOVA) for comparison of the slopes and elevations of the linear relationships between
26 VO₂ and W, and between HR and W. This comparison was performed on individual and
27 group mean relationships. Differences in individual peak limb power outputs were analysed
28 using one-way analysis of variance (ANOVA) with TUKEY'S honestly significant difference
29 (HSD) post-hoc test. Population normality was checked using Shapiro-Wilks and Levene's
30 test was used to test for equality of variances.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

The mean values for VO_{2peak} , HR_{peak} and W_{peak} from test 1 and 2 are given in Table 1, along with statistical test values for differences, CV% and intra-class correlation. A Bland and Altman plot of test-retest differences in VO_{2peak} is given in Figure 2. There were no significant differences between test 1 and 2 for any variable at exhaustion ($P < 0.05$). The CV% (with 95% CI) were 2.8% (1.4 to 3.6) for VO_{2peak} ; 3.3% (2.0 to 5.4) for HR_{peak} ; and 3.4% (1.9 to 4.8) for W_{peak} . The pooled mean values of VO_{2peak} , HR_{peak} and W_{peak} were; 3.7 ± 0.65 L.min⁻¹, 178.7 ± 6.6 b.min⁻¹ and 349.7 ± 16.5 W respectively. On average at exhaustion, the total power output from each limb was: 111 ± 12.3 W (W-RA_{peak}); 108 ± 13.1 W (W-LA_{peak}); 68.1 ± 9.1 W (W-RL_{peak}) and 62.6 ± 8.3 W (W-LL_{peak}). ANOVA revealed significant differences in these mean limb peak power values ($F = 3.6$; $P = 0.012$). The Tukey HSD multiple comparisons revealed no significant difference between right and left arm ($P < 0.05$) or between right and left leg ($P = 0.01$), but the legs produced less power than the arms ($P < 0.01$) This equated to approximate contributions of 29%, 33%, 16% and 22% for right arm, left arm, right leg and left leg respectively. The contribution to the total power output from both legs and both arms were (37.3 ± 4.1 % and 62.7 ± 5.1 % respectively). An example of the contribution made by each limb throughout an incremental test for one subject is given in Figure 3.

The mean data for VO_2 at each incremental power output (W), for test_1 and test 2 are shown in Figure 4. Comparison of the submaximal relationships between VO_2 and W revealed that in all individuals this relationship was linear (r at least 0.89; $p < 0.05$). The slopes were different in 3 of the eight participants ($P < 0.05$), which precluded comparison of elevations. In the remaining 5 participants there was no difference in the slope ($P > 0.05$) or elevation ($P > 0.05$). The group mean relationship showed no difference in either slope ($P = 0.09$) or elevation ($P = 0.07$). For the HR vs W relationship, responses in all individuals were linear (r at least 0.93; $p < 0.05$). The slopes were different in 1 participant ($P < 0.05$), but in the remaining 7 participants there was no difference in the slope ($P > 0.05$) or elevation ($P > 0.05$). Again the group mean relationship for HR vs W showed no difference in either slope ($P = 0.1$) or elevation ($P = 0.09$).

Deleted:

Deleted: s

([Figures 1 to 4, Table 1 here](#))

Deleted: -----Page Break-----

Discussion

This study demonstrated that it is possible to assess cardiopulmonary responses to whole-body exercise which mimic the front-crawl swimming movements. Furthermore, it was possible to relate the cardiopulmonary responses to power output of the limbs throughout this exercise. The peak cardiopulmonary responses to this exercise were shown to have small repeat-test coefficient of variation between 2.8 and 3.4%. There are no previous measures of reproducibility for whole-body exercise using a swimming machine, with which to compare the data from the present study, but these values compare favourably with previous values for arms-only swim bench exercise of 1.0 to 2.1% [26]. These values also compare well with reproducibility of peak cardiopulmonary responses to treadmill running [23] and cycle ergometer exercise [1].

The reproducibility of the responses recorded on the novel swimming training machine can be used to inform sample size requirement for future studies, as advocated by Hopkins [11]. For a crossover or simple test-retest study the number of participants required is based around precision, defined by the 95% confidence limit, (deriving a power 0.8). Hopkins [11] calculates sample size as $n = 8s^2/d^2$, where n is the sample size, s is the typical error and d is the smallest worthwhile effect. If d is presented as a proportion (%) of mean group score then CV% can be inserted for s . These [figures](#) are changed to $n = 32s^2/d^2$ for a study using an experimental and control group. Using 0.2 of the between-subject variation as the smallest worthwhile change [5], the sample sizes required for these types of studies (based on the VO_{2peak} data) are presented in Table 2. These calculations suggest that this machine, protocol and gas analysis system could be used with relatively small groups of swimmers to detect small changes in VO_{2peak} . Indeed, an expected change in VO_{2peak} of ~2% would require between 4 and 26 participants, based on the current findings. This analysis indicates that the methods used in the present study could comfortably detect the large (11%) training-induced changes in VO_{2peak} reported previously in recreational swimmers [16] and might be able to detect the much smaller changes consistent with ergogenic supplementation.

Deleted: s

Deleted: figures

1
2
3 [\(Table 2 here\)](#)

4 There are only two previous studies with which to compare our whole-body incremental
5 exercise measurements [13,14]. Konstantaki *et al.*, [14] reported $\text{VO}_{2\text{peak}}$ values of $2.85 \text{ L}\cdot\text{min}^{-1}$
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
There are only two previous studies with which to compare our whole-body incremental exercise measurements [13,14]. Konstantaki *et al.*, [14] reported $\text{VO}_{2\text{peak}}$ values of $2.85 \text{ L}\cdot\text{min}^{-1}$ for combined arm-stroke and leg-kick, but their measurements were made on younger swimmers. Kimura *et al.*, [13] used simultaneous arm-cranking and leg-kicking and measured $\text{VO}_{2\text{peak}}$ in varsity swimmers. In their study it was $3.6 \pm 0.3 \text{ L}\cdot\text{min}^{-1}$ and in ours it was $3.7 \pm 0.63 \text{ L}\cdot\text{min}^{-1}$, therefore our results confirm those of these previous studies. In the variety of studies that have assessed $\text{VO}_{2\text{peak}}$ during flume and tethered swimming, the values have been 2.6 to 3.2 lmin^{-1} [2, 10, 15, 20]. Therefore, our results for exercise using this novel swimming training machine appear to be similar to those of free swimming.

The individual limb peak power output values were somewhat surprising, especially the mean contribution made by the leg-kick (37.3%). These results could be suggestive that a much greater proportion of the total power output during swimming might be done by the legs than previously thought. Certainly, these relative contributions from arms and legs cast doubt on the previous suggestions that the power output delivered by the swimmer comes almost entirely from the arms and trunk [33]. Indeed, although we did not make a systematic analysis of the contribution that the legs made through all intensities of exercise, several of our swimmers generated higher power outputs from the legs than from the arms when nearing exhaustion.

There are no previous power output values with which to compare our results for whole-body power output using a swimming training machine. Swaine [28] reported separate peak power output values, at exhaustion, during incremental exercise using front crawl arm-stroke and leg-kick tests. The sum of the separate values from that study (170 W for arm-stroke and 141 W for leg-kick) is similar to our mean of 352 W but perhaps suggests that it is possible to generate a small amount of additional power by combining the upper and lower limbs in a synchronised way. Front crawl swimming is known to involve highly-skilled co-ordination of the leg-kick and arm-stroke in such a way as to achieve greatest forward propulsion [4].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Toussaint *et al.* [32] has reported power output values of swimmers from measurements using the fixed push-off pads of the MAD system [9]. However, measurements were only made for arm-stroke power output (leg-kick power was derived by subtracting arm power from whole-stroke power). Also, in that study it was not possible to precisely manipulate the exercise intensity in the same way that we did. Therefore, it is difficult to compare our power output values to those reported in the study by Toussaint *et al.* [32]. However, his reported power output values, estimated at 1000 W of 'power input', were between 50 and 120 W (as derived from arms-only swimming) which are much lower than our power output values and would represent approximately 20-30% of whole-body exercise power output at exhaustion in our subjects. Unfortunately, it is not possible to determine exactly what relative exercise intensity 1000 W represented in the study of Toussaint *et al.* [32] and therefore it is not possible to directly compare our power output values with such previous studies.

Simultaneous arm and leg work has been studied previously, using arm-cranking and cycling [22]. However, the freely-chosen contribution from the arms and legs was not assessed in this study. Rather, the contribution was set by the investigator. Nevertheless, this study showed that the additional oxygen uptake achieved when adding arm- to leg-work represented approximately 15% in this type of exercise. Similarly, Swaine and Zanker [27] showed that the VO_{2peak} during swim bench arms-only exercise was approximately $2.9 \text{ L}\cdot\text{min}^{-1}$ and our results of $3.8 \text{ L}\cdot\text{min}^{-1}$ for whole-body exercise would represent a $0.9 \text{ L}\cdot\text{min}^{-1}$ or 31% increase in arms-only VO_{2peak} , due to addition of the leg-kick. Therefore, this difference in VO_{2peak} must be considered when interpreting data from previous reports that have used swim bench exercise.

The extent to which combining the arm-stroke and leg-kick enhances the total power output of the swimmer, during whole-body exercise, remains to be determined. It is unlikely to be simply represented by the sum of the separate arm- and leg-power outputs reported by Swaine [30]. The combining of the upper and lower body during swimming is known to enhance total

1
2
3 power output by allowing transfer of energy between the limbs [17]. The combining of arm-
4 stroke and leg-kick is likely to affect the total efficiency, at any given exercise intensity.
5
6 Therefore, it might be expected that the oxygen cost for a given exercise intensity using
7
8 combined arm-stroke and leg-kick would be lower than the sum of the oxygen cost values for
9
10 separate arm-stroke and leg-kick. This could be studied by using the whole-body swimming
11
12 training machine.

13
14
15 Also, it was notable during our investigation that ventilatory thresholds (as identified using
16
17 breath-by-breath gas exchange) appeared to coincide with changes in the contribution of each
18
19 limb to the total power output. Although it was not the purpose of this study to identify
20
21 ventilatory thresholds, it appeared that quite marked changes occurred at these thresholds.
22
23 Therefore, it might be useful in future studies, to systematically investigate the pattern of
24
25 change in limb contribution, during incremental simulated swimming, and relate this pattern
26
27 to the gas exchange or lactate markers of the onset of anaerobiosis. Currently, it is not known
28
29 how the relative contributions of the arm-pull and leg-kick contribute to fatigue (lactate
30
31 accumulation) or how fatigue influences the freely-chosen relative contribution from each
32
33 limb.

34
35 Of course, we acknowledge that there are significant differences in the movement patterns of
36
37 exercise using a machine, compared to 'free swimming' and this presents a limitation to the
38
39 direct comparison between measurements made in the laboratory and during swimming itself.
40
41 For example, during exercise on a machine there is restriction of body roll, which is known to
42
43 be an important aspect of front crawl swimming [19]. However, the arrangement of the
44
45 swimmer in a suspended 'cradle' appeared to allow greater body roll than that seen in
46
47 previous swim bench work [25, 27]. Also, the 'cradle' that we used was foreshortened so that
48
49 it did not restrict the movement of the thoracic cavity as much as has been experienced when
50
51 using the swim bench [26]. However, it has been shown previously, through EMG
52
53 measurement, that the movement patterns of dry land training devices are quite different to
54
55 those of free swimming [18].
56
57
58
59
60

In addition to the restricted body roll, the swimming training machine cannot quantify power generated by the swimmer's limbs during many of the lateral or rotational movements of the hands and feet. These movements are known to be important components of the front crawl swimming technique. Indeed, the current arrangement of the pulley-ropes means that tension is only developed in the direction dictated by the fixed point of the pulley. This presents a further limitation of exercise on the swimming training machine. Therefore, machine-based exercise in swimmers has limitations in its direct application to free swimming. Of course, future studies with this novel machine would be required, to establish the extent to which power output from the limbs and cardiopulmonary responses during such exercise are reflective of swimming itself.

References

- 1 *Astrand I, Astrand P-O, Christensen EH, Hedman R.* Circulatory and respiratory adaptations to severe muscular work. *Acta Physiol Scand* 1960; 50: 250-259
- 2 *Bonen A, Wilson BA, Yarkony M, Belcastro AN.* Maximal oxygen uptake during free, tethered and flume swimming. *J App Physiol* 1980; 48: 232-235
- 3 *Bradshaw A, Hoyle J.* Correlation between sprinting and dry land power. *J Swim Res* 1993; 9: 15-18
- 4 *Bucher W.* Influence of the leg kick and the arm stroke on the total speed during the crawl stroke. In Lewillie L. Clarys JP (eds). *Proceedings of the 2nd International Symposium on Biomechanics in Swimming*. Brussels, Belgium, 1974, Baltimore: University Park Press, 1975; 180-187
- 5 *Cohen J.* *Statistical Power Analysis for the Behavioural Sciences*. 2nd Edition. Hillsdale NJ: Lawrence Erlbaum, 1988

Deleted: E

Deleted: i

Deleted: symposium

Deleted: biomechanics

Deleted: swimming

Deleted: ,

Deleted: ,

Deleted: Baltimore,

Deleted: ; ISBN: 0-83910-817-6, pp

Deleted: .

Deleted: power

Deleted: a

Deleted: behavioural

Deleted: sciences

Deleted: H

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 6 *Costill DL, King DS, Thomas RT, Hargreaves M.* Effects of reduced training on muscular power in swimmers. *Physician Sports Med* 1985; 13: 94-98; 100-101
- 7 *Harriss DJ, Atkinson G.* International Journal of Sports Medicine – Ethical Standards in Sport and Exercise Science Research. *Int J Sports Med* 2009; 30: 701-702
- 8 *Hawley JA, Williams MM, Vickovic MM, Handcock PJ.* Muscle power predicts freestyle swimming performance. *Br J Sports Med* 1992; 26: 151-155
- 9 *Hollander AP, De Groot G, van Ingen-Schenau GJ, Toussaint HM.* Contribution of the legs to propulsion in front crawl swimming. In: Ungerechs BE, Wilke K, Reischle K (eds). *Swimming Science V. Human Kinetics, Champaign, Illinois 1988; 39-45*
- 10 *Holmer I.* Energy cost of arm stroke, leg kick and the whole stroke in competitive swimming styles. *Eur J Appl Physiol* 1974; 33: 105-118
- 11 *Hopkins WG.* Measures of reliability in sports medicine and science. *Sports Med* 2000; 30: 1-15
- 12 *Johnson RE, Sharp RL, Hendrick MS.* Relationship of swimming power and dryland power to sprint freestyle performance: A multiple regression approach. *J Swim Res* 1993; 9: 10-14
- 13 *Kimura Y, Yeater RA, Martin RB.* Simulated swimming: a useful tool for evaluation the VO₂max of swimmers in the laboratory. *Br J Sports Med* 1990; 24: 201-206
- 14 *Konstantaki M, Winter EM, Swaine IL.* Peak oxygen uptake responses to free and simulated swimming using different body segments. *J Swim Res* 2004; 16: 18-25
- 15 *Magel JR, Faulkner JA.* Maximum oxygen uptake of college swimmers. *J Appl Physiol* 1967; 22: 929-938
- 16 *Magel JR, Foglia GF, McArdle WD, Gutin B, Pechar GS, Katch FL.* Specificity of swim training on maximum oxygen uptake. *J Appl Physiol* 1975; 38: 151-155

Deleted: and

Deleted: it

Deleted:

Deleted: Eds

Deleted: ,

Deleted: ;

Deleted: ISBN 0-87322-108-7, pp.

Deleted: it

Deleted: .

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 17 Maglischo E. Swimming ~~Faster~~. ~~California, USA~~: Mayfield Publishing Company, 1982; 81
- 18 Olbrecht J, Clarys JP. EMG of ~~Specific Dry Land Training~~ for the ~~Front Crawl~~. In: Huijing PA, de Groot G, Hollander JP (eds). Biomechanics and Medicine in Swimming IV, Champaign, Illinois: ~~Human Kinetics~~, 1983; 136-141
- 19 Payton CJ, Bartlett R. Estimation of propulsive forces in swimming from three-dimensional kinematic data. J Sports Sci 1995; 13: 447-454
- 20 Rinehardt KF, Kraemer RR, Gormley S, Colan S. Comparison of maximal oxygen uptakes from tethered, the 183 and 457 meter unimpeded supramaximal freestyle swims. Int J Sports Med 1991; 12: 6-9
- 21 Roberts AJ, Termin B, Reilly MF, Pendergast DR. Effectiveness of biokinetic training on swimming performance in collegiate swimmers. J Swim Res 1991; 7: 5-11
- 22 Secher NH, Ruberg-Larsen N, Binkhorst RA, Bonde-Petersen F. Maximal oxygen uptake during arm cranking and combined arm plus leg exercise. J Appl Physiol 1974; 36: 515-518
- 23 Sharp RL, Troup JP, Costill DL. Relationship between power and sprint freestyle swimming. Med Sci Sports Exerc 1982; 14: 53-56
- 24 Shephard RJ, Allen C, Benade AJS, Davies CTM, Di Prampero PE, Hedman R, Merrisman JE, Myhre K, Simmons R. The maximum oxygen uptake. An international reference standard of cardiorespiratory fitness. Bull World Health Organ 1968; 38: 757-764
- 25 Swaine IL. The relationship between physiological variables from a swim bench ramp test and middle-distance swimming performance. J Swim Res 1994; 10: 41-48
- 26 Swaine IL, Reilly T. The freely chosen swimming stroke rate in a maximal swim and on a biokinetic swim bench. Med Sci Sports Exerc 1983; 15: 370-375

Deleted: f

Deleted: California, USA,

Deleted: ; ISBN 0-87484-548-3; pp

Deleted: .

Deleted: specific

Deleted: dry

Deleted: land

Deleted: training

Deleted: front

Deleted: crawl

Deleted: .

Deleted: .

Deleted: G.

Deleted: and J.P.

Deleted: Eds

Deleted: . Human Kinetics,

Deleted: ;

Deleted: ISBN 0-931250-50-1, pp

Deleted: (5)

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 27 Swaine IL, Zanker CL. The reproducibility of cardiopulmonary responses to exercise using a swim bench. *Int J Sports Med* 1996; 17: 140-144
- 28 Swaine IL. Cardiopulmonary responses to exercise in swimmers using a swim bench and a leg-kicking ergometer. *Int J Sports Med* 1997; 18: 359-363
- 29 Swaine IL, Reavell C, Winter EM, Cooke DS, Maycock KM. A Dry-Land Ergometer for Measurement and Manipulation of Power Output During Front Crawl Arm-Pulling, Leg-Kicking and Whole Body Simulated Swimming. In: Haake SJ (ed). *The Engineering of Sport*. Oxford: Blackwell Science, 1998; 93-98
- 30 Swaine IL. Arm and leg power output in swimmers during simulated swimming. *Med Sci Sports Exerc* 2000; 32: 1288-1292
- 31 Takahashi S, Bone M, Cappaert JM, Barzdukas A, D'Acquisto L, Hollander AP, Troup JP. Validation of a Dryland Swimming Specific Measurement of Anaerobic Power. In: MacLaren D, Reilly T, Lees A, (eds). *Biomechanics and Medicine in Swimming VI*. London: E & FN Spon, 1992; 301-305
- 32 Toussaint HM, Knops W, DeGroot G, Hollander AP. The mechanical efficiency of front crawl swimming. *Med Sci Sports Exerc* 1990; 22: 402-408
- 33 Toussaint HM, Beek PJ. Biomechanics of competitive front crawl swimming. *Sports Med* 1992; 13: 8-24

Deleted: l

Deleted: p

Deleted: k

Deleted: .

Deleted: Ed

Deleted: Oxford,

Deleted: ;

Deleted: ISBN: 0-632-05048-9, pp

Deleted: dryland

Deleted: swimming

Deleted: specific

Deleted: measurement

Deleted: anaerobic

Deleted: power

Deleted: .

Deleted: Eds

Deleted: .

Deleted: London,

Deleted: ; ISBN 0-585-45686-0, pp.

1
2
3 **Table 1** Mean values and reproducibility measures for peak oxygen consumption (VO_{2peak}),
4 peak heart rate (HR_{peak}) and peak total power output (W_{peak}) from the novel
5 swimming training machine.
6
7

8 **Table 2** Sample size calculations based on the methods described by Hopkins (2000), using
9 the 95% confidence interval for the CV (1.4 – 3.6%). The smallest worthwhile
10 change (d) was derived as 0.2 of the between-subject variation from (a) the pooled
11 data from the current study, (b) the pre-intervention data of Magel et al. (1975), and
12 (c) tethered swimming data from Bonen et al. (1980).
13
14
15
16
17
18
19

20 **Figure 1** Photograph of the prototype novel swimming training machine showing, the leg air-
21 dynes to the rear, the use of a suspended ‘cradle’ for body support, and pulley ropes
22 used to drive the four air dynes.
23
24

25 **Figure 2** A Bland and Altman plot of individual test-retest differences in VO_{2peak}
26

27 **Figure 3.** An example of the relative contribution of each limb to total power output
28 during incremental whole-body simulated swimming to exhaustion.
29

30 **Figure 4.** The group mean data for VO_2 and total power output (W) during incremental
31 exercise tests 1 and 2, using the novel swimming training machine.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 1. Photograph of the novel swimming training machine.
609x450mm (96 x 96 DPI)

review




Figure 2. An example of the relative contribution of each limb to total power output during exhaustive incremental exercise using the novel swimming training machine.
164x114mm (96 x 96 DPI)




Figure 3. The group mean relationships between VO₂ and total power output (W) during incremental exercise test 1 and test 2, using the novel swimming training machine.
148x95mm (96 x 96 DPI)




Figure 4. The group mean data for VO₂ and total power output (W) during incremental exercise tests 1 and 2, using the novel swimming training machine.

Table 1 Mean values and reproducibility measures for peak oxygen consumption (VO_{2peak}), peak heart rate (HR_{peak}) and peak total power output (W_{peak}) from the novel swimming training machine.

	Mean test 1	Mean test 2	t-test	CV%	Correlation
VO_{2peak} ($L \cdot min^{-1}$)	3.68 ± 0.65	3.72 ± 0.61	$t=1.4$; $p=0.7$	3.4	$r=0.94$; $p=0.01$
HR_{peak} ($b \cdot min^{-1}$)	177.7 ± 6.6	180.2 ± 6.2	$t=3.7$; $p=0.01$	2.8	$r=0.92$; $p=0.01$
W_{peak} (Watts)	345.7 ± 15.2	356.7 ± 16.0	$t=4.9$; $p=0.01$	3.1	$r=0.90$; $p=0.02$

Table 2. Sample size calculations based on the methods described by Hopkins (2000), using the 95% confidence interval for the CV (1.4 – 3.6%). The smallest worthwhile change (*d*) was derived as 0.2 of the between subject variation from (a) the pooled data from the current study, (b) Magel et al's [15] pre intervention data, and (c) Bonen et al's [2] tethered swimming data.

VO _{2peak}	SD	<i>d</i> (0.2 SD)	Participant number required for a simple test retest experiment.	Participant number required for a study with an experimental and control group.
3.70 ^a	0.65	3.5%	2 – 9	6 – 34
3.44 ^b	0.49	2.9%	2 – 13	8 – 50
3.53 ^c	0.27	1.5%	7 – 47	28 – 184