

LETTER • OPEN ACCESS

Advancing conservation planning for western chimpanzees using IUCN SSC A.P.E.S.—the case of a taxon-specific database

To cite this article: Stefanie Heinicke *et al* 2019 *Environ. Res. Lett.* **14** 064001

View the [article online](#) for updates and enhancements.

Environmental Research Letters



LETTER

Advancing conservation planning for western chimpanzees using IUCN SSC A.P.E.S.—the case of a taxon-specific database

OPEN ACCESS

RECEIVED

23 November 2018

REVISED

25 March 2019

ACCEPTED FOR PUBLICATION

26 March 2019

PUBLISHED

22 May 2019

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Stefanie Heinicke^{1,2,29} , Roger Mundry¹, Christophe Boesch^{1,3}, Bala Amarasekaran⁴, Abdulai Barrie⁵, Terry Brncic⁴, David Brugière⁶, Geneviève Campbell⁷, Joana Carvalho⁸, Emmanuel Danquah⁹ , Dervla Dowd³, Henk Eshuis¹, Marie-Claire Fleury-Brugière¹⁰, Joel Gamys¹¹, Jessica Ganas¹², Sylvain Gatti¹³, Laura Ginn¹⁴, Annemarie Goedmakers¹⁵, Nicolas Granier¹⁶, Ilka Herbinger¹⁷, Annika Hillers^{3,18}, Sorrel Jones^{18,19}, Jessica Junker^{1,2}, Célestin Y Kouakou^{3,20,21}, Vincent Lapeyre³, Vera Leinert¹, Fiona Maisels^{22,23}, Sergio Marrocoli^{1,2}, Mary Molokwu-Odozi²⁴, Paul KN'Goran²⁵, Liliana Pacheco²⁶, Sébastien Regnaut^{3,27}, Tenekwetché Sop^{1,2}, Els Ton¹⁵, Joost van Schijndel^{1,15}, Virginie Vergnes³, Maria Voigt^{1,2}, Adam Welsh¹, Erin G Wessling^{1,2} , Elizabeth A Williamson^{23,28} and Hjalmar S Kühl^{1,2,28} 

¹ Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

² German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

³ Wild Chimpanzee Foundation (West Africa Representation), Abidjan, Côte d'Ivoire

⁴ Tacugama Chimpanzee Sanctuary, Freetown, Sierra Leone

⁵ Bumbuna Watershed Management Authority, Freetown, Sierra Leone

⁶ Projets Biodiversité et Ressources Naturelles BRL Ingénierie, Nîmes Cedex, France

⁷ The Biodiversity Consultancy Ltd, Cambridge, United Kingdom

⁸ Faculty of Computing Science and Mathematics, University of Stirling, Stirling, United Kingdom

⁹ Department of Wildlife and Range Management, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

¹⁰ Independent consultant, Nîmes, France

¹¹ Conservation International, Monrovia, Liberia

¹² Independent Biodiversity Consultant, Slinger, WI, United States of America

¹³ West African Primate Conservation Action, Accra, Ghana

¹⁴ Anthropology Center for Conservation, Environment and Development, Oxford Brookes University, Oxford, United Kingdom

¹⁵ Chimbo Foundation, Amsterdam, The Netherlands

¹⁶ Behavioural Biology Unit, University of Liège, Liège, Belgium

¹⁷ WWF Germany, Berlin, Germany

¹⁸ RSPB Centre for Conservation Science, The Royal Society for the Protection of Birds, Sandy, United Kingdom

¹⁹ School of Biological Sciences, Royal Holloway University of London, Egham, United Kingdom

²⁰ Université Jean Lorougnon Guédé, Daloa, Côte d'Ivoire

²¹ Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire

²² Global Conservation Program, Wildlife Conservation Society, Bronx, NY, United States of America

²³ Faculty of Natural Sciences, University of Stirling, Stirling, Scotland, United Kingdom

²⁴ Fauna & Flora International, Monrovia, Liberia

²⁵ World Wide Fund for Nature, Regional Office for Africa - Yaoundé Hub, Yaoundé, Cameroon

²⁶ Jane Goodall Institute Spain, Dindefelo Community Nature Reserve, Kédougou, Senegal

²⁷ International Union for Conservation of Nature, West and Central Africa Programs, Ouagadougou, Burkina Faso

²⁸ Shared last author.

²⁹ Author to whom any correspondence should be addressed.

E-mail: stefanie_heinicke@eva.mpg.de

Keywords: species distribution model, *Pan troglodytes verus*, development corridor, West Africa

Supplementary material for this article is available [online](#)

Abstract

Even though information on global biodiversity trends becomes increasingly available, large taxonomic and spatial data gaps persist at the scale relevant to planning conservation interventions. This is because data collectors are hesitant to share data with global repositories due to workload, lack of incentives, and perceived risk of losing intellectual property rights. In contrast, due to greater conceptual and methodological proximity, taxon-specific database initiatives can provide more direct benefits to data collectors through research collaborations and shared authorship. The IUCN SSC Ape Populations, Environments and Surveys (A.P.E.S.) database was created in 2005 as a repository for data on great apes and other primate taxa. It aims to acquire field survey data and make different types of data accessible, and

provide up-to-date species status information. To support the current update of the conservation action plan for western chimpanzees (*Pan troglodytes verus*) we compiled field surveys for this taxon from IUCN SSC A.P.E.S., 75% of which were unpublished. We used spatial modeling to infer total population size, range-wide density distribution, population connectivity and landscape-scale metrics. We estimated a total abundance of 52 800 (95% CI 17 577–96 564) western chimpanzees, of which only 17% occurred in national parks. We also found that 10% of chimpanzees live within 25 km of four multi-national ‘development corridors’ currently planned for West Africa. These large infrastructure projects aim to promote economic integration and agriculture expansion, but are likely to cause further habitat loss and reduce population connectivity. We close by demonstrating the wealth of conservation-relevant information derivable from a taxon-specific database like IUCN SSC A.P.E.S. and propose that a network of many more such databases could be created to provide the essential information to conservation that can neither be supplied by one-off projects nor by global repositories, and thus are highly complementary to existing initiatives.

1. Introduction

In conservation planning there is an increasing need for detailed information on the density distribution of species, population trends, and habitat suitability to support evidence-based decision-making (Schwartz *et al* 2018). To derive these parameters different types of data are needed across large areas, an extent that usually exceeds the scope of individual research projects. Consequently, the curation of existing data has been the focus of various databases, many of them compiling data at a global scale, such as the Global Biodiversity Information Facility (GBIF 2018), Map of Life (Jetz *et al* 2012), and Living Planet Index (Collen *et al* 2009). However, large data gaps remain regarding spatial and taxonomic coverage and type of data, especially for Africa and the Middle East, and occurrence data are more readily available than abundance or trend data (Boakes *et al* 2010, Kindsvater *et al* 2018, Peterson and Soberón 2018).


For many taxa the challenge is not necessarily that data are not available. In contrast, researchers and conservation organizations go to great lengths to collect species survey data, which requires extensive human and financial resources. However, only a fraction of these data are shared, despite their value for broad-scale and comparative analyses (Costello *et al* 2013). Impediments to data sharing include first and foremost a lack of perceived benefit, workload, and concern of losing intellectual property (Thessen and Patterson 2011). In addition, the amount of survey data published in the gray literature greatly exceeds that published in peer-reviewed journals (Corlett 2011). To inform conservation planning, these data need to be centralized, standardized, and quality checked, whilst assuring data collectors of their intellectual property rights (Reichman *et al* 2011, Thessen and Patterson 2011, Costello *et al* 2013).

Apes are particularly well studied (Wich and Marshall 2016), and western chimpanzees (*Pan troglodytes verus*) are no exception, as illustrated by the IUCN Status Survey and Conservation Action Plan (Kormos *et al* 2003). However, the action plan stated that

information available at the time was insufficient because data were only available for specific sites, and large data gaps remained. Consequently, the identification of priority areas for conservation activities was based on expert opinion (Kormos and Boesch 2003). Besides the call for filling data gaps (Kormos and Boesch 2003), conservationists and researchers saw the need of compiling available ape survey datasets and make them accessible through a platform to better inform conservation planning.

The IUCN SSC Ape Populations, Environments and Surveys database (A.P.E.S.; Kühl *et al* 2007) was initiated in 2005, and its creation was facilitated by a collaboration between the section on Great Apes of the IUCN Primate Specialist Group, ape range country authorities, academic institutions, and conservation organizations. IUCN SSC A.P.E.S. contains geo-referenced survey data of all 14 taxa of extant great apes covering 21 of the 23 ape range countries (data not available for South Sudan, and Cabinda Province in Angola, figure 1). The database holds almost 500 standardized and quality-checked datasets consisting of more than three million records, including information on abundance, density, population trends, presence-absence, and spatial distribution (as of November 2018).

For western chimpanzees, data deposited in IUCN SSC A.P.E.S. have been used to predict the distribution of habitat suitability and its trends (Junker *et al* 2012, Jantz *et al* 2016), and to determine their population trend as well as geographic range (Kühl *et al* 2017). These assessments estimated a population decline of 80% and a range reduction of 20% within 24 years (Kühl *et al* 2017). As a result, western chimpanzees were uplisted to Critically Endangered by the IUCN Red List of Threatened Species (Humble *et al* 2016). Currently, the conservation action plan for this taxon is being updated. As various researchers and conservation organizations conducted surveys on this ape in the past 15 years and shared their data with IUCN SSC A.P.E.S., we were now able to use a dataset representative of the entire range of western chimpanzees, 75% of which had not been published. For the first time, we



could thus model the range-wide density distribution for this ape. We then derived information on western chimpanzees important for the update of the conservation action plan, including areas with high chimpanzee densities, or those where population connectivity has been reduced, estimated total abundance, and proportion of chimpanzees occurring in proximity to settlements and infrastructure.

2. Methods

2.1. IUCN SSC A.P.E.S. database

The IUCN SSC A.P.E.S. database currently holds 498 survey datasets contributed by more than 200 conservation scientists, wildlife authorities, and non-governmental organizations. A dataset is defined as a set of data that was collected for a specific area and

time period. Datasets had been collected as part of single or repeated surveys, and range from small scale (20 km²) to large areas, or even entire countries. The database stores different types of data that are standardized and quality-checked, including point and reconnaissance survey data that can be used to determine presence and absence, line transect data which are the basis for density estimates, and camera trap data. IUCN SSC A.P.E.S. also holds 280 abundance polygons, meaning abundance estimates for various resource management areas such as protected areas or resource concessions. The available data also include 24 spatial layers, for example, species density distribution and range layers, abundance layers, and suitable ecological conditions layers. Additionally, IUCN SSC A.P.E.S. contains nest decay datasets, which are needed to convert counts from ape nest surveys into individual ape density and abundance estimates. Most datasets not only include sightings of the ape taxa targeted in the survey, but also include phenology of ape food plants, records of other taxa, human signs, and records of covariates such as vegetation type and slope. Furthermore, the database stores 950 publications and reports, of which 280 are unpublished field survey reports (as of November 2018). For this study we used 52 chimpanzee nest count datasets, only 11 of which had been published to date (figure 1, table S1 is available online at stacks.iop.org/ERL/14/064001/mmedia).

2.2. Modeling chimpanzee density distribution

We followed a commonly used procedure to predict ape density distributions (Murai *et al* 2013, Wich *et al* 2016, Strindberg *et al* 2018, Voigt *et al* 2018). Specifically, we first fitted a full model to establish the relationship between chimpanzee densities and several social-ecological predictor variables, and then predicted chimpanzee density distribution based on multi-model inference (Burnham and Anderson 2002).

The response variable in the full model was the number of nests per transect with a sample size of 17 109 transects and a total survey effort of 10 929 km, covering all western chimpanzee range states (figure 1(a)). For the model output to directly express number of individuals per km² and to account for varying transect lengths, we included an offset term comprising transect length, effective strip width, proportion of nest builders, nest production rate, and nest decay time (details supplementary material). We then extracted 20 predictor variables for each transect using publicly available satellite and aggregated household-survey data which approximate known drivers of chimpanzee density including both environmental variables and anthropogenic pressure (details in table S2 and table S3). We originally started with a model comprising the same predictors used in an earlier study to identify drivers of chimpanzee densities

(Heinicke *et al* 2019), but the initial evaluation of the derived density distribution revealed an underestimation of chimpanzee densities for protected areas. We therefore added ‘protected area’ as a binary predictor, meaning whether the midpoint of a transect was within the boundaries of a protected area designated as ‘national park’ or IUCN category I or II based on data from the World Database of Protected Areas (UNEP-WCMC and IUCN 2017). For the full model (table S4), we fitted a Generalized Linear Mixed Model (Baayen 2008) with a negative binomial error distribution (Hilbe 2011). Details on model implementation, namely spatial autocorrelation, random effects, check for multicollinearity and overdispersion can be found in the supplementary material.

We then extracted all predictors across the entire range of western chimpanzees by deriving a grid with a resolution of half a minute (ca. 0.9 km) and identifying the coordinates of each cell center. The total area was approximately 523 000 km². For each cell we extracted, processed and transformed the predictors using the same procedure and parameters as for the transect data (table S3).

To avoid nuisance parameters, namely parameters with an overestimated contribution, and model selection uncertainty, we based the range-wide density prediction on qAICc-weighted multi-model inference (Burnham and Anderson 2002). Specifically, we first derived all possible models on the basis of the test predictors (5824 models). Six of those models did not converge, and we used the remaining 5818 models to derive a density prediction for each grid cell ($n = 620\,043$ cells) for the year 2015. These predictions were made in link space and weighted by the corresponding models’ qAICc, summed for each cell, and were finally exponentiated to produce chimpanzee densities (Cade 2015). We calculated 95% confidence intervals based on non-parametric bootstrapping ($n = 1\,000$) with the sampling units being the datasets (Manly 1997).

2.3. Identifying populations and low-connectivity areas

To estimate where connectivity between chimpanzee populations might be reduced, we first identified grid cells with a high likelihood of chimpanzee presence based on modeled chimpanzee density and expert opinion (details in supplementary material). We then determined patches of connected presence cells. Cells were iteratively assigned to the same patch when they were within a threshold distance or connected via cells separated by no more than the threshold distance. There is little information for dispersal distances between chimpanzee communities, for example when females transfer from their natal group. Published maximum daily travel distances range from 9 km in rainforest habitat (Herbinger *et al* 2001) to 16 km in drier habitat (Humble *et al* 2011), and may be larger in

very dry areas where chimpanzees have larger home ranges (Pruetz 2018). However, this is likely only the case within suitable habitat and in the absence of barriers such as areas densely populated by humans. As this is a broad-scale analysis, we did not account for conditions between presence cells. Therefore, we present three scenarios for possible dispersal distances, namely 5, 15, and 25 km, to identify areas where connectivity might be low or be reduced in the near future in case of land-use change or increase of other threats.

2.4. Spatial distribution of chimpanzees in relation to infrastructure

Large-scale land-use change across West Africa is mainly driven by the expansion of agricultural areas, resource extraction, and development of associated infrastructure (Norris *et al* 2010, Edwards *et al* 2014, Laurance *et al* 2015). While most of the land surface has essentially been divided into mining and timber concessions, as well as areas for renewable energy production, such as hydropower plants, spatial data are not available for the entirety of the western chimpanzee range. We therefore focused on only one of these planned development projects, namely proposed ‘development corridors’ (Laurance *et al* 2015), to illustrate how such developments could affect western chimpanzees if they were implemented. Development corridors center on the expansion of roads, railroads, pipelines, and ports, to improve the movement of people and goods between remote areas and urban centers. The aim is to enable rural communities’ access to markets and social services, and ultimately improve agricultural productivity, market integration, and regional trade (Mulenga 2013, Weng *et al* 2013, Laurance *et al* 2015). However, these infrastructure projects could lead to environmental damage by opening up formerly inaccessible areas and intersecting protected areas (Laurance *et al* 2015, Sloan *et al* 2017). Four corridors have been proposed for West Africa: Conakry-Buchanan (Guinea, Liberia, Sierra Leone), Dakar-Port Harcourt (Mali, Senegal), Gulf of Guinea (Côte d’Ivoire, Ghana, Liberia), and Sekondi/Ouagadougou (Burkina Faso, Ghana) (Laurance *et al* 2015). The recent \$22.7 Mio agreement between the Economic Community of West African States and the African Development Bank to upgrade roads on the ‘Gulf of Guinea’ corridor (African Development Bank 2019) and feasibility studies for the upgrade of the Dakar-Bamako railroad on the ‘Dakar-Port Harcourt’ corridor (PIDA 2018) suggest that these developments might threaten apes and their habitat (Laurance 2018). To estimate how many chimpanzees occur in proximity to these corridors, we overlaid the 50-km wide corridor bands from Laurance *et al* (2015) and Sloan *et al* (2017) with the modeled chimpanzee density distribution.

To provide further contextual information for conservation planning we determined the proportion

Table 1. Estimated western chimpanzee abundance by country (within geographic range delineated by IUCN SSC A.P.E.S. database).

Country	Estimated chimpanzee abundance (95% CI)	% chimpanzees living in national parks and IUCN category I or II protected areas
Guinea	33 139 (8796–68 203)	12.21
Liberia	6050 (2902–13 690)	14.22
Sierra Leone	5925 (1951–12 668)	31.20
Senegal	2642 (1077–13 293)	31.55
Guinea-Bissau	1908 (923–6121)	34.45 ^a
Mali	2029 (322–9228)	10.00
Côte d’Ivoire	1093 (329–3299)	49.52 ^b
Ghana	24 (1–212)	14.40
Total	52 811 (17 577–96 564)	17.03

^a As the spatial outline of Boé and Dulombi National Parks provided by the World Database of Protected Areas is not up to date (A Goedmakers pers. obs.), we used the outline provided by the ‘Instituto da Biodiversidade e das Áreas Protegidas’ (Agency of Guinea-Bissau government responsible for national parks) for this calculation.

^b It is noteworthy that while this number seems high, chimpanzees have declined by more than 90% across Côte d’Ivoire including regional extinctions resulting in a strong contraction of their range (Campbell *et al* 2008, Kühl *et al* 2017).

of chimpanzees in three habitat types based on the Global land cover dataset (Friedl *et al* 2010): forest (‘broadleaf forest’, ‘mixed forest’), savanna-mosaic (‘savanna’, ‘woody savanna’, ‘open shrubland’, ‘closed shrubland’), and cropland (‘cropland’, ‘cropland/natural vegetation mosaic’). We also determined the distance of each grid cell to the closest road (FAO 2005) and settlement (Esch *et al* 2012) to estimate how many chimpanzees live within 5 km and 10 km of roads and settlements. All analyses were implemented in R (vers. 3.4x, R Core Team 2018).

3. Results

3.1. Modeled chimpanzee density distribution

We estimated a total western chimpanzee abundance of 52 811 (95% confidence interval: 17 577–96 564), with the highest numbers in Guinea, Liberia and Sierra Leone (table 1). Densities ranged between <0.01 and 6.3 individuals km⁻². The highest densities were predicted for the Fouta Djallon highland region (figure 2). We estimated that 7.66% of western chimpanzees range in high-level protected areas (i.e. national parks and IUCN Cat I + II) as of 2015. Since then several new national parks have specifically been created for the protection of western chimpanzees, e.g. Boé and Dulombi (Guinea-Bissau), and Gola and Grebo-Krahn (Liberia), while Moyen Bafing (Guinea) is currently being created. Consequently, 8.79% of the current range is now a high-level protected area which




Figure 2. Modeled western chimpanzee density distribution (within geographic range delineated by IUCN SSC A.P.E.S. database, protected area outline from UNEP-WCMC and IUCN 2017, Boé and Dulombi National Parks outline from Instituto da Biodiversidade e das Áreas Protegidas in Guinea-Bissau). For plotting purposes we truncated densities at two because <0.04% of cells had estimated values larger than two, and otherwise low density cells would not be discernible. Maps of lower and upper confidence limit are in the supplementary material (figure S1).

corresponds to 17.03% of the estimated western chimpanzee population.

3.2. Population connectivity analysis

We estimated that there is one large chimpanzee population across the Fouta Djallon highland region and adjacent areas, extending from Senegal and Guinea-Bissau, across Guinea and Mali and into Sierra Leone (figure 3). This population comprises at least half of the remaining chimpanzees in West Africa (>33 000 individuals, details table S5). The southern population that extends from eastern Guinea across Liberia to Tai National Park in western Côte d'Ivoire comprises the remaining half of western chimpanzees (table S5). Our analysis revealed that connectivity between these two populations might be low in certain areas, specifically across the Upper Niger Basin in Guinea, and where the three countries Guinea, Liberia and Sierra Leone meet (green and blue patch in figure 3(a)), and in the Zone Forestière in southern Guinea (green and red patch in figure 3(b)), population estimates for all scenarios in table S5 and results for additional minimum density thresholds in figure S2).

3.3. Spatial distribution of chimpanzees in relation to infrastructure

We estimated that 10.44% of chimpanzees lived within 25 km of the four aforementioned development



corridors. The planned Dakar-Port Harcourt corridor in Senegal and Mali would intersect the northernmost distribution of western chimpanzees, while the Conakry-Buchanan corridor would intersect Outamba-Kilimi National Park and the above identified low-connectivity area at the tri-national border of Guinea, Liberia, and Sierra Leone (figure 4). Azagny and Banco National Park in Côte d'Ivoire, which still hold small chimpanzee populations, are entirely within 25 km of the proposed Gulf of Guinea corridor.


For further contextual information relevant for conservation planning, we estimated that 77.93% of western chimpanzees live in savanna-mosaic habitat, 16.38% in forest habitat, and 5.32% in cropland habitat. We also estimated that 38.59% of chimpanzees live within 5 km and 67.43% within 10 km of settlements, while 59.25% live within 5 km and 88.11% within 10 km of roads (figure 5).

4. Discussion

4.1. Implications for western chimpanzee conservation

As IUCN SSC A.P.E.S. provides access to a large number of ape survey datasets, many of which have not been published in peer-reviewed journals, we were able to compile a dataset representative of the





entire range of western chimpanzees. For the first time, we could thus model range-wide density distribution of this taxon and derive information on chimpanzee status important for conservation planning.


The analysis showed that two main populations can be distinguished (figure 3). Chimpanzees in the Fouta Djallon and adjacent areas appear to be a large population characterized by savanna-mosaic habitat. Despite challenging ecological conditions (Wessling *et al* 2018), chimpanzees persist here at high densities due to relatively low levels of anthropogenic threats and a high prevalence of hunting taboos (Boesch *et al* 2017, Heinicke *et al* 2019). Chimpanzees in the second largest population live in a habitat mostly characterized by rainforest. Our analysis indicated that these two populations might have low connectivity at present (figure 3). The other chimpanzee populations remaining in Côte d’Ivoire and Ghana are already completely isolated (Kühl *et al* 2017), and due to their small size these populations are particularly vulnerable to stochastic events, such as disease outbreaks, that can cause local extinctions (Knight *et al* 2016).

If implemented as planned, the multi-national infrastructure projects, including development corridors, hydropower plants and powerlines, could pose multiple threats to chimpanzees (Laurance 2018). First, infrastructure development is likely to incur direct loss of chimpanzee communities, because chimpanzees cannot shift their home range to move away from disturbances as they are highly territorial (Morgan *et al* 2018). Large-scale habitat loss can especially ensue from infrastructure developments that entail expansion of settlements and agriculture, one of the explicit objectives of development corridors (Laurance *et al* 2015). For example, large-scale deforestation and wide-spread hunting as a result of industrial agriculture led to strong declines of chimpanzee populations in Côte d’Ivoire and Ghana (Kormos *et al* 2003, Campbell *et al* 2008). Beyond the direct reduction in chimpanzee abundance this may also lead to loss in behavioral diversity, as even neighboring

communities can differ in their behavioral repertoire (Luncz *et al* 2012). Second, infrastructure expansions often carry secondary threats, especially an increase in hunting due to the arrival of more people, who potentially also do not adhere to established hunting taboos (Golden and Comaroff 2015). For example, in northern Congo the development of a logging concession with road construction and influx of workers led to a 64% increase in bushmeat supply (Poulsen *et al* 2009). However, a recent study of apes in Western Equatorial Africa underlined that this is context dependent, as central chimpanzee densities were significantly lower close to roads, but this effect disappeared with the presence of law enforcement (Strindberg *et al* 2018). Third, infrastructure projects lead to habitat fragmentation and act as dispersal barriers for a wide range of species (Laurance *et al* 2009). Although, chimpanzees have been observed to cross unpaved roads (Hockings *et al* 2015), genetic studies demonstrate that habitat fragmentation reduced chimpanzee population connectivity (Knight *et al* 2016, da Silva Borges 2017). A notable example is the case of Bossou, Guinea, where habitat fragmentation led to the isolation of a chimpanzee group, and there has been no female immigration from neighboring groups for the last 30 years (Matsuzawa *et al* 2011). What the impact on western chimpanzees will be remains to be seen and will strongly depend on the context. However, the list of potential negative impacts emphasizes the necessity to apply the mitigation hierarchy during the planning and construction of infrastructure (i.e. avoid, minimize, restore, and offset; BBOP 2013).

4.2. Comparison to previous estimates and data gaps

Overall, our abundance estimates are in line with previous studies that estimated around 7000 chimpanzees in Liberia (Tweh *et al* 2015), 5600 in Sierra Leone (Brncic *et al* 2015), and 17 700 in the Guinean part of the Fouta Djallon (WCF 2012). Kühl *et al* (2017) compiled abundance data for 35 sites across West Africa, covering 40% of the western chimpanzee range, and estimated a



minimum of 35 000 chimpanzees. Our predicted abundance for Senegal is notably higher than previous estimates of 200–400 which were extrapolated from small-scale surveys (Kormos *et al* 2003). Similarly, our estimate is higher than the 600–1000 estimated for Guinea-Bissau, but that estimate was not based on quantitative data (Kormos *et al* 2003).

Modeled estimates are associated with uncertainties (figure S1) due to the possibility of missing predictors, and differences in spatial scale of different predictor datasets (table S2). Uncertainties are more pronounced for areas with fewer data and those that differ strongly from surveyed areas regarding predictor space coverage. While substantial data are already available for large parts of the western chimpanzee's range, notable data gaps exist (figure 6). Identifying such gaps can inform where further surveys are needed, and shows where results are uncertain and need to be interpreted with care (see also figure S1). In particular, further field surveys are needed for Mali, for which only few data points are available. However, this is difficult due to the ongoing political instability there. Considering that our estimates are significantly higher than previous estimates for Senegal and Guinea-Bissau and have large confidence intervals, more intensive and representative future surveys in these countries would be instrumental in verifying our estimates. Similarly, high densities were predicted for the Kourandou and Simandou mountain ranges in eastern Guinea, but considering that this region is very dry, densities might have been over-estimated and further surveys would be needed to

validate these predictions. The aim of the model was thus to predict general patterns in chimpanzee density distribution across its range. However, its usefulness is limited at the local scale, for which site-based surveys are clearly superior.

4.3. Contribution of a taxon-specific database

Our study exemplifies the multi-faceted advantages of the IUCN SSC A.P.E.S. database. First, it can design data sharing policies reflecting the data sharing culture of its research field (Thessen and Patterson 2011), and by building trust and collaborations with a variety of data collectors, it can compile data in greater depth and provide access to unpublished datasets for a wider community. It also provides contextual information on the data, such as ecological and anthropogenic variables. Second, IUCN SSC A.P.E.S. not only ensures data attribution to data collectors so that their efforts are credited (Thessen and Patterson 2011), but it can also provide direct benefits to data collectors, for example, by collaborating on research projects and sharing authorship. In addition, database staff can support data collectors during study design and data analysis. Third, IUCN SSC A.P.E.S. hosts different types of data which can be used to derive different population parameters, including species abundance, density distribution, population trend, population connectivity, and habitat suitability (table 2). Fourth, databases can pool expertise and technical skills to process data to be directly fed into conservation decision-making. For example, data contained in

Table 2. Information on western chimpanzees that has been or could be derived from datasets archived in IUCN SSC A.P.E.S.

Type of analysis	Results for western chimpanzees	Example studies
Abundance	52 811 (17 577–96 564) individuals	This study
Climate change	To be investigated	
Density distribution	Figure 2 this study	This study
Ecosystem service provision through habitat protection (e.g. carbon storage, watershed protection, human cultural traditions associated with the habitat)	To be investigated	
Geographic range	524 100 km ²	Kühl <i>et al</i> 2017
IUCN Red List assessment	Critically Endangered	Humle <i>et al</i> 2016
Landscape-scale metrics	Figure 5 this study	This study
Land-use across species range	77.93% in savanna-mosaic, 16.38% in forest habitat, 5.32% in croplands	This study
Population trend	Annual decline of 5.96%, decline of 80.2% between 1990–2014	Kühl <i>et al</i> 2017
Population connectivity	Figure 3 this study	This study
Presence estimation for individual grid cells	Figure 3 this study	This study
Protected area coverage	17.03% of chimpanzees occur in high-level protected areas	This study
Species co-occurrence	To be investigated	
Suitable environmental conditions	Decline of area with suitable environmental conditions from 1990 to 2000	Junker <i>et al</i> 2012, Jantz <i>et al</i> 2016
Survey gaps	Figure 6 this study	This study

IUCN SSC A.P.E.S. had been used for a range of applications, including assessments for the IUCN Red List (e.g. Fruth *et al* 2016, Humle *et al* 2016, Maisels *et al* 2018), conservation action plans (e.g. IUCN and ICCN 2012, IUCN 2014), CITES, UNEP, and funding organizations (e.g. GRASP and IUCN 2018). Fifth, due to their knowledge of the field, database staff can also convey contacts to relevant experts and thereby improve information flow between different stakeholders, for example for the verification of study results. Thereby, research efforts can be streamlined and duplication avoided. Lastly, funders and journals increasingly require that research results are made publicly available. Accordingly, study results based on IUCN SSC A.P.E.S. data are also stored in the database (table 2). IUCN SSC A.P.E.S. thus enables a two-way information and knowledge exchange and functions as an intermediary to bridge the gap between stakeholders collecting data and those basing their research or decision-making on that data.

Taxon-specific databases already exist for different taxonomic groups and regions, for example the North American Breeding Bird Survey (Sauer *et al* 2013), African Elephant Database (Thouless *et al* 2016), Biological Records Centre in the UK (Pocock *et al* 2015), or Entomofauna Germanica (Bleich *et al* 2019). Despite their fundamental role in curating data relevant for conservation planners and policy makers, databases face severe funding shortages. We suggest that their contribution to filling data gaps for under-represented taxa and regions needs to be recognized, and that many more databases could be established. Similar to initiatives in the biomedical sciences that set up database networks, such as bioDBnet (Mudunuri *et al* 2009) and BioMart Central Portal (Guberman

et al 2011), a network of taxon-specific databases could be grown incrementally to complement global data repositories. Within such a network each database would pool data and expertise of the respective research field, while data users can retrieve data available across the entire network, for example, for a specific country or region. Taxon-specific databases can thus fill the niche between local data collectors and global data repositories.

Acknowledgments

We would like to thank the governments and national authorities for providing permission to collect data in chimpanzee range countries in West Africa. We also thank all organizations, their field staff and funders for providing chimpanzee survey data, specifically Chimbo Foundation, Conservation International, Conservation Society of Sierra Leone, Fauna and Flora International, Jane Goodall Institute Spain, Royal Society for the Protection of Birds, Pan African Programme, Society for the Conservation of Nature of Liberia, Tacugama Chimpanzee Sanctuary, West African Primate Conservation Action, Wild Chimpanzee Foundation, and World Wide Fund for Nature. We thank Manuel Llana, Amanda Barciela, Irene Gutiérrez and Paula Álvarez for support in the field. We further thank Emmanuelle Normand and Elleni Vendras for providing chimpanzee survey data for Côte d'Ivoire and Ghana, respectively. We thank Thomas Esch for providing Global Urban Footprint data, and William Laurance and Sean Sloan for providing spatial data for the development corridors. We thank Christian Krause for help with implementation on the high-performance cluster. We would also like to thank the participants of the Guinean

National Chimpanzee Conservation Action Planning workshop in Conakry and the Regional Chimpanzee Conservation Action Planning workshop in Monrovia in 2017 for helpful feedback. We thank two anonymous reviewers for helpful comments on earlier drafts of this manuscript. Author GC was affiliated with the Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany during the time of research and is now employed by the company The Biodiversity Consultancy Ltd, Cambridge, UK, which had no influence on the research conducted. We would like to thank the Max Planck Society and the Robert Bosch Foundation for funding and support. The raw data and the modeled density distribution for western chimpanzees will be made available via the IUCN SSC A.P.E.S. database (<http://apesportal.eva.mpg.de>).

ORCID iDs

Stefanie Heinicke  <https://orcid.org/0000-0003-0222-5281>

Emmanuel Danquah  <https://orcid.org/0000-0002-8305-5706>

Erin G Wessling  <https://orcid.org/0000-0001-9661-4354>

Hjalmar S Kühl  <https://orcid.org/0000-0002-4440-9161>

References

- African Development Bank 2019 African Development Bank, ECOWAS, sign agreement for study on Abidjan-Lagos Corridor Highway *Afr. Dev. Bank* (<https://afdb.org/en/news-and-events/african-development-bank-ecowas-sign-agreement-for-study-on-abidjan-lagos-corridor-highway-18962/>)
- Baayen RH 2008 *Analyzing Linguistic Data: A Practical Introduction to Statistics Using R* (Cambridge: Cambridge University Press)
- BBOP 2013 *To No Net Loss and Beyond: An Overview of the Business and Biodiversity Offsets Programme (BBOP)* (Washington, DC: BBOP) (<https://www.forest-trends.org/publications/to-no-net-loss-and-beyond/>)
- Bleich O, Gürlich S and Köhler F 2019 Verzeichnis und Verbreitungsatlas der Käfer Deutschlands (<http://coleokat.de>)
- Boakes EH, McGowan P J K, Fuller R A, Chang-qing D, Clark N E, O'Connor K and Mace G M 2010 Distorted views of biodiversity: spatial and temporal bias in species occurrence data *PLOS Biol.* **8** e1000385
- Boesch L, Mundry R, Kühl H S and Berger R 2017 Wild mammals as economic goods and implications for their conservation *Ecol. Soc.* **22** 36
- Bronic T, Amarasekaran B, McKenna A, Mundry R and Kühl H S 2015 Large mammal diversity and their conservation in the human-dominated land-use mosaic of Sierra Leone *Biodivers. Conserv.* **24** 2417–38
- Burnham K P and Anderson D R 2002 *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* 2nd edn (New York: Springer)
- Cade B S 2015 Model averaging and muddled multimodel inferences *Ecology* **96** 2370–82
- Campbell G, Kuehl H, N'Goran Kouamé P and Boesch C 2008 Alarming decline of West African chimpanzees in Côte d'Ivoire *Curr. Biol.* **18** R903–4
- Collen B, Loh J, Whitmee S, McRae L, Amin R and Baillie J E M 2009 Monitoring change in vertebrate abundance: the living planet index *Conserv. Biol.* **23** 317–27
- Corlett R T 2011 Trouble with the gray literature *Biotropica* **43** 3–5
- Costello M J, Michener W K, Gahegan M, Zhang Z-Q and Bourne P E 2013 Biodiversity data should be published, cited, and peer reviewed *Trends Ecol. Evol.* **28** 454–61
- da Silva Borges F F 2017 *A Country-Level Genetic Survey of the IUCN Critically Endangered Western Chimpanzee (*Pan troglodytes verus*) in Guinea-Bissau Master Thesis* (Porto: University of Porto)
- Edwards D P, Sloan S, Weng L, Dirks P, Sayer J and Laurance W F 2014 Mining and the African environment *Conserv. Lett.* **7** 302–11
- Esch T, Taubenböck H, Roth A, Heldens W, Felbier A, Thiel M, Schmidt M, Müller A and Dech S 2012 TanDEM-X mission—new perspectives for the inventory and monitoring of global settlement patterns *J. Appl. Remote Sens.* **6** 061702–1
- FAO 2005 Roads of the World (Vmap0) (<http://www.fao.org/geonetwork>)
- Friedl M A, Sulla-Menashe D, Tan B, Schneider A, Ramankutty N, Sibley A and Huang X 2010 *Global land cover MCD12Q1, Collection 5.1, IGBP Land Cover* (Boston, MA: Boston University) (<http://landcover.org/data/lc/>)
- Fruth B 2016 *Pan paniscus*. The IUCN Red List of Threatened Species (<http://iucnredlist.org>)
- GBIF 2018 GBIF Home Page (<https://gbif.org/>)
- Golden C and Comaroff J 2015 Effects of social change on wildlife consumption taboos in northeastern Madagascar *Ecol. Soc.* **20** 41
- GRASP and IUCN 2018 *Report to the CITES Standing Committee on the Status of Great Apes* United Nations Environment Programme Great Apes Survival Partnership, Nairobi, and International Union for Conservation of Nature, Gland
- Guberman J M *et al* 2011 BioMart central portal: an open database network for the biological community *Database* **2011** bar041
- Heinicke S *et al* 2019 Characteristics of positive deviants in western chimpanzee populations *Front. Ecol. Evol.* (<https://doi.org/10.3389/fevo.2019.00016>)
- Herbinger I, Boesch C and Rothe H 2001 Territory characteristics among three neighboring chimpanzee communities in the Taï National Park, Côte d'Ivoire *Int. J. Primatol.* **22** 143–67
- Hilbe J M 2011 *Negative binomial regression* (Cambridge: Cambridge University Press)
- Hockings K J *et al* 2015 Apes in the anthropocene: flexibility and survival *Trends Ecol. Evol.* **30** 215–22
- Humle T, Boesch C, Campbell G, Junker J, Koops K, Kühl H and Sop T 2016 *Pan troglodytes ssp. verus*. The IUCN Red List of Threatened Species 2016 (<http://iucnredlist.org>)
- Humle T, Colin C, Laurans M and Raballand E 2011 Group release of sanctuary chimpanzees (*Pan troglodytes*) in the Haut Niger National Park, Guinea, West Africa: ranging patterns and lessons so far *Int. J. Primatol.* **32** 456–73
- IUCN 2014 *Regional Action Plan for the Conservation of Western Lowland Gorillas and Central Chimpanzees 2015–2025* (Gland: IUCN SSC Primate Specialist Group)
- IUCN and ICCN 2012 *Bonobo (*Pan paniscus*): Conservation Strategy 2012–2022* (Gland: IUCN/SSC Primate Specialist Group & Institut Congolais pour la Conservation de la Nature)
- Jantz S M, Pintea L, Nackoney J and Hansen M C 2016 Landsat ETM plus and SRTM data provide near real-time monitoring of chimpanzee (*Pan troglodytes*) habitats in Africa *Remote Sens.* **8** 427
- Jetz W, McPherson J M and Guralnick R P 2012 Integrating biodiversity distribution knowledge: toward a global map of life *Trends Ecol. Evol.* **27** 151–9
- Junker J *et al* 2012 Recent decline in suitable environmental conditions for African great apes *Divers. Distrib.* **18** 1077–91
- Kindsvater H K, Dulvy N K, Horswill C, Juan-Jorda M-J, Mangel M and Matthiopoulos J 2018 Overcoming the data crisis in biodiversity conservation *Trends Ecol. Evol.* **33** 676–88

- Knight A, Chapman H M and Hale M 2016 Habitat fragmentation and its implications for endangered chimpanzee *Pan troglodytes* *Conservation Oryx* **50** 533–6
- Kormos R and Boesch C 2003 *Regional Action Plan for the Conservation of Chimpanzees in West Africa*. (Washington DC: IUCN/SSC Primate Specialist Group and Conservation International)
- Kormos R, Boesch C, Bakarr M I and Butynski T 2003 *West African Chimpanzees. Status survey and Conservation Action Plan* (Gland and Cambridge: IUCN/SSC Primate Specialist Group)
- Kühl H S *et al* 2017 The Critically Endangered western chimpanzee declines by 80% *Am. J. Primatol.* **79** e22681
- Kühl H S, Williamson L, Sanz C, Morgan D and Boesch C 2007 A.P.E.S., IUCN SSC A.P.E.S. database (<http://apesportal.eva.mpg.de/>)
- Laurance W F 2018 Towards more sustainable infrastructure: challenges and opportunities in ape range states of Africa and Asia *State of the APES: Infrastructure Development and Ape Conservation*, ed. Arcus Foundation (Cambridge: Cambridge University Press)
- Laurance W F, Goosem M and Laurance S G W 2009 Impacts of roads and linear clearings on tropical forests *Trends Ecol. Evol.* **24** 659–69
- Laurance W F, Sloan S, Weng L and Sayer J A 2015 Estimating the environmental costs of Africa's massive 'development corridors' *Curr. Biol.* **25** 3202–8
- Luncz L V, Mundry R and Boesch C 2012 Evidence for cultural differences between neighboring chimpanzee communities *Curr. Biol.* **22** 922–6
- Maisels F, Bergl R A and Williamson E A 2018 *Gorilla gorilla*. The IUCN Red List of Threatened Species 2018 (<http://iucnredlist.org>)
- Manly B F J 1997 *Randomization, Bootstrap and Monte Carlo Methods in Biology* 2nd edn (Boca Raton, FL: CRC Press)
- Matsuzawa T, Ohashi G, Humle T, Granier N, Kourouma M and Soumah A G 2011 Green corridor project: planting trees in the savanna between Bossou and Nimba *The Chimpanzees of Bossou and Nimba Primatology Monographs* ed T Matsuzawa, T Humle and Y Sugiyama (Tokyo: Springer)
- Morgan D, Mundry R, Sanz C, Ayina C E, Strindberg S, Lonsdorf E and Kühl H S 2018 African apes coexisting with logging: comparing chimpanzee (*Pan troglodytes troglodytes*) and gorilla (*Gorilla gorilla gorilla*) resource needs and responses to forestry activities *Biol. Conserv.* **218** 277–86
- Mudunuri U, Che A, Yi M and Stephens R M 2009 bioDBnet: the biological database network *Bioinformatics* **25** 555–6
- Mulenga G 2013 *Developing Economic Corridors in Africa: Rationale for the Participation of the African Development Bank* (Tunis: African Development Bank)
- Murai M, Ruffler H, Berlemont A, Campbell G, Esono F, Agbor A, Mbomio D, Ebana A, Nze A and Kühl H S 2013 Priority areas for large mammal conservation in equatorial Guinea *PLOS One* **8** e75024
- Norris K, Asase A, Collen B, Gockowski J, Mason J, Phalan B and Wade A 2010 Biodiversity in a forest-agriculture mosaic—the changing face of West African rainforests *Biol. Conserv.* **143** 2341–50
- Peterson A T and Soberón J 2018 Essential biodiversity variables are not global *Biodivers. Conserv.* **27** 1277–88
- PIDA 2018 Dakar-Bamako Standard Gauge Railway, Virtual PIDA Information Centre, Programme for Infrastructure Development in Africa (<http://au-pida.org/view-project/939/>)
- Pocock M J O, Roy H E, Preston C D and Roy D B 2015 The biological records centre: a pioneer of citizen science *Biol. J. Linn. Soc.* **115** 475–93
- Poulsen J R, Clark C J, Mavah G and Elkan P W 2009 Bushmeat supply and consumption in a tropical logging concession in northern Congo *Conserv. Biol.* **23** 1597–608
- Pruetz J D 2018 Nocturnal behavior by a diurnal ape, the West African chimpanzee (*Pan troglodytes verus*), in a savanna environment at Fongoli, Senegal *Am. J. Phys. Anthropol.* **166** 541–8
- R Core Team 2018 *R: A Language and Environment for Statistical Computing* (Vienna: R Foundation for Statistical Computing) (<https://R-project.org/>)
- Reichman O J, Jones M B and Schildhauer M P 2011 Challenges and opportunities of open data in ecology *Science* **331** 703–5
- Sauer J, Link W, Fallon J, Pardieck K and Ziolkowski D 2013 The North American breeding bird survey 1966–2011 *North Am. Fauna* **79** 1–32
- Schwartz M W, Cook C N, Pressey R L, Pullin A S, Runge M C, Salafsky N, Sutherland W J and Williamson M A 2018 Decision support frameworks and tools for conservation *Conserv. Lett.* **11** e12385
- Sloan S, Bertzky B and Laurance W F 2017 African development corridors intersect key protected areas *Afr. J. Ecol.* **55** 731–7
- Strindberg S *et al* 2018 Guns, germs, and trees determine density and distribution of gorillas and chimpanzees in Western Equatorial Africa *Sci. Adv.* **4** eaar2964
- Thessen A E and Patterson D J 2011 Data issues in the life sciences *ZooKeys* **150** 15–51
- Thouless C R, Dublin H T, Blanc J J, Skinner D P, Daniel T E, Taylor R D, Maisels F, Frederick H L and Bouché P 2016 *African elephant status report 2016. An Update from the African Elephant Database. Occasional Paper of the IUCN Species Survival Commission No. 60.* IUCN/SSC African Elephant Specialist Group (Gland, Switzerland: IUCN)
- Tweh C G, Lormie M M, Kouakou C Y, Hillers A, Kühl H S and Junker J 2015 Conservation status of chimpanzees *Pan troglodytes verus* and other large mammals in Liberia: a nationwide survey *Oryx* **49** 710–8
- UNEP-WCMC and IUCN 2017 *Protected Planet: The World Database on Protected Areas (WDPA), Online Jan 2017* (Cambridge: UNEP-WCMC and IUCN) (www.protectedplanet.net)
- Voigt M *et al* 2018 Global demand for natural resources eliminated more than 100 000 bornean orangutans *Curr. Biol.* **28** 761–9
- WCF 2012 *Etat de la faune et des menaces dans les aires protégées terrestres et principales zones de forte biodiversité de Rep. de Guinée* (Sangaredi, Republic of Guinea: Wild Chimpanzee Foundation Sangaredi, Republic of Guinea) (<http://www.wildchimps.org>)
- Weng L, Boedihartono A K, Dirks P H G M, Dixon J, Lubis M I and Sayer J A 2013 Mineral industries, growth corridors and agricultural development in Africa *Glob. Food Secur.* **2** 195–202
- Wessling E G, Deschner T, Mundry R, Pruetz J D, Wittig R M and Kühl H S 2018 Seasonal variation in physiology challenges the notion of chimpanzees (*Pan troglodytes verus*) as a forest-adapted species *Front. Ecol. Evol.* **6** 60
- Wich S A and Marshall A J 2016 *An Introduction to Primate Conservation* (Oxford: Oxford University Press)
- Wich S A *et al* 2016 Land-cover changes predict steep declines for the Sumatran orangutan (*Pongo abelii*) *Sci. Adv.* **2** e1500789