

1 **EFFICACY TESTING OF AN IMMERSION VACCINE AGAINST *AEROMONAS***  
2 ***SALMONICIDA* AND IMMUNOCOMPETENCE IN BALLAN WRASSE (*LABRUS***  
3 ***BERGYLTA*, *ASCANIUS*)**

4 Athina Papadopoulou<sup>1,2</sup>, Sean J. Monaghan<sup>1</sup>, Nicola Bagwell<sup>2</sup>, Mickael Teixeira Alves<sup>2</sup>, David  
5 Verner-Jeffreys<sup>2</sup>, Tim Wallis<sup>3</sup>, Andrew Davie<sup>1,a</sup>, Alexandra Adams<sup>1</sup>, Herve Migaud<sup>1\*</sup>

6

7 <sup>1</sup>Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA,  
8 UK

9 <sup>2</sup>Centre for Environment, Fisheries and Aquaculture Science, Barrack Road, The Nothe,  
10 Weymouth, Dorset, DT4 8UB, UK

11 <sup>3</sup>Ridgeway Biologicals Ltd. a Ceva Santé Animale company, Units 1-3 Old Station Business  
12 Park, Compton, Berkshire RG20 6NE UK

13 <sup>a</sup>Current address: Aquascot Ltd, Fyrish Way, Alness, Ross-shire, IV17 0PJ, UK.

14 Correspondence

15 Herve Migaud, Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling,  
16 Stirling FK9 4LA, UK

17 Email: [herve.migaud@stir.ac.uk](mailto:herve.migaud@stir.ac.uk)

18

19 **Abstract**

20 The development of effective vaccines is a critical step towards the domestication of emerging  
21 fish species for aquaculture. However, traditional vaccine delivery through intraperitoneal (i.p.)  
22 injection requires fish to reach a minimum size and age and therefore cannot provide  
23 protection at early developmental stages when infection may occur. This study investigated  
24 the effectiveness of immersion vaccination with respect to immunocompetence in a cleaner  
25 fish species (ballan wrasse, *Labrus bergylta*, *Ascanius*) used in Atlantic salmon farming as an  
26 alternative means to control sea lice. The species is susceptible to atypical strains of  
27 *Aeromonas salmonicida* (aAs) at early life stages (<15 g), when i.p. vaccination is not  
28 applicable. While immersion vaccination is currently used in commercial hatcheries, the  
29 optimal fish size for vaccination, and efficacy of the vaccine delivered by this route has not yet  
30 been established. Importantly, efficacy depends on the capability of the species immune

31 system to recognise antigens and process antigens to trigger and produce an adaptive  
32 immune response, (process known as immunocompetence). In this study, the efficacy of a  
33 polyvalent autogenous vaccine administered by immersion in juvenile ballan wrasse and the  
34 subsequent immune response induced was investigated after prime and booster vaccination  
35 regimes. In addition, temporal expression (0 - 150 days post hatch) of adaptive immune genes  
36 including major histocompatibility complex (MHC II CD74 molecule) and immunoglobulin M  
37 (IgM) was assessed using quantitative PCR (qPCR). Prime and/or boost vaccination by  
38 immersion of juvenile ballan wrasse (0.5 g and 1.5 g corresponding to 80 and 170 days post  
39 hatch (dph), respectively) did not provide significant protection against aAs *vapA* V after bath  
40 challenge under experimental conditions. Despite no evident protection > 80 dph, MHC II and  
41 IgM transcripts were first reported at 35 and 75 dph, respectively, suggesting a window of  
42 immunocompetence. The results provide important new information on the onset of adaptive  
43 immunity in ballan wrasse and highlight that immersion vaccination in the species for  
44 protection against aAs should be performed at later developmental stages (>1.5 g) in the  
45 hatchery.

46 **Key words:** ballan wrasse, immersion vaccination, *Aeromonas salmonicida*, adaptive  
47 immunity, IgM, MCH II.

## 48 1. Introduction

49 Atypical strains of *Aeromonas salmonicida* (aAs) are currently the most significant cause of  
50 mortalities in hatchery and post cage deployment of farmed ballan wrasse (*Labrus bergylta*,  
51 *Ascanius*) [1]. The bacterium is highly heterogenous infecting a variety of fresh and marine  
52 fish species and host specificity of strains has been reported [2]. Recently, partial sequencing  
53 of the paracrystalline surface array protein gene (*vapA* gene) has successfully been used to  
54 distinguish typical and atypical *As* strains to 23 *vapA* types [3, 4]. Ballan wrasse is susceptible  
55 to aAs *vapA* type V and VI in Scotland and Norway, respectively [4]. The successful  
56 development and application of vaccines to regulate bacterial challenges is increasingly  
57 important for ballan wrasse, as an emerging species, however such developments are  
58 hampered by the lack of basic understanding of immunity and immunocompetence in the  
59 species [5].

60 Vaccination is the most common disease prevention strategy in the finfish aquaculture industry  
61 [6]. Vaccines can be administered by either injection (intraperitoneally (i.p.) or intramuscularly  
62 (i.m.), immersion (dipping or bathing) and in some cases orally in feed [7]. Each practise has  
63 advantages and disadvantages. For instance, i.p. injection is the most commonly used method  
64 for juvenile fish (>15 g) generating greater protection than immersion especially facilitated by

65 including effective adjuvants, and provides a platform for standardised dosing of vaccine for  
66 each individual fish [6, 8]. However, immersion is the preferred method for small fish (0.5 – 25  
67 g) as it is less invasive, therefore minimising handling stress, is often more affordable and less  
68 labour intensive [6, 8]. A combination of these practises is usually applied in marine fish which  
69 are immunised initially by immersion (prime and booster may be required) at an early life stage  
70 and then i.p. injection vaccinated before being deployed to sea cages for on growing or, in the  
71 case of cleaner fish, for sea lice control [9].

72 Following immersion vaccination, immune related cells in the mucosa-associated lymphoid  
73 tissue (MALT) uptake the antigen [10]. Several studies have investigated the efficacy of  
74 immersion vaccination and antigen uptake however our understanding of the mucosal immune  
75 response in fish still remains limited [11]. A prerequisite for successful vaccination is a fully  
76 developed functional immune system able to develop immunological memory. The ontogeny  
77 of the immune system has been studied in various teleost fish species, showing species  
78 specific windows ranging from 8 days pre-hatching in rainbow trout (*Oncorhynchus mykiss*,  
79 L.) up to 5 and 10 month post hatch for carp (*Cyprinus carpio*, L.) and zebrafish (*Danio rerio*,  
80 Hamilton), respectively [12]. The anterior kidney and thymus are considered to be fully  
81 developed prior to hatching in rainbow trout (*Oncorhynchus mykiss*, L.) and Atlantic salmon  
82 [13], while immunocompetence (presence of IgM) occurs later during alevin development and  
83 is size dependant (fish of 20 – 30 mm in length) [14, 15]. In marine fish species, like sea bass  
84 (*Dicentrarchus labrax*, L.), spotted wolffish (*Anarhichas minor*, L.) and Atlantic cod (*Gadus*  
85 *morhua*, L.) studies have shown IgM positive lymphocytes between 1 and 10 weeks after  
86 hatching [15]. There are three main immunoglobulin (Igs) isotypes in teleost fish including IgM,  
87 IgD and IgT. Expression of functional Igs gives an indication of immune competence this would  
88 facilitate the identification of an immersion vaccination window prior to i.p. injection and after  
89 immunocompetence has been achieved [16]. IgM is the first antibody produced in teleost fish  
90 and it consistently appears at the mucosal sites in contrast with IgT that is not found in every  
91 fish species [17]. Thus, IgM is broadly used as a marker for immune competence. Another  
92 important component of the adaptive immune system in fish is the major histocompatibility  
93 complex (MHC) which is responsible for the recognition and presentation of antigen epitopes  
94 on T cell receptors (TCRs) [18]. The MHC genes produce two membrane glycoproteins class  
95 I and II [19]. MHC II express glycoproteins which are present only in antigen presenting cells  
96 (APCs) such as B cells and trigger the production of specific antibodies against exogenous  
97 antigens [20]. The binding with T helper cells (Th) activates B cells and memory cells  
98 differentiation [20]. Literature on ballan wrasse immunity is very limited and the ontogeny of  
99 the adaptive immune response has not been studied to date. Haugland, et al. [21] previously  
100 characterised the cellular innate response, and Bilal, et al. [22], [23] have reported adaptive

101 immune responses with regards to the presence of IgM, T cell receptors (TCRs) and somatic  
102 mutations of lymphocytes. In addition, the relative expression of antibodies (IgM, IgT and IgD)  
103 and TCR genes has been studied in the gut of ballan wrasse [24].

104 In an attempt to address disease challenges due to aAs (*vapA* type V), juvenile ballan wrasse  
105 in commercial Scottish hatcheries have typically been bath vaccinated at ca. 0.5 g, followed  
106 by booster vaccination at 1.0 – 1.5 g (1:10 dilution for 3 – 4 h) and then administered an i.p.  
107 injection booster between 10 and 15 g with a polyvalent autogenous vaccine (Ridgeway  
108 Biologicals, DDT Germany). In recent experimental trials, the injectable version of this  
109 commercial vaccine offered high protection to ballan wrasse (25 – 50 g) when fish were  
110 challenged with heterologous and autologous strains of aAs *vapA* type V (RPS 79 % and 91  
111 %, respectively) [25]. However, it became apparent that the species can encounter and  
112 succumb to bacterial disease in commercial hatcheries at earlier developmental stages (prior  
113 to i.p. vaccination) [26] and immunisation at earlier developmental stages is therefore needed.  
114 The efficacy of the autogenous immersion vaccine currently used in wrasse hatcheries has  
115 not been assessed and the size at which fish can be successfully vaccinated by immersion  
116 remains to be determined.

117 The aims of this study were to 1) assess the efficacy of the immersion autogenous vaccine  
118 vaccination against aAs to evaluate if the current commercial strategy of bath vaccination was  
119 effective for the autogenous vaccine and 2) to determine the ontogeny of the farmed ballan  
120 wrasse adaptive immune system using temporal gene expression analysis of immune genes  
121 for antigen presentation (MHC II) and antibody production (IgM). Results will ultimately help  
122 to optimise the vaccination strategy for farmed juvenile ballan wrasse.

## 123 **2. Materials and Methods**

### 124 **2.1. Immersion vaccination and challenge trial**

#### 125 **2.1.1. Fish acclimation**

126 Two batches of juvenile ballan wrasse ( $0.3 \pm 0.15$  g at 60 days post hatch (dph) and  $1.0 \pm 0.2$   
127 g at 150 dph) produced from eggs collected from a common broodstock population were  
128 obtained from a joint venture hatchery owned by MOWI (Scotland, UK) and Scottish Sea  
129 Farms and transported to the Centre for Environment, Fisheries and Aquaculture Science  
130 (Cefas, Weymouth UK). Fish were kept in aerated aquaria (150 L) during an acclimation period  
131 of three weeks at a constant temperature of  $15 \pm 1$  °C and photoperiod of 20 h light: 4 h dark.  
132 Water flow was set at 4.0 L / min and dissolved oxygen (DO) maintained at  $8.0 \pm 0.5$  mg / L.  
133 Fish were fed with Otohime Fish Diet B2 & C1 (Biokyowa, Japan) at 7 – 8 % of the body weight  
134 for 7 h / day using automatic feeders.

135 **2.1.2. Fish health assessment**

136 The health status of the fish was assessed prior to transportation as described in a previous  
137 study [25]. Briefly, whole larvae were screened for pathogens including common isolated  
138 bacteria (aAs and Vibrionaceae) as well as amoebic gill disease (AGD) by histology,  
139 bacteriology and molecular techniques [27-29]. In addition, fish health was assessed upon  
140 arrival to Cefas facilities for biosecurity and precautionous experimental measures and to confirm  
141 that fish were not carrying any notifiable viral diseases as listed in the OIE manual of diagnostic  
142 tests for aquatic animals [30]. Furthermore, whole fish samples were fixed in 10% neutral  
143 buffered formalin to confirm by histology that fish did not have any pathology prior to challenge.

144 **2.1.3. Immersion vaccination and sampling**

145 The two fish size cohorts post acclimation (small (S) - 0.5 g  $\pm$  0.2 g at 80 dph and large (L) -  
146 1.5  $\pm$  0.4 g at 170 dph) herein designated as small and large, were bath vaccinated with a  
147 formalin inactivated multivalent autogenous vaccine provided by Ridgeway Biologicals Ltd  
148 (IDT Biologika, Germany) in static sea water for 3 h at 15°C. The isolates contained in the  
149 vaccine were recovered from diseased ballan wrasse sampled at commercial hatcheries in  
150 Scotland and characterised as part of a previous study [26]. The vaccine used contained 9  
151 bacterial strains, 4 of which were aAs (3 strains of *vapA* type V; and 1 strain of *vapA* type VI;) and  
152 5 isolates belonging to the *Vibrionaceae* family (Table 1). The vaccine was diluted 1:100  
153 and the final vaccination doses are given in Supplementary Table 1. Mock vaccinated fish  
154 were exposed to sterile sea water (33 ppt). Tissues (liver (L), spleen (S) and head kidney  
155 (HK)) were sampled from vaccinated and mock vaccinated groups (n= 6 per group) at pre-  
156 vaccination (0 h) and 24 h post prime and booster vaccination. Samples were preserved in  
157 RNA stabilisation buffer [31] for later immune genes expression analysis.

158

**Table 1.** Bacterial isolates in the autogenous immersion vaccine.

Isolate ID	Bacterial species	Year of isolation	<i>vapA</i> type and /or isolate #
TW3/14	<i>Aeromonas salmonicida</i>	2014	V
TW4/14	<i>Aeromonas salmonicida</i>	2014	V
TW187/14	<i>Aeromonas salmonicida</i>	2014	V
TW184/16*	<i>Aeromonas salmonicida</i>	2016	VI
TW242/16	<i>Aliivibrio logei</i>	2016	Isolate 1
TW322/16	<i>Aliivibrio salmonicida</i>	2016	Isolate 1

TW130/16	<i>Vibrio splendidus</i>	2016	Isolate 1
TW319/16	<i>Vibrio ichthyenteri</i>	2016	Isolate 1
TW138/16	<i>Photobacterium indicum</i>	2016	Isolate 1

(\*) isolated from lumpsucker.

159

#### 160 **2.1.4. Bacteria preparation for bath challenge**

161 From previous studies, the aAs *vapA* type V strain (TW 4/14) isolated from infected farmed  
 162 ballan wrasse in Scotland and used in the formulation of the vaccine, caused mortalities > 60  
 163 % in fish exposed by bath and i.p. injection under experimental conditions [25, 32]. This  
 164 homologous strain was used for the bath challenge to assess the efficacy of the immersion  
 165 vaccine and it was grown in Tryptone Soya Broth (TSB) at 22°C for 48 h with continuous  
 166 shaking at 150 rpm. The bacteria were centrifuged at 2,000 x g for 10 min, bacterial pellets  
 167 were then washed with phosphate buffered saline (PBS) and re – suspended in sterile sea  
 168 water to an OD<sub>550</sub> 0.8. Colony forming unit (CFU) per mL were counted by the 10 – fold dilution  
 169 method.

#### 170 **2.1.5. Bath challenge and sampling**

171 Fish were challenged as described in an established challenge model for the species [32].  
 172 Briefly, all groups were starved for 24 h prior to challenge. Triplicate groups of vaccinated and  
 173 mock vaccinated fish were bath challenged with aAs *vapA* type V strain (TW 4 / 14) after 654-  
 174 degree days (DD) post prime immunisation and after 495 DD post booster in small aerated  
 175 aquaria of 5 L for the small cohort and 8 L for the large cohort (50 fish / aquarium). The bacteria  
 176 were adjusted to an OD<sub>550</sub> 0.8 (10<sup>8</sup> - 10<sup>9</sup> CFU / mL in 5 or 8 L of static sea water) and fish were  
 177 challenged for 4 h at 15 °C, as shown in Supplementary Figure 1. Control groups were  
 178 exposed to sterile sea water. The fish were then allocated to 30 L holding aquaria each with  
 179 water flows of 0.5 L / min, while all other parameters remained the same as outlined prior. The  
 180 fish were monitored for up to 30 days post infection (dpi). Ballan wrasse with obvious signs of  
 181 disease were considered as moribund and were sacrificed using an overdose of MS-222 (40  
 182 ppm; Tricaine methane sulphonate, Sigma). Head kidney swabs were sampled from moribund  
 183 fish (n= 5 per tank) and plated onto tryptone soya agar (TSA) to confirm specific mortality due  
 184 to aAs. Suspected aAs colonies were confirmed with the agglutination test (Mono-AQUA  
 185 BIONOR™, BIONOR). In addition, liver, spleen and head kidney tissue samples (n= 299; 100  
 186 from L, 99 from S and 100 from HK) from all moribund fish were fixed in 100% ethanol for  
 187 bacterial species / type confirmation with previously published conventional PCRs and qPCR  
 188 as described below [33, 34]. Tissues samples (L, S and HK; n= 6 per treatment for vaccinated,

189 mock vaccinated and control groups) were fixed in RNA stabilisation buffer after termination  
 190 of the challenge trial for gene expression analysis.

191 **2.1.6. DNA extraction and quality check**

192 DNA was extracted from preserved tissues using a salt precipitation method as described in  
 193 Khanam, et al. [35]. The extracted DNA was re-suspended in 5 mM Tris, concentration  
 194 measured and standardised at 100 ng /  $\mu$ L using NanoDrop® ND-1000 (Thermo Fisher  
 195 Scientific). Sample integrity was validated by gel electrophoresis. The samples were stored at  
 196 - 20°C until use.

197 **2.1.7. Conventional PCRs and Quantitative PCR (qPCR) for *Aeromonas salmonicida***

198 The presence of bacterial DNA was assessed on the V3-V4 hypervariable region of the 16S  
 199 rRNA gene – 16S PCR; if samples were negative no further testing was carried out as bacterial  
 200 load in the samples was considered below the assay detection limit. If positive, samples were  
 201 then screened with a species-specific PCR (*vapA*) for presence of *As* DNA using a previously  
 202 published PCR protocol targeting the hypervariable region of *vapA* gene – *vapA* PCR [4]. An  
 203 a*As* *vapA* type V specific PCR (a*As* type V specific PCR) previously developed by our group  
 204 [36] was used to determine the presence or absence of a*As* *vapA* type V. The primer pairs  
 205 used for the tissue samples screening with conventional PCRs are listed in Table 2. The  
 206 relative molecular weight of the amplicons was compared against a 100 bp gene ruler (Thermo  
 207 Scientific) on 1 % agarose gel. *Aeromonas salmonicida* bacterial loads were assessed on  
 208 tissue samples that were confirmed positive for *As* DNA with the conventional PCRs using a  
 209 modified protocol [34] as described in [36]. The primers and qPCR conditions are listed in  
 210 Table 1. Any samples above 35 Cross point were considered negative.

211 **Table 2.** Primers used for amplification of broad-range bacterial DNA, *Aeromonas salmonicida*  
 212 and atypical *Aeromonas salmonicida* *vapA* type V specific with conventional PCR and  
 213 quantitative PCR.

Gene	Oligo sequence	Annealing (°C)	Product size (bp)	Publication
<b>Bacterial rRNA 16S</b>	F341: CCTACGGGNGGCWGCAG R805: GACTACHVGGGTATCTAATCC	54.0	465	[37]
<b><i>vapA</i> partial</b>	F2: CTGGACTIONTCTCCACTGCTCA R3: ACGTTGGTAATCGCGAAATC	53.0	626	[4]
<b><i>vapA</i> partial</b>	Vspec-F: CAACGGTTTCTGGAGTAATAACTTT Vspec - R: TGCATCAGCAACAGCGGTAGT	57.0	254	[33]

---

*vapA*

F: ACTGTCTGTTACCCTGCCA  
R: GCTACTTCACCCTGATTGG

60.0

[34]

---

214

215 **2.1.8. Immune gene expression**

216 **2.1.8.1. RNA extraction and cDNA synthesis**

217 RNA was extracted from samples preserved in TRI Reagent (Sigma, UK) following the  
218 manufacturer's instructions. The quality and quantity of the RNA was assessed by  
219 electrophoresis and was measured with a Nanodrop 1000 spectrophotometer (NanoDrop  
220 Technologies, Thermo Fisher Scientific), respectively. The RNA pellet was eluted in 15  $\mu$ L of  
221 nuclease-free water prior to cDNA synthesis with QuantiTect® Reverse Transcription kit  
222 (Qiagen) following the manufacturer's protocol. Samples were stored at -20 °C until use.

223 **2.1.8.2. Primers design and specificity**

224 Primers were designed for two target genes (MHC II - CD74 molecule and IgM) to assess the  
225 effect of bath vaccination on juvenile ballan wrasse adaptive immune responses. Furthermore,  
226 a primer pair was also designed for a housekeeping gene ( $\beta$ -actin), while previously published  
227 elongation factor  $\alpha$  (ELF $\alpha$ ) primers [24] were used as the second housekeeping gene for  
228 expression normalisation. The primers used for qPCR are provided in Table 3. Primers were  
229 designed based on sequencing similarity of > 88 % using NCBI Primer – Blast software  
230 (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) [38] and following good primer design criteria  
231 [39]. Validation of the newly designed primers was carried out and amplification products sent  
232 for sequencing. In brief, separate PCR amplifications were performed for all the primers pairs  
233 in a total reaction of 10  $\mu$ L each including; 1  $\mu$ L (1/20 cDNA synthesised of 250 ng of total RNA  
234 from whole larvae, HK, L, S, 0.1  $\mu$ M of each primer, 1 x Q5® Hot start high fidelity Master mix  
235 (2x) (NEB biolabs) and nuclease free water to reach the desirable final volume. The following  
236 thermal cycling conditions were used in a Biometra, TAdvanced: 1 cycle of initial denaturation  
237 at 98 °C for 1 min, 30 cycles at 95 °C for 15 sec (denaturation), 56 °C for 15 sec (annealing)  
238 and 72 °C for 15 sec (extension), followed by final extension 1 cycle at 72 °C for 2 min and a  
239 cooling step at 4 °C. The relative molecular weight of the amplicons was calculated against a  
240 100 bp marker (Thermo Fisher Scientific). Resulting PCR products were sequenced by  
241 Eurofins to confirm identity.

242 **Table 3.** Primer pairs used for quantitative PCR (qPCR) to assess ballan wrasse adaptive  
243 immune responses.



Target gene	Oligo name	Sequence	Annealing (°C)	Accession number	Source
Major histocompatibility complex II (MHC II)	MHC II F_5	CCAATGTTGCGTCTCCGATG	56	XM_020645324.2	This study
	MHC II R_5	CACTGACTACAGCGTCCTGC			
Immunoglobulin M (IgM)	IgM F1	AGTCCCGTATGTCCTGTGGT	56	KX688616.1	This study
	IgM R1	CCCTTGCTCGTCTGTCCAAT			
Beta actin $\beta$ -actin	$\beta$ -actin F4	GCCAGAAGGACAGCTACGTT	56	XM_029281812.1	This study
	$\beta$ -actin R4	TCGATGGGGTACTTCAGGGT			
Elongation factor alpha (ELF- $\alpha$ )	ELF $\alpha$ F	ATTGATGCCCTGGACAC	56		[24]
	ELF $\alpha$ R	CCTCAAACCTCACCGACACC			

244


### 245 2.1.8.3. Quantitative PCR (qPCR)

246 The expression of target genes; MHC II and IgM and housing keeping genes;  $\beta$ -actin and  
247 ELF $\alpha$  was measured on a LightCycler®480 instrument (Roche Diagnostics). Target gene  
248 expression was normalised against the  $\beta$ -actin and ELF $\alpha$  housekeeping genes. Every reaction  
249 contained a final volume of 10  $\mu$ L, containing 1  $\mu$ L of cDNA (either sample or negative control;  
250 nuclease free water), 5  $\mu$ L of Luminaris Color HiGreen q PCR MasterMix (Thermo Fisher  
251 Scientific), 0.5  $\mu$ L of each forward and reverse primers, and 3.0  $\mu$ L of nuclease free water.  
252 Cycling conditions consisted of an initial activation of DNA polymerase at 95 °C for 10 min,  
253 followed by 40 cycles of denaturation at 95 °C for 15 sec, annealing at 56 °C for 30 sec, and  
254 extension at 72 °C for 30 sec. The assay included duplicates of each sample, non-template  
255 control and a serial dilution (from 10 to 10<sup>8</sup> copies per 1  $\mu$ l) of a linearised plasmid construct  
256 pGEM-T Easy vector (Promega, France) which incorporated a PCR fragment of the target  
257 gene. Melt curves were checked to ensure specific amplification. Crossing points (Cp) values  
258 were calculated employing the second derivative maximum method in the LightCycler  
259 software (Supplementary Table 2).

### 260 2.2. Ontogeny of the adaptive immune system

261 Whole ballan wrasse larvae were sampled and fixed in RNA stabilisation buffer from the same  
262 production tank at nine time points from 0, 15, 30, 60, 90, 115, 130 and 150 (dph) (0.3  $\pm$  0.1 g  
263 at 60 dph and 1.0  $\pm$  0.2 g at 150 dph) from two commercial hatcheries (site A and B) in Scotland  
264 (n= 20 per time point per hatchery). The samples were stored at - 20 °C until processing.

265 Whole larvae were processed for times 0 – 60 dph. The head, tail, and dorsal area (dorsal fin  
266 and muscle) were removed prior to RNA extraction for larvae from 90 - 150 dph to enrich  
267 transcripts associated with target lymphoid organs and gut regions as shown in Figure 1. The  
268 ontogeny of immune response was assessed with qPCR as described below.



269

270 **Figure 1.** Juvenile ballan wrasse from > 90 days post hatch (dph) used in ontogeny study.  
271 Dashed line shows the regions where the head, tail, and dorsal area (dorsal fin and muscle)  
272 were removed from the larvae prior to RNA extraction to enrich transcripts associated with  
273 target lymphoid organs and gut.

### 274 **2.3. Statistical analysis**

275 The geometric mean was calculated for housekeeping gene concentration and used for  
276 normalisation of target genes concentration (MHC II –CD 74 and IgM). The mean and the  
277 standard error were calculated for each target gene. MHC II–CD 74 and IgM expression was  
278 assessed and visualised on a histogram graph using Sigma Plot (SYSTAT inc Software, Inc.,  
279 San Jose California USA). Statistical analysis was conducted using Minitab software  
280 (Minitab18). Gene expression transcript concentrations were assessed with General Linear  
281 Model (GLM) for normality and homogeneity of variance and data were normalised using Log<sub>10</sub>  
282 or natural log (LN). Determination of whether a difference in expression occurred for either of  
283 the genes during the ontogeny and challenge trial studies was tested by ANOVA with Tukey's  
284 post hoc test (significance level;  $p < 0.05$ ). The results from qPCR for *As* bacterial load in  
285 tissues samples from challenged fish in this experiment were tested using Mood's median  
286 non-parametric test (significance level;  $p < 0.05$ ). Statistical analysis on bath challenge data  
287 were performed in statistical software R [40] for Kaplan – Meier survival curves, log-rank non-

288 parametric tests test (significance level;  $p < 0.05$ ) [41, 42] and pairwise comparison test  
289 (significance level;  $p < 0.05$ ) [43].

## 290 **2.4. Ethics statement**


291 Animal procedures for bacterial infection and vaccination were performed under UK  
292 Government Home Office project licence PC5CD1B81 following approval by the Animal  
293 Welfare and Ethical Review Body (AWERB) at the Centre for Environment, Fisheries and  
294 Aquaculture Science (Cefas) and University of Stirling. Ballan wrasse were treated in  
295 accordance with the Animals (Scientific Procedures) Act 1986 Amended Regulations (SI  
296 2012/3039).

## 297 **3. Results**

### 298 **3.1. Immersion vaccine efficacy trial**

#### 299 **3.1.1. Cumulative mortalities**

300 Morbidities in the small cohorts for both the mock and vaccinated groups started at 8 dpi and  
301 10 dpi, respectively, while morbidities in the large cohorts for prime vaccinated groups were  
302 noted at 9 dpi (mock) and 13 dpi (vaccinated) (Figure 2). Morbidities were first recorded in the  
303 booster vaccinated group from the large cohort at 8 dpi followed by the mock vaccinated at 9  
304 dpi except from a single tank (T08 – 34) for which morbidities started earlier at 3 dpi even  
305 though these fish received a similar infection dose than the others from the same group. The  
306 survival rate from this tank was the lowest from all groups reaching 33.3 % (Figure 2). The  
307 mean cumulative mortalities for the large fish challenged after booster vaccination were  
308 calculated with and without tank T08 – 34 (Table 4). Mean cumulative mortalities (%) for  
309 vaccinated and unvaccinated groups (either prime or booster) for both cohorts (small and  
310 large) were not statistically different (Table 4). Survival rates (%) did not differ between  
311 vaccinated and mock vaccinated groups (Figure 2). Control fish from both prime and booster  
312 vaccinated groups in the large population experienced 4 and 2 % mortalities, respectively, but  
313 bacteriology and molecular analysis showed that they were not caused by aAs. No mortalities  
314 were recorded in the control fish from the small cohort.



317 **Figure 2.** Survival curves for juvenile ballan wrasse bath challenged with atypical *Aeromonas*  
 318 *salmonicida vapA* type V upon prime (both cohorts) and booster (large cohort only) immersion  
 319 vaccination with an autogenous vaccine at two developmental stages. Small (0.5 g ± 0.2 g or  
 320 ca. 80 days post hatch; dph) (A) and large cohort (1.5 ± 0.4 g or ca. 170 dph) (B) following  
 321 prime vaccination and large cohort (C) upon booster vaccination. The vaccinated group is  
 322 presented as Vac (1-3) and mock vaccinated as Mock vac (1-3) on the legend.

324 **Table 4.** Challenge dose (CFU / mL) of atypical *Aeromonas salmonicida vapA* type V and  
 325 mean cumulative mortalities (%) with standard deviation ( $\pm$ SD) for ballan wrasse juveniles (n=  
 326 50). Small and large cohorts were prime vaccinated, and only a group of fish from the large  
 327 cohort received booster vaccination. The immunisation period for prime and booster groups  
 328 were 645 and 420 DD, respectively. No mortalities were recorded for control groups. Non-  
 329 specific mortalities were recorded in one of the replicates from the vaccinated group of the  
 330 large cohort (T08-34). The mean cumulative mortalities were calculated with and without this  
 331 tank. No statistical difference ( $p < 0.05$ ) was observed between the different groups or cohorts.

Cohort	Vaccination status	Mean CFU / mL	Mean cumulative mortalities (%)
Small	Vaccinated	2.98E+09	28.0 $\pm$ 8.7 <sup>a</sup>
	Mock vaccinated	5.23E+09	23.0 $\pm$ 13.6 <sup>a</sup>
Large	Vaccinated	5.65E+08	34.0 $\pm$ 2.0 <sup>b</sup>
	Mock vaccinated	9.95E+08	37.0 $\pm$ 7.0 <sup>b</sup>
	Booster vaccinated	3.77E+09	30.0 $\pm$ 7.0 (*54 $\pm$ 15.7) <sup>c,d</sup>
	Mock booster vaccinated	5.11E+09	40.0 $\pm$ 12.4 <sup>c,d</sup>

332

333 \*Mean cumulative mortalities taking into account T08 – 34.

334 a,b,c,d statistical difference at trial termination  $p = 0.29$ ,  $p = 0.232$ ,  $p = 0.232$ ,  $p = 0.691$  without  
 335 (T08 – 34), respectively.

### 336 3.1.2. Bacteria presence and load in morbid fish

337 Head kidney swabs from moribund fish plated on TSA had small pin head creamy aAs like  
 338 colonies and were positive with the agglutination test, except for three samples from tank T08-  
 339 34 between 3 and 8 dpi. The As qPCR loads were similar for all tissue samples (HK, L and S)  
 340 from both vaccinated and mock vaccinated fish with Cp value ranging from 21 to 35 (Table 5).  
 341 The conventional PCR specific for aAs type V confirmed the presence of aAs type V bacteria  
 342 in the same samples as qPCR. Large moribund fish (n= 3) from tank T08-34 which had been

343 booster vaccinated and challenged with aAs type V were negative for all 3 screening methods  
 344 and bacteria did not grow on TSA plates.

345 **Table 5.** Bacterial load in tissues samples (head kidney, liver and spleen) from small and large  
 346 cohort ballan wrasse (n= 15 per treatment 10 %) challenged with atypical *Aeromonas*  
 347 *salmonicida vapA* type V following vaccination (prime and booster; only large) with a  
 348 polyvalent autogenous vaccine or mock vaccination. Load was assessed by quantitative PCR  
 349 and expressed as mean cross point (Cp), Cp values > 35 considered negative. No statistically  
 350 significant differences between treatments were observed..


Cohort	Vaccination status	Mean bacterial load (Cp) in tissue		
		Kidney	Liver	Spleen
Small	Prime vaccinated	27.64 ± 4.71	25.87 ± 4.47	29.64 ± 5.79
	Prime mock vaccinated	27.90 ± 4.21	26.92 ± 4.73	26.88 ± 3.14
Large	Prime vaccinated	29.14 ± 2.55	28.74 ± 4.22	28.15 ± 3.31
	Prime mock vaccinated	27.62 ± 5.06	25.17 ± 3.10	24.50 ± 2.32
	Booster vaccinated	29.18 ± 3.77	27.94 ± 3.51	27.51 ± 3.72
	Booster mock vaccinated	27.69 ± 2.11	25.60 ± 3.20	28.39 ± 3.70

351

### 352 3.1.3 Gene expression at pre- and post- vaccination and post- challenge

353 Overall, gene expression (MCH II –CD 74 and IgM) was not significantly different in tissues or  
 354 between groups (pre - and post- vaccination and vaccinated and mock vaccinated) after  
 355 challenge with aAs *vapA* type V (Figures 3 and 4) with a few exceptions. The relative  
 356 expression of both target genes was statistically significantly higher in head kidney and spleen  
 357 of both small and large booster control groups in comparison with challenged, mock and  
 358 vaccinated fish of the same cohorts.


359



360

361 **Figure 3.** Box and whisker plots showing the normalised number of MHC II and IgM transcripts  
 362 obtained by qPCR from head kidney (A, B), liver (C, D) and spleen (E, F) from small cohort  
 363 ballan wrasse at pre- and post- prime (24 h) vaccination by immersion and post – bath  
 364 challenge with atypical *Aeromonas salmonicida vapA* type V (n= 6 for each group). Pre-  
 365 vaccinated and vaccinated groups are presented as pre and vac, respectively and mock  
 366 vaccinated (vaccination control) as mock in the legend. Boxes represent the interquartile

367 range, the line inside the box shows the median value and the whiskers, outside the boxes,  
 368 relate to the maximum and minimum values. Statistically significant differences (Tukey's test,  
 369  $p < 0.05$ ) between groups are represented by superscripts.






371 **Figure 4.** Box and whisker plots showing the normalised number MHC II and IgM transcripts  
372 obtained by qPCR from head kidney (A, B), liver (C, D) and spleen (E, F) tissues of large  
373 cohort ballan wrasse at pre- and post- prime (24 h) vaccination by immersion and post – bath  
374 challenge with atypical *Aeromonas salmonicida vapA* type V (n= 6 for each group). Boxes  
375 represent the interquartile range, the line inside the box shows the median value and the  
376 whiskers, outside the boxes, relate to the maximum and minimum values. Pre-vaccinated and  
377 vaccinated groups are presented as pre and vac, respectively and mock vaccinated  
378 (vaccination control) as mock in the legend. Statistically significant differences (Tukey's test,  
379  $p < 0.05$ ) between the groups are represented by superscripts.

### 380 **3.2. Ontogeny**

381 A significant increase in expression of both MHC II and IgM was noticed in both surveyed  
382 hatchery site populations. Expression levels of MHC II were significantly higher than basal  
383 levels (0 dph) from 35 dph (Figure 5 A). Temporal gene expression continued to increase until  
384 115 dph when expression remained unchanged until the end of the study (150 dph).  
385 Additionally, the IgM expression levels significantly increased by 75 dph in comparison to  
386 earlier time points and continued to increase significantly until 130 dph (Figure 5 B). Similar  
387 trends were noticed both for MHC II and IgM expression for site B (Figure 5 C and D).



388

389 **Figure 5.** Box and whisker plots showing the normalised number of MHC II and IgM transcripts  
 390 obtained by qPCR from whole ballan wrasse larvae (n= 6 for each group) from 0 to 150 days  
 391 post hatch (dph) at site A (A, B) and B (C, D). Boxes represent the interquartile range, the line  
 392 inside the box shows the median value and the whiskers, outside the boxes, relate to the  
 393 maximum and minimum values. Statistically significant differences (Tukey's test,  $p < 0.05$ )  
 394 between the time points are represented by superscripts.

#### 395 4. Discussion

396 This study aimed to assess the efficacy of an immersion autogenous vaccine (Ridgeway  
 397 Biologicals LTB) used for early juvenile developmental stages (<15 g) in ballan wrasse  
 398 hatcheries in Scotland and then determine the ontogeny of key representative components of  
 399 the adaptive immune system of the species. This would ultimately assist the establishment of  
 400 a vaccination timeframe for farmed ballan wrasse for protection against aAs. The results  
 401 showed no significant differences in the mean cumulative mortalities for immersion vaccinated  
 402 juvenile ballan wrasse (both small cohort and large cohorts) challenged with aAs by bath  
 403 inoculation post immunisation. The expression of MHC II (CD74 molecule) and IgM transcripts  
 404 were first identified at 35 and 75 dph, respectively and were observed to increase until

405 termination of the study at 150 dph which suggests initiation towards maturation of adaptive  
406 immunity.

407 In the present study all vaccination groups (prime small (ca. 0.5 g) and large (ca. 1.5 g), and  
408 booster large) along with mock vaccinated fish were challenged against aAs type V post  
409 immunisation (prime at 654 DD and booster at 495 DD) using a challenge model we developed  
410 previously [32]. Morbidities between vaccinated and mock vaccinated fish were not statistically  
411 different for any of the three groups indicating that the vaccine did not confer protection in the  
412 fish under the conditions used in this study. This confirms previous reports in which immersion  
413 vaccination was shown to have low potency [9]. Chakraborty, et al. [14] reported no protection,  
414 but an apparent delayed infection (ca. 3 weeks) in juvenile lumpsuckers (ca. 2.9 g) vaccinated  
415 with a commercial polyvalent vaccine against *Vibrio anguillarum* following a full vaccination  
416 regime (prime and booster bath and i.p. injection). Vaccinated fish showed higher bacterial  
417 loads in some cases, however no significant differences were found in the present study  
418 between vaccinated and mock vaccinated fish. Importantly, it must be acknowledged that  
419 while qPCR detected the bacterial load within the tested tissues, it could not distinguish  
420 between live and dead cells.

421 The vaccine protection was also evaluated in prime and booster immunised fish after  
422 challenge with aAs by profiling the relative expression of adaptive immune related genes  
423 (MHC II – CD74 molecule and IgM). No statistically significant differences were noted between  
424 treatments (mock and vaccinated fish) post challenge (up to 30 dpi) for any of the three groups.  
425 The relative expression levels of both target genes were higher in the head kidney and spleen  
426 sampled from control fish in both the small and large cohorts in comparison to the challenged  
427 mock and vaccinated fish of the same cohort. MHC II (CD74 molecule) expression was higher  
428 in the spleen of all groups but not in other tissues at the end of the challenge period. MHC II  
429 expression was higher in liver tissues for all cohorts post infection in this study, suggesting a  
430 functional role of MHC II in this tissue. These results agree with a previous report for miiuy  
431 croaker, *Miichthys miiuy* [44]. Antigen presentation of MHC II molecules may occur in other  
432 organs like the liver, which very recently has been considered to have an immunological role  
433 in mammals [45, 46]. Yang, et al. [47] and Causey, et al. [48] suggested a similar role of the  
434 liver in teleost fish as lymphocytes are present in the organ. In the present study, the total IgM  
435 expression levels were similar in the lymphoid organs for both mock and vaccinated groups  
436 suggesting no vaccine-associated immunity against aAs. In a previous study, expression of  
437 membrane IgM (mIgM) was located in different tissues in fish challenged with *Photobacterium*  
438 *damselae subsp. piscicida* prior (spleen and gut) and post vaccination (only in spleen) [49].  
439 Thus, comparative analysis of gene expression with ELISA for specific antibody presence (IgM

440 and IgT) was undertaken previously by the authors and confirmed that vaccination resulted in  
441 a systemic IgM response while the challenge resulted in a local IgT response only [49]. Despite  
442 several attempts to collect blood samples using a capillary tube, blood isolation was not  
443 successful in the present study due to the small size of the fish, and as such serology tests  
444 could not be performed although we previously demonstrated enhanced serum IgM levels in  
445 i.p. vaccinated wrasse that correlated with protection [25].

446 While the organogenesis in ballan wrasse has been described previously [50, 51], the onset  
447 of the expression of immune related genes in tissues of ballan wrasse species has not been  
448 described to date and the immune system of *L. bergylta* is currently only partially characterised  
449 [21-23]. In this study, the temporal expression of MHC II (CD74) and IgM transcripts was  
450 investigated in newly hatched ballan wrasse larvae (0 dph) until 150 dph using qPCR in two  
451 commercial hatcheries in Scotland. MHC II and IgM expression levels significantly increased  
452 at 35 and 75 dph, respectively, and continued to do so in larvae originating from site A. MHC  
453 I and II are responsible for antigen recognition and presentation and play a significant role in  
454 activation of adaptive immune response and memory [44]. Hence, the detection of MHC II  
455 gene expression prior to IgM in the study was expected considering the role that MHC II plays  
456 in immunological processing in phagocytes [45]. Saravanan, et al. [52] were the first to  
457 document expression of MHC II in Rohu (*Labeo rohita*) at 28 dph. Earlier expression of MHC  
458 II molecules has been observed for sea bass (*Dicentrarchus labrax*) at 4 dph [47] and common  
459 carp (*Cyprinus carpio*) at 28 dpf [48]. IgM mRNA transcripts were first detected at 5 dph with  
460 a significant increase observed at 35 dph for olive flounder, *Paralichthys olivaceus* [49]. In  
461 teleost fish maternal immune elements have a significant role in survival of eggs and newly  
462 hatched larvae. Innate (lysozyme and antibacterial peptides) and adaptive (IgM) immune  
463 elements can pass from broodstock to offspring [15, 53, 54]. However, maternal immunity has  
464 only a short duration of several weeks post hatch (3 – 4) [55]. For example, IgM decreased in  
465 striped trumpeter (*Latris lineata*) by 100 dph [56]. In the present study, the temporal expression  
466 profile observed for MCH II and IgM in ballan wrasse from 35 dph and 75 dph, respectively  
467 continued to increase suggesting a relationship with the animal's size and immune  
468 responsiveness appeared therefore to not be derived from maternal transfer. Differences in  
469 expression patterns and relative gene expression were observed for MHC II (CD 74) and IgM  
470 between the two hatcheries. This could be explained by the fact larvae from site B were mixed  
471 with larvae from other developmental stages (earlier and later) during husbandry procedures  
472 (*i.e.* grading) in contrast with larvae from site A whereby the same batch and tank of fish were  
473 sampled during the entire study. Thus, the results from site A are considered more reliable.

474 The ontogeny results indicated that expression of MHC II and IgM should have been initiated  
475 at least for the large cohort used in the immersion vaccination study prior to vaccination (ca.  
476 170 dph at prime vaccination). Indeed, the overall relative expression of MHC II and IgM was  
477 higher in the tissues (HK, L and S) of both fish sizes at vaccination in comparison with  
478 ontogeny fish samples, which included multiple pooled organs including gut tissue.  
479 Nevertheless, the mean cumulative mortality and the gene expression results of the  
480 immunised and bath challenged fish in this study indicates that the vaccine did not elicit  
481 protection. It is important to mention that the detection and expression of these adaptive  
482 immune genes only provides an indication of the onset of immunocompetence but not full  
483 immunological maturity [57]. Similarly, previous studies have demonstrated that presence of  
484 lymphoid organs and initiation of IgM gene expression does not guarantee  
485 immunocompetence [58-61] which may explain the result of the immersion vaccination. In  
486 addition, there are various factors apart from immunocompetence that need to be considered  
487 for a successful vaccine e.g. the vaccine formulation, vaccine dose, the vaccination route,  
488 vaccination time for immersion vaccines, the use of an adjuvant, and the challenge trial  
489 including pathogen pressure / load, exposure time and virulence of the pathogen [62-64]. In  
490 addition, multivalent vaccines need rigorous testing and careful selection of strains or different  
491 species included in the formulation as combinations of antigens could interact synergistically  
492 or antagonistically and stimulate, cross-react, compete and lead to immunosuppression [65].  
493 Nikoskelainen, et al. [66] reported that different multivalent vaccines composition against *A.*  
494 *salmonicida subsp. salmonicida*, *Listonella anguillarum* and *Flavobacterium psychrophilum*  
495 had an effect in antibody production in rainbow trout (*Oncorhynchus mykiss*). The results  
496 obtained from this study further illustrate the challenges of immersion vaccination against  
497 typical and atypical strains of *As*, for which historically immersion vaccination has proven  
498 challenging and only few experimental successes have been reported with demonstrated  
499 protection to fish [2, 67-69].

500 Overall, the results of the vaccine trial conducted in this study suggested that immersion  
501 vaccination for farmed ballan wrasse should be applied at a later stage of the production cycle  
502 (e.g. size > 1.5 g), at least for the aAs component of the vaccine. In addition, we reported for  
503 first time, the onset of adaptive immune genes expression for ballan wrasse (MHC II – CD74  
504 at 35 dph and IgM at 75 dph), which is critical for commercial vaccination strategies. Further  
505 research is needed to better understand the adaptive immune system of juvenile ballan wrasse  
506 including, its maturation and uptake and processing of antigens for ballan wrasse juveniles.  
507 All of the above will help to optimise vaccine formulation and vaccination regimes in hatcheries  
508 and mitigate the high mortalities occurring during early developmental stages. Protection at  
509 this early stage is vital to boost the cleaner fish sector productivity to supply the increasing

510 demand from salmon producers for healthy, disease free and effective delousing ballan  
511 wrasse.

## 512 **5. Acknowledgements**

513 The authors would like to thank MOWI Scotland and Scottish Sea Farms for providing the  
514 experimental animals, supported the fish transportation and overall collaboration during the  
515 project. We also thank Ridgeway Biologicals for providing the bacteria isolates and the  
516 vaccine, and the aquarists and bacteriology team at Cefas for their technical assistance during  
517 the challenge trial.

518

## 519 **6. Funding information**

520 This project was co-funded through the Sustainable Aquaculture Innovation Centre (SAIC)  
521 and University of Stirling (grant number: SL-2015-01).

522

## 523 **7. Conflict of interest Statement**

524 The authors declare no conflict of interest.

525

## 526 **8. References**

527 [1] B. Hjeltnes, B. Bang-Jensen, G. Bornø, A. Haukaas, C.S. Walde, The Health Situation in  
528 Norwegian Aquaculture 2018, Norwegian Veterinary Institute, Oslo, Norway, (2019), p. 132.

529 [2] B. Austin, D.A. Austin, Aeromonadaceae representative (*Aeromonas salmonicida*),  
530 Bacterial fish pathogens, Springer 2016, pp. 215-321.

531 [3] S. Gulla, S. Bayliss, B. Bjornsdottir, I. Dalsgaard, O. Haenen, E. Jansson, U. McCarthy,  
532 F. Scholz, M. Vercauteren, D. Verner-Jeffreys, T. Welch, T. Wiklund, D.J. Colquhoun,  
533 Biogeography of the fish pathogen *Aeromonas salmonicida* inferred by *vapA* genotyping,  
534 FEMS Microbiol. Lett. 366(7) (2019) fnz074. <https://doi.org/10.1093/femsle/fnz074>.

535 [4] S. Gulla, V. Lund, A. Kristoffersen, H. Sørum, D. Colquhoun, *vapA* (A-layer) typing  
536 differentiates *Aeromonas salmonicida* subspecies and identifies a number of previously  
537 undescribed subtypes, J Fish Dis 39(3) (2016a) 329-342. <https://doi.org/10.1111/jfd.12367>.

538 [5] A.J. Brooker, A. Papadopoulou, C. Gutierrez, S. Rey, A. Davie, H. Migaud, Sustainable  
539 production and use of cleaner fish for the biological control of sea lice: recent advances and  
540 current challenges, Vet Rec 183(12) (2018) 383. 10.1136/vr.104966.

541 [6] M. Dadar, K. Dhama, V.N. Vakharia, S.H. Hoseinifar, K. Karthik, R. Tiwari, R. Khandia, A.  
542 Munjal, C. Salgado-Miranda, S.K. Joshi, Advances in aquaculture vaccines against fish

- 543 pathogens: global status and current trends, *Rev Fish Sci Aquac* 25(3) (2017) 184-217.  
544 <https://doi.org/10.1080/23308249.2016.1261277>.
- 545 [7] I. Sommerset, B. Krossoy, E. Biering, P. Frost, Vaccines for fish in aquaculture, *Expert*  
546 *Rev Vaccines* 4(1) (2005) 89-101. <https://doi.org/10.1586/14760584.4.1.89>.
- 547 [8] B.E. Brudeseth, R. Wiulsirod, B.N. Fredriksen, K. Lindmo, K.E. Lokling, M. Bordevik, N.  
548 Steine, A. Klevan, K. Gravningen, Status and future perspectives of vaccines for  
549 industrialised fin-fish farming, *Fish Shellfish Immunol* 35(6) (2013) 1759-68.  
550 <https://doi.org/10.1016/j.fsi.2013.05.029>.
- 551 [9] A. Toranzo, J. Romalde, B. Magariños, J. Barja, Present and future of aquaculture  
552 vaccines against fish bacterial diseases, *Options Mediterraneennes* 86 (2009) 155-176.
- 553 [10] J. Bogwald, R.A. Dalmo, Review on immersion vaccines for fish: An update 2019,  
554 *Microorganisms* 7(12) (2019) 627. <https://doi.org/10.3390/microorganisms7120627>.
- 555 [11] E. Soto, M.J. Griffin, J.A. Tobar, 11 - Mucosal vaccines, in: B.H. Beck, E. Peatman  
556 (Eds.), *Mucosal Health in Aquaculture*, Academic Press, San Diego, 2015, pp. 297-323.  
557 <https://doi.org/10.1016/B978-0-12-417186-2.00011-X>.
- 558 [12] A. Zapata, B. Diez, T. Cejalvo, C. Gutierrez-de Frias, A. Cortes, Ontogeny of the  
559 immune system of fish, *Fish Shellfish Immunol* 20(2) (2006) 126-36.  
560 [10.1016/j.fsi.2004.09.005](https://doi.org/10.1016/j.fsi.2004.09.005).
- 561 [13] B. Razquin, A. Castillo, P. Lopez-Fierro, F. Alvarez, A. Zapata, A. Villena, Ontogeny of  
562 IgM-producing cells in the lymphoid organs of rainbow trout, *Salmo gairdneri* Richardson: An  
563 immuno- and enzyme-histochemical study, *Journal of Fish Biology* 36(2) (1990) 159-173.  
564 <https://doi.org/10.1111/j.1095-8649.1990.tb05592.x>.
- 565 [14] S. Chakraborty, T. Cao, A. Hossain, H. Gnanagobal, I. Vasquez, D. Boyce, J.  
566 Santander, Vibrogen-2 vaccine trial in lumpfish (*Cyclopterus lumpus*) against *Vibrio*  
567 *anguillarum*, *J Fish Dis* 42(7) (2019) 1057-1064. <https://doi.org/10.1111/jfd.13010>.
- 568 [15] B. Magnadottir, S. Lange, S. Gudmundsdottir, J. Bogwald, R.A. Dalmo, Ontogeny of  
569 humoral immune parameters in fish, *Fish Shellfish Immunol* 19(5) (2005) 429-39.  
570 [10.1016/j.fsi.2005.03.010](https://doi.org/10.1016/j.fsi.2005.03.010).
- 571 [16] M.-G. Mao, J.-L. Lei, P.-M. Alex, W.-S. Hong, K.-J. Wang, Characterization of RAG1  
572 and IgM (mu chain) marking development of the immune system in red-spotted grouper  
573 (*Epinephelus akaara*), *Fish Shellfish Immunol* 33(4) (2012) 725-735.  
574 <https://doi.org/10.1016/j.fsi.2012.06.011>.
- 575 [17] H. Gong, Q. Wang, Y. Lai, C. Zhao, C. Sun, Z. Chen, J. Tao, Z. Huang, Study on  
576 immune response of organs of *Epinephelus coioides* and *Carassius auratus* after immersion  
577 vaccination with inactivated *Vibrio harveyi* vaccine, *Front. Immunol.* 11 (2021) 3766.
- 578 [18] C. Monzón-Argüello, C.G. De Leaniz, G. Gajardo, S. Consuegra, Eco-immunology of  
579 fish invasions: the role of MHC variation, *Immunogenetics* 66(6) (2014) 393-402.  
580 <https://doi.org/10.1007/s00251-014-0771-8>.
- 581 [19] A. Sato, F. Figueroa, B.W. Murray, E. Málaga-Trillo, Z. Zaleska-Rutczynska, H.  
582 Sültmann, S. Toyosawa, C. Wedekind, N. Steck, J. Klein, Nonlinkage of major  
583 histocompatibility complex class I and class II loci in bony fishes, *Immunogenetics* 51(2)  
584 (2000) 108-116.

- 585 [20] A.B. Wilson, MHC and adaptive immunity in teleost fishes, *Immunogenetics* 69(8)  
586 (2017) 521-528. [10.1007/s00251-017-1009-3](https://doi.org/10.1007/s00251-017-1009-3).
- 587 [21] G.T. Haugland, A. Ronneseth, H.I. Wergeland, Flow cytometry analyses of phagocytic  
588 and respiratory burst activities and cytochemical characterization of leucocytes isolated from  
589 wrasse (*Labrus bergylta* A.), *Fish Shellfish Immunol* 39(1) (2014) 51-60.  
590 <https://doi.org/10.1016/j.fsi.2014.04.023>.
- 591 [22] S. Bilal, K.K. Lie, O.A. Karlsen, I. Hordvik, Characterization of IgM in Norwegian cleaner  
592 fish (lumpfish and wrasses), *Fish Shellfish Immunol* 59 (2016) 9-17.  
593 <https://doi.org/10.1016/j.fsi.2016.09.063>.
- 594 [23] S. Bilal, K.K. Lie, O. Saele, I. Hordvik, T Cell receptor alpha chain genes in the teleost  
595 ballan wrasse (*Labrus bergylta*) are subjected to somatic hypermutation, *Front. Immunol.*  
596 9(1101) (2018) 1101. <https://doi.org/10.3389/fimmu.2018.01101>.
- 597 [24] S. Bilal, K.K. Lie, A.S. Dalum, O.A. Karlsen, I. Hordvik, Analysis of immunoglobulin and  
598 T cell receptor gene expression in ballan wrasse (*Labrus bergylta*) revealed an  
599 extraordinarily high IgM expression in the gut, *Fish Shellfish Immunol* 87 (2019) 650-658.  
600 <https://doi.org/10.1016/j.fsi.2019.02.007>.
- 601 [25] J.G. Ramirez-Paredes, D. Verner-Jeffreys, A. Papadopoulou, S.J. Monaghan, L. Smith,  
602 D. Haydon, T.S. Wallis, A. Davie, A. Adams, H. Migaud, A commercial autogenous injection  
603 vaccine protects ballan wrasse (*Labrus bergylta*, Ascanius) against *Aeromonas salmonicida*  
604 *vapA* type V, *Fish Shellfish Immunol* 107 (2020) 43-53.  
605 <https://doi.org/10.1016/j.fsi.2020.09.040>.
- 606 [26] A. Papadopoulou, T. Wallis, J.G. Ramirez-Paredes, S.J. Monaghan, A. Davie, H.  
607 Migaud, A. Adams, Atypical *Aeromonas salmonicida vapA* type V and *Vibrio* spp. are  
608 predominant bacteria recovered from ballan wrasse *Labrus bergylta* in Scotland, *Diseases of*  
609 *Aquatic Organisms* 140 (2020) 47-54.
- 610 [27] A. Klindworth, E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, F.O. Glockner,  
611 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-  
612 generation sequencing-based diversity studies, *Nucleic Acids Res* 41(1) (2013) e1.  
613 <https://doi.org/10.1093/nar/gks808>.
- 614 [28] S. Yamamoto, H. Kasai, D.L. Arnold, R.W. Jackson, A. Vivian, S. Harayama, Phylogeny  
615 of the genus *Pseudomonas*: intrgeneric structure reconstructed from the nucleotide  
616 sequences of *gyrB* and *rpoD* genes, *Microbiology (Reading)* 146 ( Pt 10)(10) (2000) 2385-  
617 2394. <https://doi.org/10.1099/00221287-146-10-2385>.
- 618 [29] N.D. Young, I. Dykova, B.F. Nowak, R.N. Morrison, Development of a diagnostic PCR to  
619 detect *Neoparamoeba perurans*, agent of amoebic gill disease, *J Fish Dis* 31(4) (2008) 285-  
620 95. <https://doi.org/10.1111/j.1365-2761.2008.00903.x>.
- 621 [30] Office International des Epizooties (OIE) Manual of diagnostic tests for aquatic animals,  
622 Chapter 2.3, (2018).
- 623 [31] A. Oboh, M.B. Betancor, D.R. Tocher, O. Monroig, Biosynthesis of long-chain  
624 polyunsaturated fatty acids in the African catfish (*Clarias gariepinus*): Molecular cloning and  
625 functional characterisation of fatty acyl desaturase (*fads2*) and elongase (*elovl2*) cDNAs7,  
626 *Aquaculture* 462 (2016) 70-79. <https://doi.org/10.1016/j.aquaculture.2016.05.018>.



- 627 [32] A. Papadopoulou, K. Garvey, T. Hill, J.G. Ramirez-Paredes, S.J. Monaghan, J.L. Baily,  
628 A. Davie, I. Katsiadaki, D. Verner-Jeffreys, T. Wallis, H. Migaud, A. Adams, Novel atypical  
629 *Aeromonas salmonicida* bath challenge model for juvenile ballan wrasse (*Labrus bergylta*,  
630 *Ascanius*), *J Fish Dis* (2020). 10.1111/jfd.13312.
- 631 [33] A. Papadopoulou, A. Davie, S.J. Monaghan, H. Migaud, A. Adams, Development of  
632 diagnostic assays for differentiation of atypical *Aeromonas salmonicida vapA* type V and VI  
633 in ballan wrasse (*Labrus bergylta*, *Ascanius*), *J Fish Dis* In review (2020).
- 634 [34] S. Gulla, S. Duodu, A. Nilsen, I. Fossen, D. Colquhoun, *Aeromonas salmonicida*  
635 infection levels in pre-and post-stocked cleaner fish assessed by culture and an amended  
636 qPCR assay, *J Fish Dis* 39(7) (2016b) 867-877. <https://doi.org/10.1111/jfd.12420>.
- 637 [35] T. Khanam, A. Davie, B.J. McAndrew, D.J. Penman, DNA sampling from mucus in the  
638 Nile tilapia, *Oreochromis niloticus*: minimally invasive sampling for aquaculture-related  
639 genetics research, *Aquaculture Research* 47(12) (2016) 4032-4037.  
640 <https://doi.org/10.1111/are.12809>.
- 641 [36] A. Papadopoulou, A. Davie, S.J. Monaghan, H. Migaud, A. Adams, Development of  
642 diagnostic assays for differentiation of atypical *Aeromonas salmonicida vapA* type V and  
643 type VI in ballan wrasse (*Labrus bergylta*, *Ascanius*), *J Fish Dis* n/a(n/a) (2021).  
644 <https://doi.org/10.1111/jfd.13334>.
- 645 [37] D.P. Herlemann, M. Labrenz, K. Jürgens, S. Bertilsson, J.J. Waniek, A.F. Andersson,  
646 Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea,  
647 *The ISME journal* 5(10) (2011) 1571-1579. <https://doi.org/10.1038/ismej.2011.41>.
- 648 [38] J. Ye, G. Coulouris, I. Zaretskaya, I. Cutcutache, S. Rozen, T.L. Madden, Primer-  
649 BLAST: a tool to design target-specific primers for polymerase chain reaction, *BMC*  
650 *Bioinformatics* 13(1) (2012) 134. 10.1186/1471-2105-13-134.
- 651 [39] S. Taylor, M. Wakem, G. Dijkman, M. Alsarraj, M. Nguyen, A practical approach to RT-  
652 qPCR-Publishing data that conform to the MIQE guidelines, *Methods* 50(4) (2010) S1-5.  
653 <https://doi.org/10.1016/j.ymeth.2010.01.005>.
- 654 [40] R.C. Team, R: A language and environment for statistical computing. R 3.4. 4, Vienna,  
655 Austria: R Foundation for Statistical Computing., (2018).
- 656 [41] T. Therneau, A Package for Survival Analysis in S. version 2.38 URL: [https://CRAN.R-](https://CRAN.R-project.org/package=survival)  
657 [project.org/package= survival](https://CRAN.R-project.org/package=survival) (2015).
- 658 [42] T. Therneau, P.M. Grambsch, *The Cox model, Modeling survival data: extending the*  
659 *Cox model*, Springer, 2000, pp. 39-77.
- 660 [43] A. Kassambara, M. Kosinski, *Survminer: drawing survival curves using 'ggplot2'R*  
661 *package version 0.4. 0. 2017, (2017)*.
- 662 [44] T. Xu, Y. Sun, G. Shi, Y. Cheng, R. Wang, Characterization of the major  
663 histocompatibility complex class II genes in miiuy croaker, *PLoS One* 6(8) (2011) e23823.  
664 <https://doi.org/10.1371/journal.pone.0023823>.
- 665 [45] M.W. Robinson, C. Harmon, C. O'Farrelly, Liver immunology and its role in inflammation  
666 and homeostasis, *Cell. Mol. Immunol.* 13(3) (2016) 267-76.  
667 <https://doi.org/10.1038/cmi.2016.3>.

- 668 [46] P.A. Knolle, The Liver as a Lymphoid Organ, in: M.E. Gershwin, J.M. Vierling, M.P.  
669 Manns (Eds.), Liver Immunology, Springer International Publishing, Cham, 2014, pp. 55-64.  
670 10.1007/978-3-319-02096-9\_5.
- 671 [47] D. Yang, Q. Liu, M. Yang, H. Wu, Q. Wang, J. Xiao, Y. Zhang, RNA-seq liver  
672 transcriptome analysis reveals an activated MHC-I pathway and an inhibited MHC-II  
673 pathway at the early stage of vaccine immunization in zebrafish, BMC Genomics 13(1)  
674 (2012) 319. <https://doi.org/10.1186/1471-2164-13-319>.
- 675 [48] D.R. Causey, M.A.N. Pohl, D.A. Stead, S.A.M. Martin, C.J. Secombes, D.J. Macqueen,  
676 High-throughput proteomic profiling of the fish liver following bacterial infection, BMC  
677 Genomics 19(1) (2018) 719. <https://doi.org/10.3390/cells6040048>.
- 678 [49] M.C. Piazzon, J. Galindo-Villegas, P. Pereiro, I. Estensoro, J.A. Calduch-Giner, E.  
679 Gómez-Casado, B. Novoa, V. Mulero, A. Sitjà-Bobadilla, J. Pérez-Sánchez, Differential  
680 modulation of IgT and IgM upon parasitic, bacterial, viral, and dietary challenges in a  
681 Perciform fish, Front. Immunol. 7 (2016) 637. <https://doi.org/10.3389/fimmu.2016.00637>.
- 682 [50] E. Dunaevskaya, A.B. Amin, O.H. Ottesen, Organogenesis of ballan wrasse *Labrus*  
683 *bergylta* (Ascanius 1767) larvae, Journal of Aquaculture Research & Development 3(5)  
684 (2012) 142.
- 685 [51] M.R. Gagnat, P.-A. Wold, T. Bardal, G. Øie, E. Kjørsvik, Allometric growth and  
686 development of organs in ballan wrasse (*Labrus bergylta* Ascanius, 1767) larvae in relation  
687 to different live prey diets and growth rates, Biology open 5(9) (2016) 1241-1251.
- 688 [52] K. Saravanan, K. Rajendran, P. Gireesh-Babu, C. Purushothaman, M. Makesh,  
689 Ontogenic expression of adaptive immune genes in rohu, *Labeo rohita* (Hamilton-Buchanan,  
690 1822), (2017).
- 691 [53] T. Ellingsen, C. Strand, E. Monsen, J. Bogwald, R.A. Dalmo, The ontogeny of  
692 complement component C3 in the spotted wolffish (*Anarhichas minor* Olafsen), Fish  
693 Shellfish Immunol 18(5) (2005) 351-8. <https://doi.org/10.1016/j.fsi.2004.09.002>.
- 694 [54] M. Seppola, H. Johnsen, S. Mennen, B. Myrnes, H. Tveiten, Maternal transfer and  
695 transcriptional onset of immune genes during ontogenesis in Atlantic cod, Dev Comp  
696 Immunol 33(11) (2009) 1205-11. <https://doi.org/10.1016/j.dci.2009.06.013>.
- 697 [55] P. Swain, S. Dash, J. Bal, P. Routray, P. Sahoo, S. Sahoo, S. Saurabh, S. Gupta, P.  
698 Meher, Passive transfer of maternal antibodies and their existence in eggs, larvae and fry of  
699 Indian major carp, *Labeo rohita* (Ham.), Fish Shellfish Immunol 20(4) (2006) 519-527.  
700 <https://doi.org/10.1016/j.fsi.2005.06.011>.
- 701 [56] J. Covello, S. Bird, R. Morrison, A. Bridle, S. Battaglione, C. Secombes, B. Nowak,  
702 Isolation of *RAG-1* and *IgM* transcripts from the striped trumpeter (*Latris lineata*), and their  
703 expression as markers for development of the adaptive immune response, Fish Shellfish  
704 Immunol 34(3) (2013) 778-788.
- 705 [57] Y. Corripio-Miyar, S. Bird, J.W. Treasurer, C.J. Secombes, *RAG-1* and *IgM* genes,  
706 markers for early development of the immune system in the gadoid haddock,  
707 *Melanogrammus aeglefinus*, L, Fish Shellfish Immunol 23(1) (2007) 71-85.  
708 <https://doi.org/10.1016/j.fsi.2006.09.006>.

- 709 [58] N.M. dos Santos, N. Romano, M. de Sousa, A.E. Ellis, J.H. Rombout, Ontogeny of B  
710 and T cells in sea bass (*Dicentrarchus labrax*, L.), Fish Shellfish Immunol 10(7) (2000) 583-  
711 96. <https://doi.org/10.1006/fsim.2000.0273>.
- 712 [59] M. Mughal, M. Manning, Antibody responses of young carp, *Cyprinus carpio*, and grey  
713 mullet, *Chelon labrosus*, immunized with soluble antigen by various routes, Fish  
714 immunology, Elsevier, 1985, pp. 313-325. [https://doi.org/10.1016/B978-0-12-469230-  
715 5.50030-1](https://doi.org/10.1016/B978-0-12-469230-5.50030-1).
- 716 [60] L. Patrie-Hanson, A.J. Ainsworth, Humoral immune responses of channel catfish  
717 (*Ictalurus punctatus*) fry and fingerlings exposed to *Edwardsiella ictaluri*, Fish Shellfish  
718 Immunol 9(8) (1999) 579-589. <https://doi.org/10.1006/fsim.1999.0215>.
- 719 [61] I. Mulero, M.P. Sepulcre, I. Fuentes, A. García-Alcázar, J. Meseguer, A. García-Ayala,  
720 V. Mulero, Vaccination of larvae of the bony fish gilthead seabream reveals a lack of  
721 correlation between lymphocyte development and adaptive immunocompetence, Molecular  
722 Immunology 45(10) (2008) 2981-2989. <https://doi.org/10.1016/j.molimm.2008.01.017>.
- 723 [62] H.M. Munang'andu, B.N. Fredriksen, S. Mutoloki, R.A. Dalmo, O. Evensen, Antigen  
724 dose and humoral immune response correspond with protection for inactivated infectious  
725 pancreatic necrosis virus vaccines in Atlantic salmon (*Salmo salar* L), Vet Res 44(1) (2013)  
726 7. <https://doi.org/10.1186/1297-9716-44-7>.
- 727 [63] H.M. Munang'andu, S. Mutoloki, O. Evensen, Acquired immunity and vaccination  
728 against infectious pancreatic necrosis virus of salmon, Dev Comp Immunol 43(2) (2014)  
729 184-96. <https://doi.org/10.1016/j.dci.2013.08.008>.
- 730 [64] H.M. Munang'andu, S. Mutoloki, O. Evensen, An Overview of challenges limiting the  
731 design of protective mucosal vaccines for finfish, Front. Immunol. 6 (2015) 542.  
732 <https://doi.org/10.3389/fimmu.2015.00542>.
- 733 [65] R. Busch, Polyvalent vaccines in fish: the interactive effects of multiple antigens, Dev.  
734 Biol. Stand. 90 (1997) 245-256.
- 735 [66] S. Nikoskelainen, S. Verho, S. Järvinen, J. Madetoja, T. Wiklund, E.-M. Lilius, Multiple  
736 whole bacterial antigens in polyvalent vaccine may result in inhibition of specific responses  
737 in rainbow trout (*Oncorhynchus mykiss*), Fish Shellfish Immunol 22(3) (2007) 206-217.
- 738 [67] K.R. Villumsen, I. Dalsgaard, L. Holten-Andersen, M.K. Raida, Potential role of specific  
739 antibodies as important vaccine induced protective mechanism against *Aeromonas*  
740 *salmonicida* in rainbow trout, PLoS One 7(10) (2012).  
741 <https://doi.org/10.1371/journal.pone.0046733>.
- 742 [68] Y. Santos, S. Garcia-Marquez, P.G. Pereira, F. Pazos, A. Riaza, R. Silva, A. El Morabit,  
743 F.M. Ubeira, Efficacy of furunculosis vaccines in turbot, *Scophthalmus maximus* (L.):  
744 evaluation of immersion, oral and injection delivery, J Fish Dis 28(3) (2005) 165-72.  
745 <https://doi.org/10.1111/j.1365-2761.2005.00610.x>.
- 746 [69] M.R. Arkoosh, J.P. Dietrich, M.B. Rew, W. Olson, G. Young, F.W. Goetz, Exploring the  
747 efficacy of vaccine techniques in juvenile sablefish, *Anoplopoma fimbria*, Aquaculture  
748 Research 49(1) (2018) 205-216. <https://doi.org/10.1111/are.13449>.

749

750

751 **Supplementary Tables**

752

753 **Supplementary Table 1.** Vaccination doses (mL) are given below including calculated and  
 754 actual amounts for small and large cohorts.

Size (g) at vaccination	Vaccination status	Fish number	Biomass at vaccination (kg)	Tank (L)	Actual vaccine volume (mL)
0.5	Vaccinated	207.0	0.10	141.0	39.0
0.5	Mock	272.0	0.14	159.0	44.0
1.5	Vaccinated	207.0	0.31	165.0	46.0
1.5	Mock	272.0	0.41	151.0	42.0
2.5	Booster vaccinated	210.0	0.32	155.0	43.0
2.5	Booster mock	275.0	0.41	151.0	42.0

755

756 **Supplementary Table 2.** Standard curves quality testing on transformed plasmids containing  
 757 target and housekeeping genes (*IgM*, *MHC II*,  *$\beta$ -actin* and *ELF $\alpha$* ). Table contains the Cp upper  
 758 and lower limit, efficiency and slope of the each assay.

Target gene	Cp lower / upper limit	Efficiency	Slope
MHC II	10.51 / 33.7	1.958	-3.437
IgM	9.99 / 34.1	1.929	-3.504

---

$\beta$ -actin	10.4 / 34.6	1.976	-3.381
----------------	-------------	-------	--------

---

ELF $\alpha$	10.5 / 32.3	1.924	-3.519
--------------	-------------	-------	--------

---


759

760

761 **Supplementary Figures**

762

763



764

765 **Supplementary Figure 1.** Challenge of ballan wrasse juveniles with atypical *Aeromonas*  
766 *salmonicida vapA* type V in small aerated aquaria (8 L).

767

768

769 **Highlights**

770

- 771 • Autogenous vaccine offers no protection in farmed ballan wrasse at 0.5 g - 1.5 g
- 772 against atypical *Aeromonas salmonicida* (aAs) *vapA* V.
- 773 • Immersion vaccination for aAs should be performed at bigger sizes > 1.5 g.
- 774 • Onset of adaptive immune genes responses reported for first time.
- 775 • MHC II and IgM transcript were first reported at 35 and 75 days post hatch.
- 776

777