

1        **Inclusion of oil from transgenic *Camelina sativa* in feed effectively**  
2        **supplies EPA and DHA to Atlantic salmon (*Salmo salar*) grown to**  
3        **market size in seawater pens**

4  
5        Douglas R. Tocher<sup>1,2\*</sup> Matthew Sprague<sup>1</sup>, Lihua Han<sup>3</sup>, Olga Sayanova<sup>3</sup>, Fernando  
6        Norambuena<sup>4</sup>, Johnathan A. Napier<sup>3</sup> and Mónica B. Betancor<sup>1</sup>

7  
8        <sup>1</sup>Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA,  
9        United Kingdom

10        <sup>2</sup>Guangdong Provincial Key Laboratory of Marine Biotechnology, Institute of Marine Sciences,  
11        Shantou University, Shantou 515063, China

12        <sup>3</sup>Rothamsted Research, Harpenden AL5 2JQ, United Kingdom

13        <sup>4</sup>BioMar AS, Havnegata 9, 7010 Trondheim, Norway

14  
15        \* Corresponding author

16        E-mail: [monica.betancor@stir.ac.uk](mailto:monica.betancor@stir.ac.uk)

19 **Abstract**

20 Atlantic salmon were fed either a diet reflecting current commercial feeds with added oil supplied by  
21 a blend of fish oil and rapeseed oil (COM), or a diet formulated with oil from transgenic *Camelina*  
22 *sativa* containing 20% EPA+DHA (TCO). Salmon were grown from smolt to market size (>3kg) in  
23 sea pens under semi-commercial conditions. There were no differences in growth, feed efficiency or  
24 survival between fish fed the TCO or COM diets at the end of the trial. Levels of EPA+DHA in flesh  
25 of salmon fed TCO were significantly higher than in fish fed COM. A 140g fillet from TCO-fed  
26 salmon delivered 2.3g of EPA+DHA, 67% of the weekly requirement level recommended by many  
27 health agencies, and 1.5-fold more than the 1.5g of EPA+DHA for COM-fed fish. Oil from transgenic  
28 *Camelina* supported growth and improved the nutritional quality of farmed salmon in terms of  
29 increased “omega-3” supply for human consumers.

30

31 **Keywords:** Aquaculture; farmed salmon; omega-3; transgenic oil; EPA and DHA

## 32 **1. Introduction**

33 It is well established and widely accepted that the omega-3 (n-3) long-chain polyunsaturated fatty  
34 acids (LC-PUFA), eicosapentaenoic acid (20:5n-3; EPA) and docosahexaenoic acid (22:6n-3; DHA),  
35 have health-promoting effects in the human diet (Calder, 2018; Innes and Calder, 2020). Based on  
36 these health benefits, many national and international agencies across the world have set levels for  
37 recommended EPA and DHA intake with 250-500 mg per day being reported commonly as required  
38 to maintain and support cardiac health (e.g. ISSFAL, 2004; EFSA, 2010; Richter et al., 2016). While  
39 fish and seafood are the main sources of these important nutrients, the capture fisheries that  
40 traditionally supplied them are, at best, stagnant or at worst, in decline, and so over 50 % of all fish  
41 and seafood are now farmed (FAO, 2022). Paradoxically, while the growth of aquaculture has ensured  
42 that the demand for fish and seafood from the increasing human population can be met, it has not  
43 been able to ensure the supply of EPA and DHA (Tocher 2015; Tocher et al., 2019). This is because  
44 many farmed fish like Atlantic salmon (*Salmo salar*) themselves also require a dietary supply of EPA  
45 and DHA to ensure maximum growth and optimum health (Tocher 2010, 2015; NRC, 2011). This  
46 was historically supplied in feeds by the inclusion of fishmeal (FM) and fish oil (FO), also derived  
47 from feed fisheries that are similarly at their sustainable limit and unable to supply the increasing  
48 demand from aquaculture (Cottrell et al., 2020; Naylor et al., 2021; Tacon 2020; Tacon et al., 2022).  
49 The high use of FM and FO was an unsustainable practice, which prompted the development of more  
50 sustainable feeds based on raw materials such as plant meals and vegetable oils (Turchini et al., 2011,  
51 2022). However, these ingredients derived from terrestrial agriculture are devoid of n-3 LC-PUFA  
52 and, therefore, their increased use resulted in reduced levels of EPA and DHA in farmed fish, as has  
53 been well documented in salmon (Sprague et al., 2016; Reksten et al., 2022). Lower levels of dietary  
54 EPA and DHA not only impacts human consumers, but also has potential consequences for the health  
55 of the farmed fish themselves (Tocher and Glencross, 2015; Lufti et al., 2022).

56 While the gap between the demand for n-3 LC-PUFA to satisfy human dietary requirements and  
57 the available supply from all sources is clearly a global issue (Salem et al., 2015), it was felt

58 particularly acutely in fish farming, and the constantly increasing demand from the aquaculture  
59 industry was key in prompting the search for alternative sources of EPA and DHA (Tocher et al.,  
60 2019). Two main research directions were developed, both based on the fact that marine microalgae  
61 are the main organisms responsible for the primary production of EPA and DHA. While one line of  
62 research focused on mass cultivation of microalgae, particularly heterotrophic species (Sprague et al.,  
63 2017), another line of research utilised transgenic approaches to combine the trait for n-3 LC-PUFA  
64 production found in microalgae with the trait for the production and accumulation of oil in large  
65 quantities found in oilseed crops (Napier et al., 2015, 2019; Petrie et al., 2020; Napier and Betancor,  
66 2023). The transgenic approach came with the benefits that oilseeds bring as major agricultural  
67 commodity products, with well-established and highly organised infrastructure that supports the  
68 cultivation, harvest, and processing of oilseeds, along with distribution, marketing and utilisation of  
69 the resultant vegetable oils (VO) (Salunkhe et al., 1992). Furthermore, VO had been the main  
70 alternatives to dietary FO as primary lipid sources in sustainable fish feed formulations (Turchini et  
71 al., 2011; Ytrestøyl et al., 2015; Aas et al., 2019, 2022). Finally, while no VO contains LC-PUFA,  
72 several such as false flax *Camelina sativa*, a member of the Brassicaceae family, can be rich in  $\alpha$ -  
73 linolenic acid (18:3n-3) and, thus, potentially suited for genetic modification to promote the  
74 production of EPA and DHA from the precursor form (18:3n-3) (Napier et al., 2015, 2020).

75 Consequently, in recent years, *C. sativa* crops genetically-modified to produce EPA or EPA and  
76 DHA in their seeds were developed (Ruiz-Lopez et al., 2014; Usher et al., 2017), and have been  
77 evaluated extensively as replacements for dietary FO in feeds for Atlantic salmon (Betancor et al.,  
78 2015a,b, 2016a, 2017), gilthead sea bream (*Sparus aurata*) (Betancor et al., 2016b), European sea  
79 bass (*Dicentrarchus labrax*) (Betancor et al., 2021), rainbow trout (*Oncorhynchus mykiss*) (Osmond  
80 et al., 2021) and Atlantic bluefin tuna (*Thunnus thynnus*) (Betancor et al., 2022). Specifically, in  
81 previous studies in Atlantic salmon, oils from 1<sup>st</sup> and 2<sup>nd</sup> iterations of transgenic *Camelina* supplying  
82 either 20 % EPA or 6 % each of EPA and DHA, respectively, were compared initially with “gold  
83 standard” feeds formulated with high FM and FO (Betancor et al., 2015a,b, 2016a), and the 2<sup>nd</sup>

84 iteration oil (EPA+DHA) was tested subsequently in comparison with more commercially-  
85 representative feeds formulated with lower levels of both FM and FO (Betancor et al., 2017). All the  
86 above studies in salmonids and marine fish species showed highly encouraging results, with the oils  
87 from transgenic *Camelina* supporting good fish growth and enabling the deposition and accumulation  
88 of n-3 LC-PUFA in flesh.

89 The success of the transgenic *Camelina* oils in the above-mentioned trials prompted the  
90 development of a third-generation oil that contained almost 28 % of total fatty acids as n-3 LC-PUFA  
91 including of 10.5 % EPA and 9 % DHA, levels greater than those found in many FO (Betancor et al.,  
92 2018). This oil was tested in salmon using feeds that more closely reflected commercial feeds for  
93 salmon, with even lower levels of FM and FO (Betancor et al., 2018). The diet formulated with the  
94 oil from transgenic *Camelina* showed no negative effects on growth, survival or health of the salmon,  
95 and flesh n-3 LC-PUFA levels were more than 2-fold higher compared with those of fish fed the diet  
96 with a commercial-like formulation containing 30 % FM and 5 % FO (Betancor et al. 2018). The data  
97 demonstrated that the oil from the 3<sup>rd</sup>-generation transgenic *Camelina* crop could efficiently supply  
98 EPA and DHA to salmon resulting in flesh n-3 LC-PUFA levels that were similar to those found  
99 routinely in farmed salmon prior to the large-scale replacement of dietary FM and FO (Sprague et al.,  
100 2016). However, all the above trials in salmon were carried out in land-based seawater tanks in  
101 experimental research facilities with smolts grown over a period of up to 12-weeks and to a maximum  
102 size of 500 g.

103 The aim of the present study was to further validate the efficacy of the 3<sup>rd</sup>-generation transgenic  
104 *Camelina* oil as a dietary oil for farmed Atlantic salmon in a trial carried out in seawater pens and  
105 growing fish over a period of 9 months to a market size of greater than 3 kg. Triplicate groups of  
106 Atlantic salmon were fed experimental diets formulated with low FM that declined as dietary oil  
107 content increased as fish and corresponding pellet size increased during the trial. Two feeds were  
108 produced with added oil supplied either by a mixture of rapeseed oil and FO reflecting the current oil  
109 blend used in commercial salmon feeds in the northern hemisphere (Diet COM), or by 100 %

110 transgenic *Camelina* oil (Diet TCO). The impacts of diet on survival, growth performance, feed  
111 efficiency, tissue fatty acid contents and compositions, and flesh quality were assessed.

112

## 113 **2. Materials and Methods**

### 114 *2.1 Ethics statement*

115 All experimental procedures associated with the Atlantic salmon feeding trial were conducted in  
116 compliance with the Animals Scientific Procedures Act 1986 (Home Office Code of Practice. HMSO:  
117 London January 1997) under project licence PPL7007916 “Environmental Regulation of Fish  
118 Physiology” and personal licence number PIL107216B95, and in accordance with EU regulation (EC  
119 Directive 86/609/EEC). In addition, all experimentation performed by the University of Stirling is  
120 subjected to a thorough ethical review process carried out by the Animal Welfare and Ethical Review  
121 Board (AWERB) prior to any work being approved. This involves all projects, irrespective of where  
122 they are carried out, to be submitted to AWERB for approval using detailed Ethical Approval forms  
123 that require all aspects of the experimentation to be described including all animal health and welfare  
124 issues as well as other ethical considerations. The present research was assessed by the AWERB and  
125 passed the ethical review process (Ethical Approval No. AWERB/16-17/83/New ASPA].

### 126 *2.2 Production of oil from transgenic Camelina sativa*

127 Seeds for the third iteration (identified as DHA2015.1, event #39) were grown under Canadian Food  
128 Inspection Agency (CFIA) permit 17-AGQ1-406-CAM at a site in Elm Creek, Manitoba, Canada.  
129 This was managed by AgQuest LLC, as described previously (Han et al., 2020). Seed was harvested  
130 and transferred to an approved facility (POS Bio-Sciences, Saskatoon, Canada) where the oil was  
131 extracted by cold-pressing and solvent (hexane) extraction. The resulting oil was then provided to  
132 BioMar AS for the production of experimental feed.

133

### 134 2.3 Experimental feeds

135 Two isonitrogenous and isolipidic feeds were formulated to satisfy the known requirements of  
136 Atlantic salmon, and produced by vacuum coating extruded base pellets with either a blend of FO  
137 and rapeseed oil (Control/reference, Diet COM) or the high n-3 LC-PUFA *Camelina* oil (Diet TCO)  
138 (Table 1). The initial formulation (fed to 187 g smolt) provided 44 % protein, 28 % lipid and 24  
139 MJ.kg<sup>-1</sup> of energy, and changed as the fish grew to supply 36 % protein, 36 % lipid and 26 MJ.kg<sup>-1</sup>  
140 of energy to fish growing from 1.5 kg to market size. Initially the base pellet contained around 50 %  
141 plant protein sources and 10 % land animal proteins, and low FM that declined from 15 % to 7.5 %  
142 as the proportion of added oil increased from around 24 % to 32 % as the fish and corresponding  
143 pellet size increased during the trial. In addition, the ratio of rapeseed oil to FO in the COM diet  
144 varied from 0.75:1 in the smallest pellet (5 mm, weeks 1-11), to 2:1 in the larger pellet sizes (7 mm,  
145 weeks 12- 23 and 10 mm, weeks 24 to 37). The changes in the protein and oil contents of the feeds,  
146 and the oil blend ratio of the COM diet, as the fish grew reflected the commercial practices current in  
147 most salmon farming, globally. The fatty acid profiles of the diets showed that the total replacement  
148 of the commercial-type dietary oil blend with the transgenic *Camelina* oil resulted in higher  
149 percentages of all n-3 LC-PUFA, including EPA and DHA, in diet TCO compared to diet COM  
150 (Table 2). The proportions of both linoleic acid (18:2n-6) and 18:3n-3 were also higher in the TCO  
151 diet compared with the COM diet, with the overall higher proportions of PUFA in the TCO feed being  
152 balanced by lower proportions of monoenes (Table 2). The feeds were manufactured at the BioMar  
153 Tech Centre (Brande, Denmark).

154

### 155 2.4 Feeding trial

156 The nutritional feeding trial was carried out at the facilities of the Mowi Feeds Trial Unit (FTU),  
157 Ardnish, Lochailort, Scotland from May 2018 to March 2019. Smolts of the Mowi strain of Atlantic  
158 salmon were transferred to the Ardnish FTU in May 2018 and fed standard commercial transfer feed

159 from then until initiation of the feeding trial in June. A total of 900 well-adapted post-smolt Atlantic  
160 salmon (initial weight ~187 g) were distributed randomly into six 5 x 5 m square seawater pens (150  
161 fish per pen) fitted with automatic feeders (Arvo-Tec Oy, Huutokoski, Finland) and uneaten feed  
162 collection systems. The fish were fed with one of the two feeds in triplicate for a total period of 37  
163 weeks starting on 20 June 2018 with the trial terminated on 6 March 2019. During the experiment,  
164 feeds were provided by the automatic feeders at a ration based on size of the fish and water  
165 temperature as per standard feeding tables for the Ardnish FTU. The actual feed supplied was the  
166 ration + 5 %, to ensure feeding to satiation. Feeds were distributed to the fish twice daily (8.15 – 9.15  
167 am and 2.00 – 3.00 pm) with uneaten feed collected 30 min later and accurate feed intake calculated.  
168 Fish were monitored at feeding to ensure normal feeding behaviour. Growth was determined by  
169 weighing all the fish in the trial pens as appropriate time points including changes in pellet size.  
170 Mortalities were collected daily and examined for any signs of ill health. In the initial 2 weeks after  
171 stocking, mortalities were replaced from the same stock fish to maintain numbers at 150/pen but, after  
172 2 weeks, mortalities were not replaced. Water temperature, salinity, clarity and dissolved oxygen  
173 were monitored daily for the duration of the experiment and can be found in Supplementary Figure  
174 1.

175

## 176 *2.5 Sample collection*

177 At the termination of the nutritional trial, fish were starved for 24 h prior to sampling. All fish were  
178 measured (wet weight and fork length) after anaesthesia with tricaine methanesulphonate (MS222  
179 compound; Merck, Darmstadt, Germany) as per the standard protocol at the Ardnish FTU. A total of  
180 twelve fish per pen were killed by an overdose of MS222 ( $> 150 \text{ mg.l}^{-1}$ ). Four whole fish per pen  
181 were collected onto dry ice and frozen immediately as two pooled samples of two fish per pen ( $n = 6$   
182 per diet) for biochemical analysis (proximate and fatty acid compositions). A second batch of four  
183 fish from each pen were specifically selected to be most representative of harvest-size (~3.5 kg) and  
184 immediately filleted with the fillets from the right side labelled, bagged and taken immediately on ice  
185 for flesh quality analyses (Xelect, St. Andrews, Scotland). A further batch of four fish per pen were



186 used for tissue biochemical analyses with the tissues collected being flesh (Norwegian Quality Cut,  
187 NQC), liver, intestine (pyloric caeca), gills, eyes and brain. The tissue samples were collected as two  
188 pools of two fish per pen (n = 6 per diet), with samples placed in 10 ml plastic tubes and immediately  
189 frozen in liquid nitrogen.

190

## 191 *2.6 Calculations*

192 Biometric parameters were calculated using the following equations:

193 Feed conversion ratio (FCR) = feed (dry weight) consumed / weight gain (wet weight).

194 Fulton's condition factor (k) =  $100 * (W/L^3)$ , where W is the fish weight (g) and L is the total length  
195 (cm).

196 Hepato-somatic index (HSI) =  $(LW/W) * 100$ , where LW is the liver weight and W is the somatic  
197 weight.

198 Specific growth rate (SGR) =  $100 * (\ln W_t - \ln W_o) * D^{-1}$ , where  $W_o$  and  $W_t$  are the initial and end  
199 weights (tanks means, n = 3) of the fish in a specific period, respectively, and D represents the number  
200 of feeding days.

201 Thermal growth coefficient (TGC) =  $1000 * [(W_t^{1/3}) - (W_o^{1/3})] / °D$ , where  $W_o$  and  $W_t$  are the  
202 initial and end weights (tanks means, n = 3) of the fish in a specific period, respectively, and °D  
203 represents degree-days, the sum of daily temperatures in °C in the specific period (or duration in days  
204 x average temperature in period).

205 Viscero-somatic index (VSI) =  $(VW/W) * 100$ , where VW is the weight of the viscera (without liver)  
206 and W is the somatic weight.

207 Weight gain (WG, g) =  $W_t - W_o$ , where  $W_o$  and  $W_t$  are the initial and end weights (tanks means, n =  
208 3) of the fish in a specific period.

209

## 210 *2.7 Proximate compositions of whole fish*

211 Pooled whole fish (Robot Coupe R23 Vertical Food Processor; Robot-Coupe, Vincennes, France)  
212 and salmon flesh samples (NQC) (Robot Coupe Blixer® 4 V.V.) were homogenised before  
213 determination of proximate composition in samples of the resultant pates according to standard  
214 procedures (AOAC, 2000). Protein contents were determined by measuring nitrogen content (N x  
215 6.25) using automated Kjeldahl analysis (Tecator Kjeltex Auto 1030 Analyzer, Foss, Warrington,  
216 UK), while lipid contents were determined gravimetrically after extraction using the Soxhlet method  
217 (Tecator Soxtec system 2050 Auto Extraction apparatus). Moisture contents were obtained after  
218 drying in an oven at 110 °C for 24 h, while ash contents were determined by incinerating the samples  
219 in a muffle furnace at 600 °C for 20 h.

220

## 221 *2.8 Lipid content and fatty acid composition*

222 Total lipid was extracted from ground feeds, homogenised whole fish and flesh (NQC), and  
223 homogenates of liver, intestine (pyloric caeca), gill, brain and eye (MX blender; Waring, USA)  
224 prepared from the two pools of two fish per tank (n = 6 per diet) according to the method of Folch et  
225 al. (1957). Briefly, approximately 1 g samples of experimental material were homogenised in 10  
226 volumes of ice-cold chloroform/methanol (2:1, v/v) containing 0.01 % (w/v) butylated  
227 hydroxytoluene (BHT) as antioxidant using an Ultra-Turrax tissue disrupter (Fisher Scientific,  
228 Loughborough, UK), with content determined gravimetrically. Acid-catalysed transesterification at  
229 50 °C for 16 h was used to prepare fatty acid methyl esters (FAME) from total lipid (Christie, 2003).  
230 The FAME were extracted and purified as described previously (Tocher and Harvie, 1988) and  
231 quantified by gas chromatography in a Fisons GC-8160 (Thermo Scientific, Hemel Hempstead, UK)  
232 equipped with a 30 m × 0.32 mm internal diameter × 0.25 µm ZB-wax column (Phenomenex,  
233 Macclesfield, UK), on-column injector, and a flame ionisation detector. Hydrogen was used as carrier  
234 gas with an initial oven thermal gradient from 50 °C to 150 °C at 40 °C / min, and to a final

235 temperature of 230 °C at 2 °C / min. Individual FAME were identified by comparison with a standard  
236 mixture (Restek 20-FAME Marine Oil Standard; Thames Restek UK Ltd.) and by reference to  
237 published data (Tocher and Harvie, 1988). The GC data were collected and processed using  
238 Chromcard for Windows (version 1.19; Thermoquest Italia S.p.A.).

239

## 240 *2.9 Flesh quality analyses*

241 Fillets for flesh quality analyses were delivered on ice on the day of slaughter. At day three post-  
242 slaughter all fillets (n = 24, 12 per diet) were analysed for colour and gaping, then deboned and  
243 analysed for texture. To determine flesh colour, all fillets were photographed with a 10-megapixel  
244 camera (Canon PowerShot G12) together with a SalmoFan colour scale ruler (Roche, Welwyn  
245 Garden City, UK) and a white reference as a white-colour balance for analysis in ImageJ. Colour was  
246 determined by comparing with the SalmoFan Lineal colour scale ruler (Roche) in three regions of the  
247 fillet above the lateral line: anterior (A, anterior to the dorsal fin), middle (B, below the dorsal fin),  
248 and posterior (C, tail area). For each area, a colour- numbered score from the SalmoFan was assigned.  
249 The scores from the SalmoFan ruler ranged from 20 to 34. Gaping was assessed on a 5-point scale,  
250 but was found to be minimal with only one individual showing very minor gapes. All measurements  
251 were assessed independently by two people and the mean score calculated.

252 Mechanical texture analysis was performed using a TA.XTplus texture analyser (Stable Micro  
253 Systems, Godalming, UK). Firmness measurements were made using a Warner Bratzler blade, which  
254 is a blunt blade of 3 mm thickness with a V-shape notch in the cutting surface. Tensile strength was  
255 measured by mounting the sample on a Pizza Tension rig and using the skin to maintain good grip.  
256 The skin was cut with scissors between the mounts and the sample was then pulled apart while  
257 measuring the force required to do so. All test samples were cut from standardised locations on the  
258 fillet and were cut and trimmed using measured moulds to ensure maximum sample accuracy (4 x 4  
259 x 2 cm blocks for firmness, and 4 x 8 x 2 cm for tensile strength). Both measurements were performed  
260 in duplicate and were analysed by calculating the area under the force/distance curve generated. The

261 resulting value was expressed as mJ of work required to perform the standardised test movement.  
262 Full technical details of the procedure are reported in Ashton et al. (2010).

263

#### 264 *2.10 Carotenoid analysis*

265 Carotenoid contents of flesh (NQC) were determined using a modification of the method of Barua et  
266 al. (1993). Briefly, samples of approximately 1 g of homogenised NQC (see above) were added to 10  
267 mL ethanol/ethyl acetate (1:1, by volume) and thoroughly blended (Ultra-Turrax tissue disrupter;  
268 Fisher Scientific) before being centrifuged at 1000 x g for 5 min. The supernatant was collected into  
269 a clean glass tube and the pellet homogenised and centrifuged twice more, firstly in 5 mL ethyl acetate  
270 then 5mL isohexane. The combined supernatants were dried at room temperature under a stream of  
271 nitrogen and desiccated overnight *in vacuo* before being resuspended in 2 mL isohexane. Samples  
272 were analysed by HPLC on an Ultimate 300 UHPLC system (Thermo Scientific) equipped with a 50  
273 x 3 mm, 1.7  $\mu$  silica column (Synchronis; Thermo Scientific), using an isocratic solvent system  
274 consisting of isohexane/acetone/isopropanol (82:16:2, by volume) at a flow rate of 0.5 mL.min<sup>-1</sup> with  
275 detection at a wavelength of 470 nm. Astaxanthin and other carotenoids were quantified using an  
276 external standard of astaxanthin obtained from DSM (Heerlen, Netherlands).

277

#### 278 *2.11 Statistical analysis*

279 Data were presented as means  $\pm$  SD with n = 3 for fish performance data (Table 3), or n = 6 for  
280 biochemical analyses data (Tables 4-7), while flesh quality data were presented as means  $\pm$  SEM with  
281 n = 12. Percentage data were subjected to arcsin square-root transformation prior to statistical  
282 analyses, and data were tested for normality and homogeneity of variances with Levene's test prior  
283 to nested one-way analysis of variance (ANOVA) with the factor "pen" nested into "treatment"  
284 followed by a Tukey and post-hoc test to determine significant differences for multiple comparisons.  
285 All statistical analyses were performed using SPSS software (IBM SPSS Statistics 23; SPSS Inc.,

286 Chicago, IL, USA) except for the flesh quality analyses, including Pearson Correlation, that were  
287 performed in R. For all data, a P-value < 0.05 was considered significant.

288

### 289 **3. Results**

#### 290 *3.1 Fish growth performance and feed efficiency*

291 There were almost no significant differences observed in any of the growth, biometric or feed  
292 efficiency parameters evaluated at the end of the feeding trial between the fish fed the COM and TCO  
293 diets (Table 3). Overall mortality during the trial was low at around 5 % and not related to the feeds.  
294 While the average size of fish fed the TCO diet was just over 3.1 kg compared to 3.6 kg for the  
295 average size of fish fed the COM diet, the range of fish sizes obtained, especially in pens fed the  
296 COM diet, meant that this difference was not statistically significant (P = 0.0555) (Table 3).  
297 Furthermore, there were no differences in weight gain and TGC at the end of the trial. While there  
298 was also no difference in VSI, HSI was slightly, but significantly, higher in the fish fed the TCO diet  
299 at the end of the trial.

300 In contrast to the overall trial results, significant differences in final weights, weight gain, SGR  
301 and TGC between fish fed the COM and TCO diets were observed in the intermediate phase of the  
302 trial from approximately 850 g up to around 2 kg (Table 3). However, other than condition factor (k)  
303 that was significantly higher in the COM fish, there were no significant differences in any measured  
304 parameter between fish fed the COM and TCO diets in the latter phase of the trial. In addition, there  
305 were no significant differences in feed intake or feed efficiency as measured by FCR between the fish  
306 fed the two diets at any stage of the trial.

307

#### 308 *3.2 Proximate compositions of whole fish and muscle/flesh*

309 There were no significant differences in the protein, lipid, ash and moisture contents of whole fish  
310 between salmon fed the COM and TCO diets (Table 4). However, the proportion of total lipid in flesh  
311 of salmon fed the TCO diet was slightly, but significantly, lower than the proportion of lipid in flesh

312 of fish fed the COM diet, while moisture contents were higher (Table 5). Diet had no effect the  
313 proportions of protein or ash in the salmon flesh.

314

### 315 *3.3 Fatty acid compositions of whole fish and muscle/flesh*

316 In percentage terms, total lipid of whole fish of salmon fed the TCO diet showed significantly  
317 increased proportions of both n-3 and n-6 PUFA and lower proportions of saturates and, especially,  
318 monoenes compared to fish fed the COM diet (Table 4). Percentages of all individual saturated and  
319 monounsaturated fatty acids fatty acids were reduced, other than 20:1n-9, while percentages of all  
320 individual n-6 PUFA were increased. Total n-3 LC-PUFA were increased by almost 2.3-fold, with  
321 EPA, DHA, 20:4n-3 and 22:5n-3 increased by 2.1-, 1.8-, 2.9- and 4.1-fold, respectively. In addition,  
322 the proportions of 18:3n-3, 18:2n-6 and arachidonic acid (20:4n-6) were increased 2.6-, 1.4- and 6.3-  
323 fold in fish fed the TCO diet compared to fish fed the COM diet. The same significant trends in fatty  
324 acid contents were also apparent when reported in mg fatty acids per 100g of fish, absolute terms that  
325 also reflected lipid content. These data showed that fish fed the TCO diet contained almost 3.6 g and  
326 2.2 g of total n-3 LC-PUFA and EPA+DHA, respectively, compared to just under 1.7 g and 1.2 g of  
327 total n-3 LC-PUFA and EPA+DHA, respectively, in fish fed the COM diet (Table 4).

328 Similarly, in percentage terms, total lipid of flesh of fish fed diet TCO also showed significantly  
329 increased proportions of n-3 and n-6 PUFA and lower proportions of monoenes compared to fish fed  
330 the COM diet (Table 5). More specifically, total n-3 LC-PUFA were increased 2.1-fold, with EPA,  
331 DHA, 20:4n-3 and 22:5n-3 increased by 1.9-, 1.6-, 3.7- and 3.6-fold, respectively, while proportions  
332 of 18:3n-3, 18:2n-6 and 20:4n-6 increased 2.6-, 1.4- and 5.5-fold in flesh of salmon fed the TCO diet  
333 compared to fish fed the COM diet. Both the EPA:DHA and n-3:n-6 PUFA ratios increased in flesh  
334 of salmon fed TCO compared to fish fed COM. The key data with respect to human consumers  
335 showed that, in absolute terms (g fatty acids per 100g of flesh), the flesh of salmon fed the TCO diet  
336 contained 2.7 g of n-3 LC-PUFA including 0.7 g of EPA, 0.9 g of DHA and 1.6 g of EPA+DHA.  
337 These data were all significantly higher than the equivalent data in flesh of fish fed the COM diet that

338 delivered 1.45 g of n-3 LC-PUFA including 0.4, 0.65 and just under 1.1 g of EPA, DHA and  
339 EPA+DHA, respectively (Table 5).

340

#### 341 *3.4 Lipid contents and fatty acid compositions of tissues*

342 The lipid content of liver of salmon fed the TCO diet was significantly higher than the liver lipid  
343 content of fish fed the COM diet (Supplementary Table 1). In contrast, diet had no significant effects  
344 on the lipid contents of intestine (pyloric caeca) and gill (Supplementary Table 1), or brain and eye  
345 (Supplementary Table 2).

346 As reported above for whole fish and flesh, the proportions of total n-3 and total n-6 PUFA in almost  
347 all the tissues (liver, intestine, gill, and eye) were higher in fish fed the TCO diet compared to fish  
348 fed the COM diet, while the proportions of monoenes were significantly lower, and diet had no effect  
349 on the proportions of total saturated fatty acids (Supplementary Tables 1 and 2). It was clear that, of  
350 all the tissues, the fatty acid composition of brain was least affected by diet (Supplementary Table 2).  
351 However, while total n-3 LC-PUFA was significantly higher in salmon fed diet TCO compared to  
352 fish fed diet COM in all tissues except brain, this was largely due to increased proportions of EPA,  
353 20:4n-3 and 22:5n-3 while the proportions of DHA were only higher in gill, but unaffected by diet in  
354 intestine, brain and eye (tissues with the highest DHA contents) or even lower as in liver  
355 (Supplementary Tables 1 and 2).

356

#### 357 *3.5 Flesh quality*

358 A summary of the trait values across all flesh quality comparisons are presented in Table 6. On  
359 average, measurements for both firmness and tensile strength were numerically higher for the flesh  
360 of salmon fed the TCO diet in comparison to flesh of fish fed the COM diet 1 (Table 6). However,  
361 these differences in firmness and tensile strength were not statistically significant (ANOVA:  $F =$   
362  $0.206$ ,  $p = 0.66$  and  $F = 0.617$ ,  $P = 0.44$ , respectively). Interestingly, a positive significant correlation  
363 was found between tensile and firmness measurements (Pearson correlation:  $r = 0.47$ ,  $P = 0.02$ ).

364 While there was no noteworthy gaping observed among fillets, flesh colour of salmon fed the COM  
365 diet was significantly paler than those fed the TCO diet (ANOVA:  $F = 4.286$ ,  $P = 0.05$ ). However,  
366 analyses of flesh (NQC) showed that there were no significant differences in total contents or  
367 compositions ( $> 95\%$  astaxanthin,  $3\%$  astacene and  $< 2\%$  lutein) of carotenoids between fish fed  
368 the two diets.

369

## 370 **Discussion**

371 It is over 50 years since the pioneering work of Dyerberg and Bang first reported on the effects of a  
372 marine-based diet (Bang et al., 1971), which subsequently led to the discovery of the importance of  
373 the omega-3 LC-PUFA in human nutrition (Dyerberg et al., 1978). Therefore, the role of a marine  
374 diet and fish in providing the human population with the health-beneficial fatty acids, EPA and DHA,  
375 was known from the very beginning of “omega-3” research. However, it is now well established that  
376 there is a large gap between the demand for EPA and DHA required for human health and their supply  
377 from both traditional (capture fisheries) and modern (aquaculture) sources (Tocher et al., 2019).  
378 Bridging this gap is a very real problem with many human populations around the globe shown to  
379 have very low levels of EPA and DHA in the blood (Stark et al., 2016). It is within this context that  
380 present study is placed.

381 In the present trial, salmon were grown in seawater pens for 9 months during which time they  
382 grew from around 180 g to  $> 3$  kg. There were no significant differences in final average weight,  
383 weight gain, SGR or TGC between fish fed the COM and TCO diets. However, it is clear that salmon  
384 fed the TCO diet were on average smaller than fish fed the COM diet, and this was only not significant  
385 because of the large range in weights obtained in these ungraded populations. The range in fish size  
386 was more than 3-fold greater in fish fed the COM diet and due to the presence of some very large fish  
387 in that group, rather than the presence of smaller fish in the TCO group. This size difference between  
388 the dietary treatments stemmed from lower growth rate in fish fed the TCO diet during the  
389 second/intermediate phase of the trial, when significantly lower final weights, weight gain, SGR and



390 TGC were recorded. Although the difference in final weight increased in the final phase of the trial,  
391 growth in that phase was not significantly different between the dietary treatments. The difference in  
392 final weights in the present trial was not observed in the previous trial where salmon were fed the  
393 same oil used in the TCO diet with fish grown from 130 g to 400 g (Betancor et al., 2018), or in any  
394 of the earlier trials feeding salmon the oils obtained from previous iterations of GM *Camelina* and  
395 growing fish to 200 g (Betancor et al., 2015a), 400 g (Betancor et al., 2017), or 500 g (Betancor et  
396 al., 2016a).

397 Fish fed the TCO diet showed numerically lower FI (in g/fish/day) throughout the trial, being 5 -  
398 6 % lower in the first and last phases of the trial, but over 12 % lower in the intermediate phase, and  
399 8 % lower overall. Although none of these differences in FI were statistically significant, it is highly  
400 likely that lower FI was the reason for differences observed in final weights and weight gains of fish  
401 fed the TCO diet compared to fish fed the COM diet. Supporting this conclusion, FCR was not  
402 significantly different between fish fed the COM and TCO diets at any point in the trial suggesting  
403 that there were no differences in intermediary metabolism and/or metabolic performance of the diets,  
404 and that both feeds were utilised with the same efficiency. However, in a trial in European sea bass,  
405 FI and SGR were significantly lower in the first month of the 4-month trial in fish fed the TCO oil  
406 compared to fish fed a control FO diet (Betancor et al., 2021). It was speculated that the initially lower  
407 FI, which affected growth, was possibly due to reduced palatability, as it was overcome in the later  
408 phases of the trial. *Camelina sativa* is a Brassicaceae and, as such, contains glucosinolates (Berhow  
409 et al., 2013) that are known to cause the bitter/sharp taste of many cruciferous vegetables (Clarke,  
410 2010), and studies investigating feed ingredients rich in glucosinolates have shown they negatively  
411 affect palatability and reduce growth of fish (Francis et al., 2001), which may be exacerbated in feeds  
412 with limited inclusion of fishmeal. The Camelina oil used in the present study was equivalent to a  
413 “virgin” oil and received no processing post-extraction, which could cause palatability issues, but  
414 also suggests that these could be alleviated by reducing the glucosinolate level and/or the inclusion  
415 of feed additives or palatants including palatability enhancers and feed attractants (Pilmer et al.,

416 2022). This would be part of the normal process of commercial optimisation of feed formulations  
417 containing the Camelina oil, securing the best inclusion levels to achieve optimal diet performance.  
418 The reduction of glucosinolates in *C. sativa* using a biotechnological approach would be one option  
419 and a potential future target (Nour-Eldin et al., 2017). Overall, therefore, the difference in final  
420 weights reported in the present trial was likely due to the crude nature of the oil used in the TCO diet,  
421 which impacted the palatability and intake of the feed, particularly in the first part of the trial. Several  
422 options are available to mitigate this issue since it is commonly encountered in the replacement of  
423 marine ingredients with plant ingredients (Francis et al., 2001; Nagel et al., 2012).

424 While diet had no impact on the biochemical composition of whole fish, the flesh of salmon fed  
425 the COM diet had a lipid content of almost 16.5 % whereas the lipid content of flesh of fish fed the  
426 TCO diet TCO was significantly lower at 14.5 % of wet weight. As would be expected, the lower  
427 lipid content of flesh of salmon fed the TCO diet was accompanied by increased moisture content.  
428 The “target” value for flesh lipid content of farmed salmon in Scotland is around 16-17 % based on  
429 retailer and quality label specifications (e.g. Label Rouge,  $\leq 16\%$ ), so the COM fish were perfectly  
430 in this range. As the tissue that arguably represents the largest fat store in salmon, the lower flesh  
431 lipid level in fish the salmon fed TCO may reflect the lower FI and, consequently, energy intake of  
432 fish fed this diet resulting in lower lipid deposition and accumulation in flesh and, possibly, lower  
433 body weight. Lower lipid contents in whole fish and flesh of smaller (400 g) Atlantic salmon fed the  
434 TCO diet compared to fish fed a COM diet were reported previously (Betancor et al., 2018). In that  
435 study, it was suggested that the lower body and flesh lipid contents could be associated with the higher  
436 EPA and DHA contents of the TCO diet compared to the COM diet as these n-3 LC-PUFA are known  
437 to have anti-adipogenic effects in mammals (Dentin et al., 2005). In addition, microarray analysis  
438 revealed that the lipogenic gene, *acs11* (acyl-CoA synthetase long chain family member 1) was down-  
439 regulated in fish fed TCO compared to fish fed COM, possibly indicating reduced lipogenesis, and  
440 the *lpl* (lipoprotein lipase) gene was also downregulated in TCO-fed fish, which could be considered  
441 consistent with lower flesh lipid levels (Betancor et al., 2018). However, in trials in similarly smaller

442 salmon fed the oils obtained from earlier iterations of GM *Camelina* including an EPA-only oil  
443 (Betancor et al., 2015a) or an oil with 6 % each of EPA+DHA (Betancor et al., 2016a, 2017), no  
444 significant impacts on flesh lipid contents were observed.

445 The small, but significant, difference in lipid content in flesh discussed above was generally not  
446 reflected in any of the flesh quality parameters measured, which showed no difference in firmness,  
447 tensile strength or gaping between dietary treatments. However, there was a significant difference in  
448 flesh colour with fish fed the TCO diet showing higher average colour in the Roche SalmoFan Lineal  
449 colour scale. The effects of the TCO diet on flesh colour and carotenoid content had not been  
450 measured previously in our earlier trials as they were performed in land-based tanks and fish were  
451 still too small at the end of the feeding trials for impacts on pigmentation to be meaningful (Betancor  
452 et al., 2016, 2018). While this result was interesting, the underpinning reason was unclear as the flesh  
453 carotenoid content ( $\text{mg}\cdot\text{kg}^{-1}$ ) was not significantly different between fish fed the two diets, and the  
454 amount of carotenoid relative to flesh lipid level was also very similar. However, a similar result was  
455 reported in salmon fed diets containing oil (“Aquaterra”) from GM Canola and grown in sea pens to  
456 1.5 kg (Ruyter et al., 2022). In that study, red colour intensity was significantly higher in flesh of  
457 salmon fed a diet with 50 % of oil supplied by the GM Canola (replacing the VO components of the  
458 diet) compared to control fish without GM Canola, while flesh astaxanthin levels were not  
459 significantly different between dietary groups (Ruyter et al., 2022). Similarly, in another study with  
460 salmon grown from 700 g to over 4.5 kg on feeds containing increasing levels of GM Canola, no  
461 differences were reported in flesh astaxanthin and total carotenoid concentrations (Hatlen et al.,  
462 2022).

463 In contrast to flesh, lipid contents were unaffected by diet in most tissues other than liver where  
464 lipid content was over 50 % higher in salmon fed the TCO diet compared to fish fed the COM diet.  
465 This was consistent with the slightly, but significantly, higher HSI of fish fed TCO. Increased liver  
466 lipid levels are often regarded as reflecting a metabolic disturbance, potentially as a result of some  
467 lipid or nutrient imbalance. The high proportion of 18:3n-3 that would, arguably, be more likely to

468 be esterified into tissue lipids, combined with lower proportions of monoenoic fatty acids that would  
469 be more likely to promote fatty acid oxidation, may represent a metabolic imbalance that could lead  
470 to accumulation of lipid in liver of fish fed TCO compared to fish fed COM. In our previous trial  
471 with smaller fish, liver lipid contents were not significantly different between fish fed the TCO and  
472 COM diets (Betancor et al., 2018) and, similarly, the oil from GM *Camelina* containing 6 % each of  
473 EPA+DHA had no impact on liver lipid contents (Betancor et al., 2016a, 2017). In contrast, salmon  
474 fed the EPA-only oil (Betancor et al., 2015a) showed higher whole body and liver lipid contents  
475 compared to salmon fed the control FO diet (Betancor et al., 2015a).

476 The main driver for the development of new sources of EPA and DHA was to increase the  
477 availability of these critical EFA to the human population and, therefore, arguably, the most important  
478 data in the present study are those showing the impact of the TCO diet on the fatty acid compositions  
479 of the salmon. Thus, it was noteworthy that the levels of all n-3 LC-PUFA in whole fish increased  
480 considerably, and total n-3 LC-PUFA were over 2-fold greater in salmon fed the TCO diet compared  
481 to fish fed the COM diet. However, nutritional quality of the salmon in terms of EPA and DHA is  
482 based on the composition of the edible portion, flesh/muscle, and the present study showed that, in  
483 relative terms, the proportion of total n-3 LC-PUFA of flesh also more than doubled from 9.7 % of  
484 total fatty acids (TFA) in fish fed COM to 20.9 % in fish fed the TCO diet. In absolute terms, EPA,  
485 DHA, EPA+DHA and total n-3 LC-PUFA increased from 0.42, 0.65, 1.08 and 1.45 g/100g<sup>-1</sup> flesh,  
486 respectively, in fish fed the COM diet to 0.70, 0.91, 1.61 and 2.71 g/100g<sup>-1</sup> flesh, respectively, in  
487 salmon fed the TCO diet. In consequence, a standard 140 g portion of flesh of salmon fed the TCO  
488 diet would deliver almost 2.3 g of EPA+DHA and 3.8 g of total n-3 LC-PUFA (EPA, DHA, 22:5n-  
489 3, 20:4-3 and 20:3n-3) and, therefore, a single 140 g portion of salmon fed TCO would deliver 67 %  
490 of the weekly requirement level of EPA and DHA (3.5 g; 500 mg daily) recommended by many  
491 health agencies (ISSFAL, 2004; EFSA, 2010). In contrast, a 140 g portion of the salmon fed the COM  
492 diet, reflecting current farming practices, would deliver 1.5 g EPA+ DHA, similar to the level reported  
493 for commercial Scottish salmon in 2016 (Sprague et al., 2016), and 0.8 g lower than a portion of

494 salmon fed TCO, and less than half the recommended weekly intake (ISSFAL, 2004; EFSA, 2010).  
495 Salmon fed the TCO diet, and with a similar flesh lipid content (16.5 %) to the COM fish, could  
496 arguably have an even higher EPA+DHA content at around 2.6 g per 140 g portion, representing 75  
497 % of the recommended weekly intake.

498 Therefore, replacing entirely (100 %) the current added oil, blends of FO and rapeseed oil, used  
499 in commercial salmon farming with oil from transgenic *Camelina* in feed for salmon during the  
500 seawater growth phase to market size had a major beneficial impact on the nutritional quality of the  
501 flesh for human consumers in terms of substantially increased n-3 LC-PUFA including, importantly,  
502 both EPA and DHA. Two other GM crops have been developed, both from rapeseed/Canola,  
503 producing oils that are either relatively rich in DHA (“Aquaterra<sup>®</sup>”, ~ 9 % DHA and 0.5 % EPA;  
504 Davis and Devine, 2023) or EPA (“Latitude”, ~ 7 % EPA and 1 % DHA). In consequence,  
505 incorporating Aquaterra into feed increased predominantly DHA levels in juvenile (Ruyter et al.,  
506 2019), on-growing (Ruyter et al., 2022) and harvest-size (Hatlen et al., 2022) Atlantic salmon reared  
507 in seawater, while incorporation of Latitude into feed increased predominantly EPA level in rainbow  
508 trout reared to market size in freshwater (Hong et al., 2022). In addition to the oils from GM crops,  
509 the microalgal oil, “Veramaris” that has high levels of both DHA and EPA (almost 40 % and 16 %  
510 of TFA, respectively), has also been used to replace the FO component of FO/VO blends to  
511 successfully maintain EPA and increase DHA levels in flesh of Atlantic salmon grown to market size  
512 (3 kg) (Santigosa et al., 2023). Incorporating Veramaris into feeds also improved flesh DHA levels  
513 in trials where it was used to replace the FO component of FO/VO blends in both rainbow trout  
514 (Santigosa et al., 2020) and gilthead seabream (Santigosa et al., 2021). Although Veramaris has the  
515 highest levels of EPA+DHA of all the new sources, it depends on fermentation, which currently limits  
516 supply and is also costly, and so the algal oil is expensive and likely to have a much higher cost per  
517 percentage point of EPA+DHA than GM crops. Thus, while not containing as high levels as  
518 Veramaris, the oil from transgenic *Camelina* used in the TCO feed was designed to have an  
519 EPA+DHA content and composition similar to the FO traditionally used in salmon farming, with

520 higher levels of EPA+DHA (~ 20 %) and a better ratio of EPA and DHA (~ 1 : 1) than the GM Canola  
521 oils and, therefore, it perhaps represents a unique balanced solution among all the alternatives.  
522 However, it is important to stress that all the oils from GM crops, as well as algal oils, have key roles  
523 to play in improving the health and nutritional quality of farmed salmon (Tocher et al., 2019),  
524 ensuring they can again supply the high levels of EPA and DHA farmed salmon once did before  
525 large-scale replacement of marine ingredients (Sprague et al., 2016; Refksten et al., 2022). Reflecting  
526 the current very high interest in new sources of omega-3 LC-PUFA, the use of oils from GM crops  
527 in aquafeeds received a boost recently when the Aquaterra<sup>®</sup> GM Canola oil was approved by the  
528 Norwegian Food Safety Authority for use in fish feed applications in Norway (Aquaterra, 2023).  
529 Furthermore, the fact that the beneficial impacts of oils from GM *Camelina* in increasing n-3 LC-  
530 PUFA levels in flesh observed in earlier trials in smaller salmon (Betancor et al., 2016, 2018)  
531 translated to market size fish, argues that this would likely extend to other farmed fish species such  
532 as gilthead seabream (Betancor et al., 2016b), European sea bass (Betancor et al., 2021), rainbow  
533 trout (Osmond et al., 2021) and Atlantic bluefin tuna (Betancor et al., 2022), where oils from GM  
534 *Camelina* increased flesh n-3 LC-PUFA levels in trials with small/less than market size fish. This  
535 suggested that similar benefits would accrue in these species if fed oil from GM *Camelina* during  
536 grow out to market size, providing farmed fish in general with the levels of EPA+DHA expected  
537 traditionally of wild capture fish and seafood (Tocher et al. 2019).

538         Increasing the dietary levels of EPA and DHA in feeds for farmed fish not only benefits human  
539 consumers, but also the health and welfare of the farmed fish themselves. Recent studies have shown  
540 that the dietary level of EPA+DHA to support optimal health in salmon is much higher than the level  
541 of 0.5 % of feed reported commonly (Tocher, 2010; NRC, 2011). One study suggested that salmon  
542 in seawater required a minimum level of EPA+DHA of at least 2.7 % of TFA (~ 1 % of diet) based  
543 largely on growth (Rosenlund et al., 2016), and other studies studies suggested that a level of  
544 EPA+DHA of at least 1.6 % of diet was required to ensure growth and maintain robustness of farmed  
545 salmon in seawater (Sissener et al., 2016; Bou et al., 2017). Most recently, however, a further trial

546 indicated that a level of EPA+DHA of 3.5 % of diet improved health and welfare of salmon in  
547 challenging, but essentially normal, farming conditions (Lufti et al., 2022). Current levels of  
548 EPA+DHA used in salmon farming vary from < 2 % (Chile) to 3.5 % of diet (Faroes), with Norway  
549 possibly transitioning between 2.0 and 2.5 %. In the present study, the COM diet was formulated to  
550 supply EPA+DHA at 2.5 % of diet (almost 7 % of TFA), which, as indicated earlier, was the standard  
551 level in feeds for Scottish salmon at the time of the trial, while the TCO diet supplied EPA+DHA at  
552 4 % of diet (11 % of TFA), above the highest levels tested in salmon in any of the earlier studies.

553 The above highlights the importance of EPA and DHA to both fish health and nutritional quality  
554 of farmed products in not only salmon but, likely, all farmed fish. While finding alternatives to  
555 traditional fish meals as protein sources remains a major driver of research into the feed resources  
556 required to support sustainable salmon farming (Albrektsen et al., 2022), the development of entirely  
557 new, sustainable, and economically-viable sources of EPA and DHA is a challenge that has been, at  
558 least partly, solved. While recovery and recycling of EPA and DHA from fisheries and aquaculture  
559 by-products has increased in recent years, and opportunities likely exist for increased by-product  
560 utilisation and waste prevention, various economic, cultural and technical challenges remain to be  
561 overcome (Hamilton et al., 2020).

562 In conclusion, the current study represents an important step in the validation of oil from an  
563 oilseed crop, *Camelina sativa*, genetically engineered to produce high levels of EPA and DHA in  
564 seeds, as an entirely new, *de novo* source of these health-critical omega-3 LC-PUFA. The present  
565 study was performed in semi-commercial conditions in sea pens in salmon grown for 9 months from  
566 new smolt (~ 180 g) to market size (> 3 kg). Although there was a size difference at harvest, there  
567 were no differences in SGR, FCR or survival between fish fed the TCO or COM diets over the whole  
568 growth period. Nutritional quality in terms of “omega-3” content was substantially improved in fish  
569 fed the TCO, diet with total n-3 LC-PUFA level of flesh more than double that in fish fed the COM  
570 diet. Consequently, a standard 140 g portion of flesh of salmon fed a diet formulated with oil from  
571 transgenic *Camelina* would deliver a dose of EPA+DHA sufficient to cover at least two-thirds of the

572 weekly requirement level recommended by many health agencies, and over one and a half times more  
573 than the level supplied by fish fed the current commercial dietary regime.

574

## 575 **Declaration of Competing Interest**

576 The authors declare no conflict of interest exist.

577

## 578 **Acknowledgements**

579 The authors are grateful to Ed King (Feed Trials Manager, Mowi Feed) and the staff of the Mowi  
580 Ardnish FTU for their excellent technical assistance in fish rearing and sampling. This project was  
581 funded by a UK Biotechnology and Biological Sciences Research Council (BBSRC) Super Follow-  
582 On Funding Award (BB/N022157/1). Additional support from the Institute Strategic Programme  
583 Grant “*Tailoring Plant Metabolism*” BBS/E/C/000I0420 to JAN at Rothamsted Research is  
584 acknowledged.

585

## 586 **References**

- 587 Aas, T. S., Åsgård, T., Ytrestøyl, T., 2022. Utilization of feed resources in the production of Atlantic  
588 salmon (*Salmo salar*) in Norway: An update for 2020. *Aquacult. Rep.* 26, 101316.
- 589 Aas, T.S., Ytrestøyl, T., Åsgård, T., 2019. Utilization of feed resources in the production of Atlantic  
590 salmon (*Salmo salar*) in Norway: an update for 2016. *Aquacult. Rep.* 15, 100216.
- 591 Albrektsen, S., Kortet, R., Skov, P.V., et al., 2022. Future feed resources in sustainable salmonid  
592 production: A review. *Rev. Aquac.* 14, 1790–1812.
- 593 Aquaterra, 2023. [https://aquaterraomega3.com/norway-approves-aquaterra-omega-3-oil-for-use-in-](https://aquaterraomega3.com/norway-approves-aquaterra-omega-3-oil-for-use-in-aquafeed/)  
594 [aquafeed/](https://aquaterraomega3.com/norway-approves-aquaterra-omega-3-oil-for-use-in-aquafeed/). Accessed January 2024.
- 595 Ashton, T.J., Michie, I., Johnston, I.A., 2010. A novel tensile test method to assess texture and gaping  
596 in salmon fillets. *J. Fd. Sci.* 75, S182-S190.
- 597 Association of Official Analytical Chemists, 2000. *Official Methods of Analysis*. Washington, DC:  
598 AOAC.
- 599 Austreng, E., 1978. Digestibility determination in fish using chromic oxide marking and analysis of  
600 contents from different segments of the gastrointestinal tract. *Aquaculture* 13, 265–272.
- 601 Bang, H.O., Dyerberg, J., Nielsen, A.B., 1971. Plasma lipid and lipoprotein pattern in Greenlandic  
602 West-coast Eskimos. *Lancet* 1, 1143–1146.
- 603 Barua, A.B., Kostic, D., Olsen, J.A., 1993. New simplified procedure for the extraction and  
604 simultaneous high-performance liquid chromatographic analysis of retinol, tocopherols and  
605 carotenoids in human serum. *J. Chromatogr.* 617, 257–264.



606 Berhow, M.A., Polat, U., Glinski, J.A., Glensk, M., Vaughn, S.F., Isbell, T., Ayala-Diaz, I., Marek,  
607 L., Gardner, C., 2013. Optimized analysis and quantification of glucosinolates from *Camelina*  
608 *sativa* seeds by reverse-phase liquid chromatography. *Ind. Crop. Prod.* 43, 119–125.

609 Betancor, M.B., Li K., Sprague, M., Bardal, T., Sayanova, O., Usher, S., Han, L., Måsøval, K.,  
610 Torrissen, O., Napier, J.A., Tocher, D.R., Olsen, R.E., 2017. An oil containing EPA and DHA  
611 from transgenic *Camelina sativa* to replace marine fish oil in feeds for Atlantic salmon (*Salmo*  
612 *salar* L.): Effects on intestinal transcriptome, histology, tissue fatty acid profiles and plasma  
613 biochemistry. *PLoS One* 12, e0175415.

614 Betancor, M.B., Li, K., Bucerzan, V.S., Sprague, M., Sayanova, O., Usher, S., Han, L., Norambuena,  
615 F., Torrissen, O., Napier, J.A., Tocher, D.R., Olsen, R.E., 2018. Oil from transgenic *Camelina*  
616 *sativa* containing over 25 % n-3 long-chain polyunsaturated fatty acids as the major lipid source  
617 in feed for Atlantic salmon (*Salmo salar*). *Br. J. Nutr.* 119, 1378-1392.

618 Betancor, M.B., MacEwan, A., Sprague, M., Gong, X., Montero, D., Han, L., Napier, J.A.,  
619 Norambuena, F., Izquierdo, M., Tocher, D.R., 2021. Oils from transgenic *Camelina sativa* as a  
620 source of EPA and DHA in feeds for European sea bass (*Dicentrarchus labrax* L.). *Aquaculture*  
621 530, 735759.

622 Betancor, M.B., Sprague, M., González-Silvera, D., Ortega, A., de la Gándara, F., Gong, X., Napier,  
623 J.A., Tocher, D.R., Mourente, G., 2022. Oils derived from GM crops as sustainable solutions to  
624 the supply of long-chain omega-3 for on-growing Atlantic bluefin tuna (*Thunnus thynnus* L.).  
625 *Fishes* 7, 366.

626 Betancor, M.B., Sprague, M., Montero, D., Usher, S., Sayanova, O., Campbell, P.J., Napier, J.A.,  
627 Caballero, M.J., Izquierdo, M., Tocher, D.R., 2016b. Replacement of marine fish oil with *de novo*  
628 omega-3 oils from transgenic *Camelina sativa* in feeds for gilthead sea bream (*Sparus aurata* L.).  
629 *Lipids* 51, 1171-1191.

630 Betancor, M.B., Sprague, M., Sayanova, O., Usher, S., Campbell, P.J., Napier, J.A., Caballero, M.J.,  
631 Tocher, D.R., 2015b. Evaluation of a high-EPA oil from transgenic *Camelina sativa* in feeds for  
632 Atlantic salmon (*Salmo salar* L.): Effects on tissue fatty acid composition, histology and gene  
633 expression. *Aquaculture* 444, 1-12.

634 Betancor, M.B., Sprague, M., Sayanova, O., Metochis, C., Campbell, P.J., Napier, J.A., Tocher, D.R.,  
635 2016a. Nutritional evaluation of an EPA-DHA oil from transgenic *Camelina sativa* in feeds for  
636 post-smolt Atlantic salmon (*Salmo salar* L.). *PLoS One* 11, e0159934.

637 Betancor, M.B., Sprague, M., Usher, S., Sayanova, O., Campbell, P.J., Napier, J.A., Tocher, D.R.,  
638 2015a. A nutritionally-enhanced oil from transgenic *Camelina sativa* effectively replaces fish oil  
639 as a source of eicosapentaenoic acid for fish. *Sci. Rep.* 5, 8104.

640 Bou, M., Berge, G.M., Baeverfjord, G., Sigholt, T., Østbye, T.-K., Ruyter, B., 2017. Low levels of  
641 very-long-chain n-3 PUFA in Atlantic salmon (*Salmo salar*) diet reduce fish robustness under  
642 challenging conditions in sea cages. *J. Nutr. Sci.* 6, 1–14.

643 Calder, P.C., 2018. Very long-chain n-3 fatty acids and human health: Fact, fiction and the future.  
644 *Proc. Nutr. Soc.* 77, 52–72.

645 Christie, W.W., 2003. *Lipid Analysis*, 3rd ed., pp. 205–224. Bridgwater: The Oily Press.

646 Clarke, D.B., 2010. Glucosinolates, structures and analysis in food. *Anal. Methods* 2, 310–325.

647 Cottrell, R.S., Blanchard, J.L., Halpern, B.S., Metian, M., Froehlich, H.E., 2020. Global adoption of  
648 novel aquaculture feeds could substantially reduce forage fish demand by 2030. *Nature Fd.* 1, 301–  
649 308.

650 Davis, B.A., Devine, M.D., 2023. Evaluation of long-chain omega-3 canola oil on Atlantic salmon  
651 growth, performance, and essential fatty acid tissue accretion across the life cycle: a review.  
652 Aquacult. Internat. <https://doi.org/10.1007/s10499-023-01099-3>

653 Dentin, R., Benhamed, F., Pégrier, J.P., Fougelle, F., Viollet, B., Vaulont, S., Girard, J., Postic, C.,  
654 2005. Polyunsaturated fatty acids suppress glycolytic and lipogenic genes through the inhibition  
655 of ChREBP nuclear protein translocation. *J. Clin. Invest.* 115, 2843–2854.

656 Dyerberg, J., Bang, H.O., Stoffersen, E., Moncada, S., Vane, J.R., 1978. Eicosapentaenoic acid and  
657 prevention of thrombosis and atherosclerosis? *Lancet* 312, 117–119.

658 European Food Safety Authority (EFSA), 2010. Scientific opinion on dietary reference values for fat,  
659 including saturated fatty acids, polyunsaturated fatty acids, monounsaturated fatty acids, trans fatty  
660 acids, and cholesterol. EFSA panel on dietetic products, nutrition and allergies (NDA). *EFSA J.*  
661 8, 1461.

662 FAO, 2022. The State of World Fisheries and Aquaculture 2022. Towards blue transformation, Rome,  
663 FAO.

664 Folch, J., Lees, M., Sloane-Stanley, G.H., 1957. A simple method for the isolation and purification of  
665 total lipides from animal tissues. *J. Biol. Chem.* 226, 497–509.

666 Francis, G., Makkar, H.P., Becker, K., 2001. Antinutritional factors present in plant-derived alternate  
667 fish feed ingredients and their effects in fish. *Aquaculture* 199, 197–227.

668 Hamilton, H.A., Newton, R., Auchterlonie, N.A., Müller, D.B., 2020. Systems approach to quantify  
669 the global omega-3 fatty acid cycle. *Nature Fd.* 1, 59–62.

670 Han, L., Usher, S., Sandgrind, S., Hassall, K., Sayanova, O., Michaelson, L.V., Haslam, R.P., Napier,  
671 J.A., 2020. High level accumulation of EPA and DHA in field-grown transgenic Camelina - a  
672 multi-territory evaluation of TAG accumulation and heterogeneity. *Plant Biotechnol. J.* 18, 2280–  
673 2291.

674 Hatlen, B., Larsson, T., Østbye, T.-K., Romarheim, O.H., Rubio, L.M., Ruyter, B., 2022. Improved  
675 fillet quality in harvest-size Atlantic salmon fed high n-3 canola oil as a DHA-source. *Aquaculture*  
676 560, 738555.

677 Hong, J., Bledsoe, J.W., Overturf, K.E., Lee, S., Iassonova, D., Small, B.C., 2022. Latitude Oil as a  
678 sustainable alternative to dietary fish oil in rainbow trout (*Oncorhynchus mykiss*): effects on fillet  
679 fatty acid profiles, intestinal histology, and plasma biochemistry. *Front. Sustain. Food Syst.* 6,  
680 837628.

681 Innes, J.K., Calder, P.C., 2020. Marine omega-3 (N-3) fatty acids for cardiovascular health: an update  
682 for 2020. *Int. J. Mol. Sci.* 21, 1362.

683 International Society for the Study of Fatty Acids and Lipids (ISSFAL), 2004. Report of the Sub-  
684 Committee on: Recommendations for Intake of Polyunsaturated Fatty Acids in Healthy Adults.  
685 Brighton: ISSFAL.

686 Lutfi, E., Berge, G. M., Bæverfjord, G., Sigholt, T., Bou, M., Larsson, T., Mørkøre, T., Evensen, Ø.,  
687 Sissener, N. H., Rosenlund, G., Sveen, L., Østbye, T. K., Ruyter, B., 2022. Increasing dietary  
688 levels of the n-3 long-chain PUFA, EPA and DHA, improves the growth, welfare, robustness and  
689 fillet quality of Atlantic salmon in sea cages. *Br. J. Nutr.* 129, 1–19.

690 Nagel, F., von Danwitz, A., Tusche, K., Kroeckel, S., van Bussel, C. G. J., Schlachter, M., Adem, H.,  
691 Tressel, R. P., Schulz, C., 2012. Nutritional evaluation of rapeseed protein isolate as fish meal  
692 substitute for juvenile turbot (*Psetta maxima* L.) - Impact on growth performance, body  
693 composition, nutrient digestibility and blood physiology. *Aquaculture*, 356–357, 357–364.

694 Napier, J.A., Betancor, M.B., 2023. Engineering plant-based feedstocks for sustainable aquaculture.  
695 *Curr. Op. Plant Biol.* 71, 102323.

696 Napier, J.A., Haslam, R.P., Olsen, R.E., Tocher, D.R., Betancor, M.B., 2020. Agriculture can help  
697 aquaculture become greener. *Nature Fd.* 1, 680-683.

698 Napier, J.A., Olsen, R.E., Tocher, D.R., 2019. Update on GM canola crops as novel sources of omega-  
699 3 fish oils. *Plant Biotechnol.* 17, 703-705.

700 Napier, J.A., Usher, S., Haslam, R.P., Ruiz-Lopez, N., Sayanova, O., 2015. Transgenic plants as a  
701 sustainable, terrestrial source of fish oils. *Eur. J. Lipid Sci. Technol.* 117, 1317-1324.

702 National Research Council (NRC), 2011. Nutrient Requirements of Fish and Shrimp. The National  
703 Academies Press, Washington DC.

704 Naylor, R.L., Hardy, R.W., Buschmann, A.H., Bush, S.R., Cao, L., Klinger, D.H., Little, D.C.,  
705 Lubchenco, J., Shumway, S.E., Troell, M., 2021. A 20-year retrospective review of global  
706 aquaculture. *Nature* 591, 551–563.

707 Nour-Eldin, H.H., Madsen, S.R., Engelen, S., Jørgensen, M.E., Olsen, C.E., Andersen, J.S.,  
708 Seynnaeve, D., Verhoye, T., Fulawka, R., Denolf, P., Halkier, B.A., 2017. Reduction of  
709 antinutritional glucosinolates in Brassica oilseeds by mutation of genes encoding transporters. *Nat.*  
710 *Biotechnol.* 35, 377–382.

711 Osmond, A.T., Arts, M.T., Bazinet, R.P., Napier, J.A., Han, L., Colombo, S.M., 2021. Transgenic  
712 camelina oil is an effective source of eicosapentaenoic acid and docosahexaenoic acid in diets for  
713 farmed rainbow trout, in terms of growth, tissue fatty acid content, and fillet sensory properties. *J.*  
714 *World Aquacult. Soc.* 52, 961–986.

715 Petrie, J.R., Zhou, X.-R., Leonforte, A., McAllister, J., Shrestha, P., Kennedy, Y., Belide, S., Buzza,  
716 G., Gororo, N., Gas, W., Lester, G.L., Mansour, M.P., Mulder, R., Liu, Q., Tian, L., Silva, C.,  
717 Cogan, N., Nichols, P.D., Green, A., de Feyter, R., Devine, M.D., Singh, S.P., 2020. Development  
718 of a *Brassica napus* (canola) crop containing fish oil-like levels of DHA in the seed oil. *Front.*  
719 *Plant Sci.* 11, 727–741.

720 Pilmer, L.W., Woolley, L.D., Lymbery, A.J., Salini, M., Partridge, G.J., 2022. Using dietary additives  
721 to improve palatability of diets containing single-cell protein from methanotrophic bacteria in  
722 yellowtail kingfish (*Seriola lalandi*) diets. *Aquacult. Res.* 53, 5006–5017.

723 Reksten, A.M., Ho, Q.T., Nøstbakken, O.J., Markhus, M.W., Kjellevoll, M., Bøkevoll, A.,  
724 Hannisdal, R., Frøyland, L., Madsen, L., Dahl, L., 2022. Temporal variations in the nutrient  
725 content of Norwegian farmed Atlantic salmon (*Salmo salar*), 2005–2020. *Fd. Chem.* 373, 131445.

726 Richter, C.K., Skulas-Ray, A.C., Kris-Etherton, P.M., 2016. Recommended intake of fish and fish  
727 oils worldwide. In: *Fish and Fish Oil in Health and Disease Prevention* (Raatz, S.K., Bibus, D.M.,  
728 Eds.) pp. 27-48. Academic Press, Elsevier: New York.

729 Rosenlund, G., Torstensen, B.E., Stubhaug, I., Usman, N., Sissener, N.H., 2016. Atlantic salmon  
730 require long-chain n-3 fatty acids for optimal growth throughout the seawater period. *J. Nutr. Sci.*  
731 5, e19.

732 Ruiz-Lopez, N., Haslam, R.P., Napier, J.A., Sayanova, O., 2014. Successful high-level accumulation  
733 of fish oil omega-3 long-chain polyunsaturated fatty acids in a transgenic oilseed crop. *Plant J.* 77,  
734 198-208.

735 Ruyter, B., Bou, M.M., Turid, Sissener, N.H.S., Monica, Lutfi, E., Østbye, T.-K., 2022. A dose-  
736 response study with omega-3 rich canola oil as a novel source of docosahexaenoic acid (DHA) in  
737 feed for Atlantic salmon (*Salmo salar*) in seawater; effects on performance, tissue fatty acid  
738 composition, and fillet quality. *Aquaculture* 561, 738733.

739 Ruyter, B., Sissener, N.H., Østbye, T.-K., Simon, C.J., Krasnov, A., Bou, M., Berge, G.M., 2019. n-  
740 3 Canola oil effectively replaces fish oil as a new safe dietary source of DHA in feed for juvenile  
741 Atlantic salmon. *Br. J. Nutr.* 122, 1329–1345.

742 Salem, N., Jr., Eggersdorfer, M., 2015. Is the world supply of omega-3 fatty acids adequate for  
743 optimal human nutrition? *Curr. Opin. Clin. Nutr. Metab. Care* 18, 147–154.

744 Salunkhe, D.K., Adsule, R.N., Chavan, J.K., Kadam, S.S., 1992. *World Oilseeds: Chemistry,*  
745 *Technology and Utilization.* New York: VanNostrand Reinhold Company.

746 Santigosa, E., Brambilla, F., Milanese, L., 2021. Microalgae oil as an effective alternative source of  
747 EPA and DHA for gilthead seabream (*Sparus aurata*) aquaculture. *Animals* 11, 971.

748 Santigosa, E., Constant, D., Prudence, D., Wahli, T., Verlhac-Trichet, V., 2020. A novel marine algal  
749 oil containing both EPA and DHA is an effective source of omega-3 fatty acids for rainbow trout  
750 (*Oncorhynchus mykiss*). *J. World Aquacult. Soc.* 51, 649–665.

751 Santigosa, E., Olsen, R.E., Madaro, A., Trichet, V.V., Carr, I., 2023. Algal oil gives control of long-  
752 chain omega-3 levels in full-cycle production of Atlantic salmon, without detriment to  
753 zootechnical performance and sensory characteristics. *J. World Aquacult. Soc.* 54, 861-881.

754 Sissener, N.H., Waagbo, R., Rosenlund, G., Tvenning, L., Susort, S., Lea, T.B., Oaland, Ø., Chen,  
755 L., Breck, O., 2016. Reduced n-3 long chain fatty acid levels in feed for Atlantic salmon (*Salmo*  
756 *salar* L.) do not reduce growth, robustness or product quality through an entire full scale  
757 commercial production cycle in seawater. *Aquaculture* 464, 236–245.

758 Sprague, M., Betancor, M.B., Tocher, D.R., 2017. Microbial and genetically engineered oils as  
759 replacements for fish oil in aquaculture feeds. *Biotechnol. Lett.* 39, 1599-1609.

760 Sprague, M., Dick, J.R., Tocher, D.R., 2016. Impact of sustainable feeds on omega-3 long-chain fatty  
761 acid levels in farmed Atlantic salmon, 2006-2015. *Sci. Rep.* 6, 21892.

762 Stark, K.D., Van Elswyk, M.E., Higgins, M.R., Weatherford, C.A., Salem Jr, N., 2016. Global survey  
763 of the omega-3 fatty acids, docosahexaenoic acid and eicosapentaenoic acid in the blood stream  
764 of healthy adults. *Prog. Lipid Res.* 63, 132-152.

765 Tacon, A.G.J., 2020. Trends in global aquaculture and aquafeed production: 2000–2017. *Rev. Fish.*  
766 *Sci. Aquacult.* 28, 43–56.

767 Tacon, A.G.J., Metian, M., McNevin, A.A., 2022. Future feeds: suggested guidelines for sustainable  
768 development. *Rev. Fish. Sci. Aquacult.* 30, 135–142.

769 Tocher, D.R., 2010. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquaculture*  
770 *Res.* 41, 717-732.

771 Tocher, D.R., 2015. Omega-3 long-chain polyunsaturated fatty acids and aquaculture in perspective.  
772 *Aquaculture* 449, 94-107.

773 Tocher, D.R., Glencross, B.D., 2015. Lipids and fatty acids. In: *Dietary Nutrients, Additives, and Fish*  
774 *Health.* (Lee, C.-S., Lim, C., Webster, C., Gatlin III, D.M., Eds.), Ch.3. pp. 47-94, Wiley-  
775 Blackwell.

776 Tocher, D.R., Harvie, D.G., 1988. Fatty acid compositions of the major phosphoglycerides from fish  
777 neural tissues; (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout (*Salmo gairdneri*) and  
778 cod (*Gadus morhua*) brains and retinas. *Fish Physiol. Biochem.* 5, 229-239.

779 Tocher, D.R., Betancor, M.B., Sprague, M., Olsen, R.E., Napier, J.A., 2019. Omega-3 long-chain  
780 polyunsaturated fatty acids, EPA and DHA: Bridging the gap between supply and demand.  
781 *Nutrients* 11, 89.

782 Turchini, G.M., Du, Z.-Y., Olsen, R.E., Francis, D., Ringø, E., Tocher, D.R., 2022. The Lipids. In  
783 *Fish Nutrition* (Hardy, R.W., Kaushik, S.J., eds.) Fourth Edition, Ch.5, pp. 303-467, Academic  
784 Press, San Diego.

785 Turchini, G.M., Ng, W.K., Tocher, D.R. (Eds.), 2011. *Fish Oil Replacement and Alternative Lipid*  
786 *Sources in Aquaculture Feeds.* Taylor & Francis, CRC Press, Boca Raton. p.533.

787 Usher, S., Han, L., Haslam, R.P., Michaelson, L.V., Sturtevant, D., Aziz, M., Chapman, K.D.,  
788 Sayanova, O., Napier, J.A., 2017. Tailoring seed oil composition in the real world: optimising  
789 omega-3 long chain polyunsaturated fatty acid accumulation in transgenic *Camelina sativa*. Sci.  
790 Rep. 7, 6570.

791 Vera, L.M., Hamre, K., Espe, M., Hemre, G.-I., Skjaerven, K., Lock, E.-J., Prabu, A.J., Leeming, D.,  
792 Migaud, H., Tocher, D.R., Taylor, J.F., 2020. Higher dietary micronutrients are required to  
793 maintain optimal performance of Atlantic salmon (*Salmo salar*) fed a high plant material diet  
794 during the full production cycle. Aquaculture 528, 735551.

795 Ytrestøyl, T., Aas, T.S., Åsgård, T., 2015. Utilisation of feed resources in production of Atlantic  
796 salmon (*Salmo salar*). Aquaculture 448, 365-374.

797

798 Table 1. Formulations of experimental diets fed for the initial (250 – 800 g), intermediate (800 -  
799 1500 g) and final pellet sizes (1500 g – harvest).

Ingredient (g.kg <sup>-1</sup> )	Initial (5 mm)		Intermediate (7 mm)		Final (10 mm)	
	COM	TCO	COM	TCO	COM	TCO
Fishmeal	150	150	75	75	75	75
Soy protein concentrate	244	244	101	101	80	80
Maize gluten	50	50	50	50	0	0
Pea protein	74	74	54	54	121	121
Guar meal	0	0	150	150	150	150
Wheat	113	113	109	109	110	110
Land animal products	100	100	100	100	100	100
Fish oil	136	0	108	0	116	0
Rapeseed oil	102	0	208	0	205	0
Camelina oil (GM)	0	238	0	316	0	320
Premixes	32	32	36	36	35	35
Yttrium oxide	0.5	0.5	0.5	0.5	0.5	0.5

800 COM, control/reference feed reflecting current commercial practices; TCO, feed with all added oil  
801 supplied by the oil from transgenic Camelina.

Table 2. Fatty acid compositions (% total fatty acids, and mg fatty acid.100g<sup>-1</sup>) of experimental feeds.

Fatty acid	Initial				Final			
	Percentage		mg.100g <sup>-1</sup>		Percentage		mg.100g <sup>-1</sup>	
	COM	TCO	COM	TCO	COM	TCO	COM	TCO
14:0	2.37	0.61	520.10	111.63	3.16	0.23	780.83	61.17
16:0	9.84	7.78	2160.92	1433.20	9.84	6.74	2434.57	1769.62
18:0	2.27	4.49	498.79	827.58	3.24	4.38	801.67	1150.21
Total saturated <sup>1</sup>	15.34	15.63	3367.64	2878.28	18.13	14.73	4484.79	3865.06
16:1n-7	4.14	1.00	908.05	184.82	4.43	0.40	1094.80	105.62
18:1n-9	32.87	9.65	7217.02	1776.93	39.29	10.01	9717.74	2625.34
18:1n-7	3.72	1.65	816.77	304.43	3.09	1.39	763.09	365.80
20:1n-9	6.27	6.46	1377.50	1189.01	1.30	7.87	322.41	2064.11
22:1n-11	4.76	1.08	1044.97	198.42	0.11	0.00	27.88	0.00
Total monoenes <sup>2</sup>	54.05	21.75	11866.86	4006.63	49.14	22.23	12153.44	5833.39
18:2n-6	11.60	20.40	2546.36	3757.90	15.20	19.62	3759.71	5147.45
18:3n-6	0.08	1.86	16.55	342.43	0.10	1.48	25.62	387.88
20:2n-6	0.17	1.65	37.37	303.35	0.10	1.58	25.34	414.43
20:3n-6	0.07	0.83	15.80	153.08	0.08	0.47	19.71	124.12
20:4n-6	0.31	2.59	68.21	477.38	0.40	1.68	98.27	439.79
22:4n-6	0.00	0.67	0.00	123.50	0.00	0.37	0.00	96.37
22:5n-6	0.08	0.09	16.55	16.19	0.09	0.05	22.25	12.83

Total n-6 PUFA	12.30	28.09	2700.83	5173.83	15.97	25.24	3950.90	6622.88
18:3n-3	4.92	9.96	1079.33	1833.93	5.81	19.10	1436.36	5011.39
18:4n-3	1.13	1.34	249.02	247.22	0.68	1.21	167.26	317.76
20:3n-3	0.07	0.97	15.30	177.91	0.00	1.15	0.00	301.35
20:4n-3	0.36	1.99	77.99	366.61	0.37	1.48	91.23	388.18
20:5n-3	4.81	8.47	1056.26	1560.59	4.85	5.70	1198.98	1494.53
22:5n-3	0.64	4.40	141.19	810.52	0.74	3.85	182.75	1009.08
22:6n-3	5.55	7.40	1217.51	1362.82	2.43	5.27	600.06	1382.04
Total n-3 PUFA	17.48	34.53	3892.01	6359.60	15.08	37.75	3730.70	9904.33
Total n-3 LC-PUFA	11.35	22.26	2492.94	4100.55	8.39	16.30	2073.02	4273.83
n-3 PUFA/n-6 PUFA	1.42	1.23	1.42	1.23	0.94	1.50	0.94	1.50

Values are means of duplicate assays. <sup>1</sup>, includes 15:0, 20:0, 22:0 and 24:0, present at up to 0.3 %. <sup>2</sup>, includes 16:1n-9, 17:1, 20:1n-11, 20:1n-7, 22:1n-9 and 24:1n-9, present at up to 0.8 %. COM, control/reference feed reflecting current commercial practices; LC-PUFA, long-chain PUFA; PUFA, polyunsaturated fatty acids; TCO, feed with all added oil supplied by the oil from transgenic Camelina.



Table 3. Effects of diet on survival, growth performance, biometric parameters, feed intake and feed efficiency.

Parameter	COM	TCO	
<b><u>OVERALL</u></b>			
Initial Weight (g)	186.6 ± 2.5	187.7 ± 1.3	
Final Weight (g)	3601.9 ± 307.6	3108.3 ± 87.0	
Final Length (cm)	63.1 ± 1.8	60.6 ± 0.8	
Weight gain (g)	3414.2 ± 307.2	2921.7 ± 85.6	
SGR	1.1 ± 0.0	1.1 ± 0.0	
TGC	3.3 ± 0.1	3.1 ± 0.0	
HSI	1.1 ± 0.0	1.2 ± 0.0	*
VSI	10.1 ± 0.3	10.0 ± 0.6	
FI (g/fish/day)	16.1 ± 0.9	14.8 ± 1.6	
FCR	1.3 ± 0.1	1.3 ± 0.1	
Survival (%)	95.8 ± 2.0	94.6 ± 1.4	
<b><u>WEEKS 0 - 11</u></b>			
Initial Weight (g)	186.7 ± 13.7	188.0 ± 13.7	
Final Weight (g)	855.0 ± 24.1	813.0 ± 11.3	
Length (cm)	40.8 ± 1.1	39.6 ± 0.4	
Weight gain (g)	668.0 ± 24.1	626.0 ± 11.3	
SGR	2.0 ± 0.0	1.9 ± 0.0	
TGC	3.2 ± 0.1	3.1 ± 0.0	
FI (g/fish/day)	8.4 ± 1.0	7.9 ± 0.5	
FCR	1.0 ± 0.1	1.0 ± 0.0	
Condition (k)	1.0 ± 0.0	1.0 ± 0.0	
<b><u>WEEKS 12 - 23</u></b>			
Initial Weight (g)	855.0 ± 24.1	813.0 ± 11.3	
Final Weight (g)	2071.2 ± 97.3	1812.4 ± 37.0	*
Length (cm)	51.6 ± 0.7	49.4 ± 0.3	*
Weight gain (g)	1216.2 ± 75.7	999.4 ± 36.3	*
SGR	1.2 ± 0.1	1.0 ± 0.0	*
TGC	3.3 ± 0.1	2.9 ± 0.1	*
FI (g/fish/day)	17.6 ± 0.9	15.4 ± 1.8	
FCR	1.1 ± 0.0	1.2 ± 0.1	
Condition (k)	0.9 ± 0.0	0.8 ± 0.0	
<b><u>WEEKS 24 - 37</u></b>			
Initial Weight (g)	2071.2 ± 97.3	1812.4 ± 37.0	*
Final Weight (g)	3601.9 ± 307.6	3108.3 ± 87.0	

Length (cm)	63.1 ± 1.8	60.6 ± 0.8	
Weight gain (g)	1530.7 ± 97.3	1295.9 ± 37.0	
SGR	0.6 ± 0.1	0.6 ± 0.0	
TGC	3.6 ± 0.4	3.4 ± 0.1	
FI (g/fish/day)	19.3 ± 1.4	18.3 ± 3.2	
FCR	1.2 ± 0.1	1.4 ± 0.3	
Condition (k)	1.4 ± 0.0	1.3 ± 0.0	*

Values are means ± SD (n = 3). An asterisk denotes a significant difference between mean values for fish fed the COM and TCO diets.

COM, control/reference feed reflecting current commercial practices;

FCR, feed conversion ratio; FI, feed intake; SGR, specific growth rate;

TCO, feed with all added oil supplied by the oil from transgenic Camelina;

TGC, thermal growth coefficient.

Table 4. Effects of diet on proximate compositions (percentage) and fatty acid compositions (percentage and mg.100 g<sup>-1</sup>) of total lipid of whole fish.

	Percentage		mg. 100g <sup>-1</sup>	
	COM	TCO	COM	TCO
<u>Proximate composition</u>				
Lipid	21.11 ± 1.75	20.08 ± 1.20	-	-
Protein	15.65 ± 0.55	15.79 ± 0.30	-	-
Ash	1.68 ± 0.18	1.65 ± 0.09	-	-
Moisture	59.42 ± 1.28	59.71 ± 0.70	-	-
<u>Fatty acid</u>				
14:0	1.94 ± 0.05	0.42 ± 0.06 *	364.4 ± 24.4	72.9 ± 10.0 *
16:0	9.28 ± 0.08	7.58 ± 0.21 *	1747.5 ± 119.9	1310.4 ± 90.2 *
18:0	2.64 ± 0.08	4.07 ± 0.10 *	496.4 ± 43.7	703.1 ± 54.0 *
Total saturated <sup>1</sup>	14.47 ± 0.10	13.64 ± 0.23 *	2722.9 ± 193.4	2357.5 ± 153.7 *
16:1n-7	3.23 ± 0.12	0.67 ± 0.09 *	606.8 ± 36.9	115.7 ± 17.8 *
18:1n-9	39.12 ± 0.69	12.57 ± 0.79 *	7360.2 ± 466.6	2179.0 ± 176.9 *
18:1n-7	4.23 ± 0.16	1.66 ± 0.15 *	795.7 ± 43.6	286.4 ± 29.1 *
20:1n-9	4.18 ± 0.08	6.61 ± 0.09 *	786.1 ± 59.9	1142.5 ± 84.1 *
22:1n-11	1.69 ± 0.15	0.32 ± 0.17 *	316.7 ± 29.3	53.7 ± 28.0 *
Total monoenes <sup>2</sup>	54.27 ± 0.98	23.65 ± 1.11 *	10208.8 ± 633.4	4086.0 ± 291.2 *
18:2n-6	13.99 ± 0.25	19.72 ± 0.04 *	2633.5 ± 182.4	3410.5 ± 246.0 *
18:3n-6	0.13 ± 0.03	1.06 ± 0.02 *	25.3 ± 6.5	183.4 ± 14.1 *
20:2n-6	0.94 ± 0.04	2.04 ± 0.28 *	177.8 ± 15.8	354.4 ± 61.2 *
20:3n-6	0.25 ± 0.03	1.17 ± 0.04 *	47.8 ± 7.7	201.8 ± 19.0 *
20:4n-6	0.30 ± 0.05	1.88 ± 0.06 *	57.0 ± 12.3	324.9 ± 26.1 *
22:4n-6	0.00 ± 0.00	0.64 ± 0.02 *	0.0 ± 0.0	110.7 ± 9.8 *

22:5n-6	0.00 ± 0.00	0.08 ± 0.02 *	0.0 ± 0.0	14.4 ± 4.2 *
Total n-6 PUFA	15.63 ± 0.33	26.60 ± 0.37 *	2941.3 ± 217.0	4600.2 ± 357.4 *
18:3n-3	5.53 ± 0.40	14.26 ± 0.16 *	1042.3 ± 115.8	2466.0 ± 181.5 *
18:4n-3	0.60 ± 0.03	1.03 ± 0.01 *	113.3 ± 8.8	178.2 ± 12.6 *
20:3n-3	0.41 ± 0.04	1.59 ± 0.06 *	78.0 ± 11.1	275.1 ± 24.7 *
20:4n-3	0.74 ± 0.05	2.17 ± 0.07 *	140.2 ± 13.7	375.4 ± 35.2 *
20:5n-3	2.69 ± 0.12	5.56 ± 0.13 *	507.3 ± 51.5	961.6 ± 80.7 *
22:5n-3	1.13 ± 0.08	4.58 ± 0.29 *	213.5 ± 26.2	792.0 ± 83.1 *
22:6n-3	3.86 ± 0.18	6.85 ± 0.38 *	726.8 ± 70.8	1185.3 ± 121.9 *
Total n-3 PUFA	15.12 ± 0.65	36.08 ± 0.87 *	2850.1 ± 274.3	6241.3 ± 522.3 *
Total PUFA	31.27 ± 0.91	62.71 ± 1.24 *	5888.9 ± 497.9	10847.9 ± 880.6 *
Total n-3 LC-PUFA	8.84 ± 0.33	20.74 ± 0.90 *	1665.7 ± 161.5	3589.4 ± 340.4 *
EPA:DHA	0.70 ± 0.05	0.81 ± 0.03 *	-	-
n-3PUFA:n-6PUFA	0.97 ± 0.03	1.36 ± 0.01 *	-	-
<u>Content (g.100g<sup>-1</sup>)</u>				
EPA	-	-	0.51 ± 0.05	0.96 ± 0.08 *
DHA	-	-	0.73 ± 0.07	1.19 ± 0.12 *
EPA+DHA	-	-	1.23 ± 0.12	2.15 ± 0.20 *
n-3LC-PUFA	-	-	1.66 ± 0.14	3.59 ± 0.28 *

Values are means ± SD (n = 6). An asterisk denotes a significant difference between mean values for fish fed the COM and TCO diets. <sup>1</sup>, includes 15:0, 20:0, 22:0 and 24:0, present at up to 0.3 %. <sup>2</sup>, includes 16:1n-9, 17:1, 20:1n-11, 20:1n-7, 22:1n-9 and 24:1n-9, present at up to 0.8 %. COM, control/reference feed reflecting current commercial practices; DHA, docosahexaenoic acid (22:6n-3); EPA, eicosapentaenoic acid (20:5n-3); LC-PUFA, long-chain PUFA; PUFA, polyunsaturated fatty acids; TCO, feed with all added oil supplied by the oil from transgenic Camelina.

Table 5. Effects of diet on proximate compositions (percentage) and fatty acid compositions (percentage and mg.100 g<sup>-1</sup>) of total lipid of muscle/flesh (NQC)

	Percentage				mg. 100g-1			
	COM		TCO		COM		TCO	
<u>Proximate composition</u>								
Lipid	16.43	± 1.50	14.50	± 0.90	*	-		-
Protein	18.85	± 0.75	19.64	± 0.44		-		-
Ash	1.64	± 0.15	1.75	± 0.13		-		-
Moisture	62.88	± 1.07	64.42	± 0.42	*	-		-
<u>Fatty acid</u>								
14:0	1.89	± 0.09	0.40	± 0.06	*	282.0	± 31.0	52.0 ± 7.9 *
16:0	9.08	± 0.12	7.46	± 0.16	*	1353.8	± 107.1	962.9 ± 49.2 *
18:0	2.62	± 0.11	3.85	± 0.11	*	391.7	± 39.0	497.0 ± 29.9 *
Total saturated <sup>1</sup>	14.27	± 0.16	13.23	± 0.25	*	2128.5	± 182.2	1707.5 ± 90.7 *
16:1n-7	3.15	± 0.10	0.63	± 0.12	*	470.7	± 45.6	81.5 ± 14.8 *
18:1n-9	39.19	± 0.99	11.97	± 1.09	*	5842.9	± 489.4	1544.4 ± 154.6 *
18:1n-7	3.49	± 0.07	1.60	± 0.09	*	520.9	± 43.1	207.1 ± 15.7 *
20:1n-9	4.00	± 0.18	6.66	± 0.09	*	597.6	± 62.9	860.0 ± 63.7 *
22:1n-11	1.54	± 0.08	0.30	± 0.03	*	228.5	± 17.4	38.3 ± 5.9 *
Total monoenes <sup>2</sup>	53.06	± 0.95	23.27	± 1.22	*	7911.8	± 662.5	3004.4 ± 232.4 *
18:2n-6	14.02	± 0.20	19.87	± 0.21	*	2092.2	± 201.0	2566.7 ± 188.8 *
18:3n-6	0.14	± 0.03	1.03	± 0.04	*	21.6	± 5.9	133.7 ± 10.3 *
20:2n-6	0.99	± 0.07	2.17	± 0.08	*	148.3	± 18.3	280.5 ± 23.7 *
20:3n-6	0.27	± 0.04	1.16	± 0.06	*	40.6	± 8.2	150.1 ± 15.3 *
20:4n-6	0.33	± 0.04	1.80	± 0.08	*	49.2	± 8.7	232.4 ± 18.7 *
22:4n-6	0.06	± 0.02	0.63	± 0.05	*	9.5	± 2.9	81.8 ± 9.8 *

22:5n-6	0.06 ± 0.01	0.07 ± 0.01		9.2 ± 1.9	8.5 ± 1.0	
Total n-6 PUFA	15.88 ± 0.37	26.73 ± 0.49 *		2370.7 ± 237.6	3453.6 ± 262.6 *	
18:3n-3	5.75 ± 0.41	14.74 ± 0.24 *		859.1 ± 116.2	1904.7 ± 143.9 *	
18:4n-3	0.61 ± 0.04	0.99 ± 0.02 *		91.6 ± 11.6	127.9 ± 8.4 *	
20:3n-3	0.44 ± 0.06	1.62 ± 0.07 *		65.5 ± 11.5	209.2 ± 18.7 *	
20:4n-3	0.78 ± 0.06	2.17 ± 0.08 *		116.9 ± 15.4	280.7 ± 26.4 *	
20:5n-3	2.85 ± 0.10	5.40 ± 0.11 *		424.6 ± 39.9	697.3 ± 44.5 *	
22:5n-3	1.29 ± 0.11	4.68 ± 0.21 *		193.4 ± 30.4	605.3 ± 58.3 *	
22:6n-3	4.36 ± 0.14	7.07 ± 0.31 *		650.5 ± 54.3	914.0 ± 85.1 *	
Total n-3 PUFA	16.25 ± 0.68	36.73 ± 0.80 *		2426.9 ± 266.6	4745.8 ± 376.1 *	
Total PUFA	32.67 ± 1.02	63.50 ± 1.23 *		4878.2 ± 510.9	8204.5 ± 634.8 *	
Total n-3 LC-PUFA	9.72 ± 0.29	20.94 ± 0.69 *		1450.9 ± 142.2	2706.5 ± 229.5 *	
EPA:DHA	0.65 ± 0.02	0.76 ± 0.03 *		-	-	
n-3PUFA:n-6PUFA	1.02 ± 0.02	1.37 ± 0.01 *		-	-	
<u>Content (g.100g<sup>-1</sup>)</u>						
EPA	-	-		0.42 ± 0.04	0.70 ± 0.04 *	
DHA	-	-		0.65 ± 0.05	0.91 ± 0.09 *	
EPA+DHA	-	-		1.08 ± 0.09	1.61 ± 0.13 *	
n-3LC-PUFA	-	-		1.45 ± 0.12	2.71 ± 0.19 *	

Values are means ± SD (n = 6). An asterisk denotes a significant difference between mean values for fish fed the COM and TCO diets. <sup>1</sup>, includes 15:0, 20:0, 22:0 and 24:0, present at up to 0.3 %. <sup>2</sup>, includes 16:1n-9, 17:1, 20:1n-11, 20:1n-7, 22:1n-9 and 24:1n-9, present at up to 0.8 %. COM, control/reference feed reflecting current commercial practices; DHA, docosahexaenoic acid (22:6n-3); EPA, eicosapentaenoic acid (20:5n-3); LC-PUFA, long-chain PUFA; PUFA, polyunsaturated fatty acids; TCO, feed with all added oil supplied by the oil from transgenic Camelina.

Table 6. Summary of flesh quality measurements for each dietary treatment

	COM	TCO
Firmness (mJ)	525.3 ± 26.3	544.2 ± 34.3
Tensile strength (mJ)	267.7 ± 14.1	282.2 ± 15.7
Gaping	nd	nd
Roche colour score A	26.8 ± 0.4	27.1 ± 0.6
Roche colour score B	26.7 ± 0.5	27.9 ± 0.6
Roche colour score C	28.7 ± 0.3	28.9 ± 0.3
Average colour ABC	27.2 ± 0.4	28.0 ± 0.4*
Total carotenoids (mg.kg <sup>-1</sup> )	2.65 ± 0.63	2.20 ± 0.42

Values are means ± SEM (n = 12) except for carotenoid content of NQC, which was n = 3. An asterisk denotes a significant difference between mean values for fish fed the COM and TCO diets. COM, control/reference feed reflecting current commercial practices; nd, no noteworthy gaping detected; TCO, feed with all added oil supplied by the oil from transgenic Camelina.