

1 Comparative evaluation of Polymerase Chain Reaction.-Restriction Enzyme Analysis
2 (PRA) and Sequencing of Heat shock protein 65 (*hsp65*) gene for Identification of Aquatic
3 Mycobacteria

4
5 **F Pourahmad^{1,2}*, K D Thompson¹, A Adams¹ and R H Richards¹**

6
7 ¹ Institute of Aquaculture, University of Stirling, Stirling, Scotland, UK

8 ² School of Veterinary Medicine, Ilam University, P.O. Box: 69315-516, Ilam, Iran

9
10
11
12
13
14
15
16
17
18 Running title: PRA and sequencing of *hsp65* gene for identification of aquatic
19 mycobacteria

20
21 Corresponding author: Fazel Pourahmad, Institute of Aquaculture, University of Stirling,
22 Stirling, FK9 4LA, Scotland / UK, Tel: +44 (0)1786 467878, Fax: +44 (0) 1786 472133,
23 Email: fp4@stir.ac.uk

1 Abstract

2 Traditional identification of mycobacteria based on cultural and biochemical tests can take
3 several weeks and may fail to provide a precise identification. Polymerase Chain Reaction-
4 restriction analysis (PRA) of the gene encoding heat shock protein 65 kDa (*hsp65*) gene
5 has been proposed as a rapid and inexpensive alternative approach. Despite being widely
6 used for differentiation of mammalian mycobacteria, this method has only been applied in
7 the identification of a small number of aquatic mycobacteria. The present study aimed to
8 evaluate the potential use of PRA of *hsp65* for the identification of aquatic mycobacteria
9 compared with sequence analysis. Seventy one mycobacterial isolates including, 10 type/
10 reference strains and the remainder field isolates, were subjected to PRA of a 441 bp
11 fragment of this gene. For 68 representative isolates, sequence analysis was performed. All
12 rapidly and slowly growing mycobacteria had best matches with 99.3% to 100% similarity
13 with their corresponding species in the databanks.

14 PRA proved to be a simple and rapid method for identifying aquatic mycobacteria.

15 However, the incidence of similar or identical restriction patterns for some species of
16 mycobacteria, and in particular, identification of new species of mycobacteria is a major
17 problem using such a method. In contrast, the nucleic acid sequencing of the *hsp65* gene
18 yielded unambiguous results.

1. Introduction

Mycobacteriosis (fish tuberculosis) is a progressive disease of a wide range of wild and captive marine and freshwater fish species (Chinabut, 1999). The zoonotic nature of the organism, the massive economical losses which have occurred to the aquaculture industry due to this disease and the lack of effective treatment highlight the need for mycobacteria to be rapidly detected and identified to species level.

Traditional identification of mycobacteria based on cultural and biochemical tests may take several weeks, and sometimes even then fail to provide a precise identification. In addition, the procedures of these tests are time consuming and complicated. Therefore, various molecular based methods have been developed for the rapid detection and identification of mycobacteria species. There are currently several commercially available methods, including the AccuProbe system (Genprobe, San Diego, Calif), INNO-LiPA (Innogenetics, Ghent, Belgium), or GenoType *Mycobacterium* assay (Hain, Lifescience, Germany). Although such tests provide desirable sensitivity and specificity, they are limited to the identification of a small number of mycobacteria.

Among other molecular methods, Polymerase Chain Reaction (PCR) -Restriction Enzyme Analysis (PRA), and sequence analysis have been applied to several mycobacteria genes. In this regard, PCR of the 441 bp of heat shock protein 65 kD gene (*hsp65*), followed by restriction fragment pattern analysis by *BstEII* and *HaeIII* enzymes proposed by Telenti et al. (1993), has provided a rapid method for identifying mycobacteria and closely related species (Brunello et al., 2001; Devallois et al., 1997). In addition, an internet database (PRASite) (<http://app.chuv.ch/prasite/index.html>) has been established that contains an algorithm

1 of *hsp65* PRA patterns to assist with identification. Three hundreds and forty eight
2 patterns of 113 species have so far been included in the PRAsite database.

3 Due to the hypervariability of the *hsp65* gene, sequence analysis of Telenti fragment
4 (441 bp) has become a routine method in taxonomical studies (Murcia et al., 2006;
5 Wilson et al., 2001) and in identification of clinical isolates of mycobacteria (Li et al.,
6 2003; McNabb et al., 2004; Ringuet et al., 1999; Turenne et al., 2006). The PRA
7 technique is relatively inexpensive and does not need specialised equipment; whereas,
8 sequencing is more discriminative but requires highly sophisticated equipment.

9 Unlike the extensive use of these methods in clinical medicine, there are limited
10 reports of the application of these techniques for detection and identification of fish
11 mycobacteria.

12 The aims of this study were (i) to evaluate the possible use of PRA and sequence
13 analysis based on the 441 bp fragment of *hsp65* gene and (ii) to compare the
14 reliability of these methods in identification and differentiation of aquatic
15 mycobacteria.

16 **2. Materials and Methods**

17 **2.1 Bacterial strains**

18 The 10 type/reference strains and 61 field isolates used in this study are listed in Table
19 1. Identification at the species level had previously been carried out by a variety of
20 molecular methods (Pate et al., 2005; Pourahmad, 2008; Puttinaowarat et al., 2002).
21 All strains were grown on Middlebrook 7H10 medium supplemented with oleic acid-
22 albumin-dextrose-catalase (OADC) and 0.5% glycerol.

2.2 DNA preparation

DNA was extracted from cultures using NucleoSpin[®] columns (Macherey–Nagel, Germany) following manufacturer's instructions.

2.3 PCR amplification of *hsp65*

PCR mixtures were prepared with 12.5 µl of a PCR master mix (AB gene[®], UK) containing 10 mM Tris-HCl (pH 8.3), 1.5 mM MgCl₂, 10% glycerol, 200 µM (each) deoxynucleoside triphosphates, 10 pmol of each primer Tb11 (5'-ACCAACGATGGTGTGTCCAT-3') and Tb12 (5'-CTTGTCGAACCGCATAACCCT-3'), as described by Telenti et al., (1993), 1.25 U of *Taq* polymerase, 2 µl of each template containing 10-50 ng DNA and ultra-pure water (Millipore) to final volume of 25 µl. PCR reactions of a 441 bp fragment of *hsp65* gene were performed in an automated T-gradient thermocycler (Biometra[®], Göttingen, Germany) with the following amplification profile: initial step of 5 min incubation at 94°C, followed by 35 cycles of 1 min at 94°C, 1 min at 60°C, 1 min at 72°C, and a final extension of 5 min at 72°C. The PCR product specificity was checked by electrophoresis on a 1.2% agarose gel for 45 min at 120 V.

2.4 Restriction enzyme analysis of the *hsp65* PCR product

Two separate restriction digests (*Bst*EII and *Hae*III, both purchased from Sigma) were enzyme, 1.5 µl of appropriate restriction buffer, and 15 µl of ultra-pure water. The mixtures were incubated overnight at 60°C for *Bst*EII digestion and for 60 min at 37°C for *Hae*III digestion. Restriction fragments were electrophoresed on a 2.5% Low Melting Point agarose gel (Promega, UK) at 70 V for 2.5 h and visualized by ethidium bromide staining. A 50 bp ladder (Sigma) served as an external molecular size marker, and was added at the both ends of each gel to reduce migration-related

errors. The restriction pattern was compared to the published algorithm (Telenti et al., 1993) and with available PRA profiles from PRAsite.

2.5 Sequencing of the *hsp65* PCR product

The remaining PCR amplicons from the PRA were purified using the Qiaquick purification kit as described by the supplier (Qiagen, UK) and quantified by Nanodrop spectrophotometry, before being sequenced in both directions, using the PCR primers (Tb11 and Tb12) as well as an in house design forward primer, F927 (5'-GGACCCGTACGAGAAGAT-3'). Sequencing was carried out using GenomeLab™ Dye Terminator Cycle Sequencing Quick Start Kit (as per manufacturer's instructions) and detected on an automated capillary sequencer, CEQ™ 8800 (Beckman Coulter, UK).

2.6 Sequence and phylogenetic analysis

Both forward and reverse sequences were assembled and edited using the SeqMan II module of Lasergene, Version 6 (DNASTAR, Inc., Madison, USA). The resulting consensus sequences were compared with published nucleotide sequences by BLASTn analysis. Phylogenetic trees were constructed, using the neighbour-joining method in MEGA version 3.1 software (Kumar et al., 2004) applying the Kimura two-parameter distance correction model. The tree was rooted using *Mycobacterium* sp. isolate DL049 (accession no AM902972) as the outgroup. Statistical confidence in tree branches were generated by performing 1000 bootstrap replicates

2.7 Nucleotide sequence accession numbers

The nucleotide sequences described herein have been deposited in the GenBank database under accession no. AM398442- AM398480; AM902952- AM902978 and EF535603-EF535604

1 **3. Results**

2 **3.1 Sequence analysis of *hsp65***

3 Primer Tb11 gave inconsistent sequencing results and was later replaced by primer
4 F927 (see Section 2.5). Use of this primer, which is 39 bp internal to primer Tb11,
5 resulted in greatly improved sequence data. Sixty-eight of the 71 isolates mentioned
6 above were sequenced. With the exception of isolates of *Mycobacterium* sp. isolate
7 126/5/03, *M. szulgai* /*M. intremedium* isolate 54/02, *M. gordonae* isolate 277/3/01 and
8 *Mycobacterium* sp. isolate DL049, all rapidly and slowly growing mycobacteria had
9 best matches with 99.3% to 100% similarity with their corresponding species in the
10 databanks. The phylogenetic relationships among these isolates is presented in Fig. 3.

11 The resulting tree demonstrated two major clusters where slowly growing and rapidly
12 growing mycobacteria could be distinguished. Sequence data resolved some
13 ambiguous results with the previous methods, i.e. conventional PCR and PRA. All
14 species screened were readily discriminated from each other. *M. fortuitum*, *M.*
15 *senegalense* and *M. conceptionense*, the three species with the highest degree of
16 similarity for the 16S rRNA gene (Kirschner et al., 1993), had sequences differing by
17 7 and 9 bp, respectively (Figs.4). Similarly, the isolates of *M. peregrinum* and *M.*
18 *fortuitum* could be differentiated by 12 nucleotides. Furthermore, using a programme
19 such as RestrictionMapper (version 3) <http://www.restrictionmapper.org>, enabled
20 precise definition of the sizes of the PRA restriction fragments from sequences (Table
21 3).

22 Despite repeating the DNA extraction and purification from *M. chelonae* isolates, 2 of
23 5 *M. chelonae* isolates, *M. chelonae* (NCIMB 13533) and *M. chelonae* MT1900,
24 could only be amplified by primers Tb11 and Tb12 when higher cycles (38 times

1 compare to 35 cycles) of PCR were applied. More interestingly, when their sequences
2 were compared against databanks, they were more closely related (99.3% similarity)
3 to *M. salmoniphilum*, a new *Mycobacterium chelonae*-like organism (Whipps et al.,
4 2007). However, using overlapping primers to Tb11 and Tb12, F927 and R927, those
5 two isolates were easily amplified and sequenced.

6 **3.2 PRA of *hsp65***

7 The results obtained from the PRA analysis of 71 mycobacterial isolates are
8 summarized in Table 2. For easy interpretation of the PRA profiles generated by each
9 field isolate, a 50 bp ladder size marker was used as a reference and the PRA profile
10 of a type strain was used as an internal control.

11 *Hae*III and *Bst*EII digestion of the 441 bp *hsp65* PCR amplicon generated fragments
12 ranging from 50 to 441 bp. Restriction fragments smaller than 50 bp were omitted in
13 order to reduce confusion with primer-dimer bands, and restriction fragment band
14 sizes were rounded to the nearest 5 bp, as recommended by Telenti et al. (1993).

15 For some species, such as *M. gordonae*, that are known to contain several subtypes,
16 their subtypes generated distinctive restriction profiles (Fig. 1, lanes 3, 6, 18).

17 Therefore, the results clearly indicated that the PRA method could differentiate
18 mycobacteria at the species and even the subspecies level.

19 All *M. marinum*, *M. shottsii*, *M. pseudoshottsii*, *M. peregrinum*, *Mycobacterium* sp.
20 DL049 and *M. gordonae* type V (Fig. 1 lane 3) displayed similar patterns on *Bst*EII
21 digestion (240 and 210 bp); however, with the exception of the first 3 species, the
22 remaining species could be discriminated from the *Hae*III digestion

1 Although 6 strains of *M. fortuitum* including two type strains (NCIMB 1294 and
2 1295) and 4 field isolates had an identical PRA pattern, they were indistinguishable
3 from *M. conceptionense* and *M. senegalense* strains due to the same digestion pattern
4 for *Bst*EII and very similar *Hae*III pattern (Fig. 2 lanes 1-8 and 12)

5 There were 6 isolates with patterns not represented in the literature or in the PRA
6 website (Fig. 1 lanes 5, 12 and 14). The type strain and field isolates of *M.*
7 *peregrinum* had the pattern of *M. peregrinum* II as described by Devallois et al.
8 (1997) and was present on the PRA website (240 and 210 bp upon digestion with
9 *Bst*EII and the pattern of 140/120/100/60 for *Hae*III digestion) as illustrated in Fig 3
10 lane 11.

11 The advantage of sequence analysis over PRA was most obvious for the *M. fortuitum*
12 group, presumed *M. gordonae* 277/3/01 and species with new PRA patterns which
13 could only be specified via sequencing (Table 3). Of the *M. fortuitum* group, 15
14 isolates were matched with *M. conceptionense* (99.7% - 100% similarity); isolates
15 S11 and S12 resembled *M. mucogenicum* (99.3% matched) and *M. gordonae* 277/3/01
16 was most closely related to *M. arupene* (99.5%). In addition, *M. stomatepieae* isolates
17 T3, T4 and T11^T, '*M. angelicum*' 126/5/03^T, '*M. aemonae*' 54/02^T were closely
18 related to *M. triplex*/*M. lentiflavum* (98% similar), *M. avium* (96.8% matched) and *M.*
19 *gordonae* (95.5% similarity), respectively.

20

21 **4. Discussion**

22 The molecular identification of mycobacteria is of growing interest for routine
23 speciation of these organisms. Several genetic based methods have been carried out

1 including sequence analysis of the 16S rRNA (Turenne et al., 2001), the spacer
2 between 16S-23S rRNA (Roth et al., 1998), the gene coding for the beta subunit of
3 RNA polymerase (*rpoB*) gene (Kim et al., 1999) and partial sequencing (Ringuelet et
4 al., 1999; McNabb et al., 2004) and PRA of *hsp65* gene (Brunello et al., 2001;
5 Devallois et al., 1997). Of these, a relatively simple method providing promising
6 results is PRA of *hsp65* gene. Despite being widely used for differentiation of
7 mammalian mycobacteria, this method has only been used in the identification of a
8 small number of aquatic mycobacteria (Rhodes et al., 2005; Ucko et al., 2002).

9 Seventy-one mycobacterial isolates were subjected to PRA of the *hsp65* gene. All
10 isolates of the well-characterised species, *M. marinum*, *M. fortuitum* and *M.*
11 *gordonae*, could be correctly identified using the PRA method. However, because of
12 the number of fragment patterns not yet reported, this technique could not be used
13 solely as a tool for identification of rare or novel species of mycobacteria. Both
14 isolates of *M. mucogenicum* included in the study had patterns most similar to *M.*
15 *chitae* type I when compared to patterns in the PRA web site.

16 The use of PRA for subtyping mycobacteria may improve our epidemiological
17 understanding of these organisms in the aquatic environment. In this study, *M.*
18 *gordonae* type I, one of the most frequently reported types of this species in clinical
19 laboratories constituted 1/3 of the fish isolates of *M. gordonae*, probably indicating the
20 importance of this type among aquatic isolates of this species. Using PRA of *hsp65*,
21 other researchers showed that *M. kansasii* type I is the most common human isolate of
22 this species (Alcaide et al., 1997; Cheunoy et al., 2005).

23 A comparison was performed between restriction fragment size from running agarose
24 gels and the real size of fragments deduced from sequence analysis (Table 3). The

1 agarose gel-based PRA fragments normally differed in size between the two methods
2 by up to 9 bp, apart from patterns for *HaeIII* digestion of *M. gordonae* IV which were
3 very different (200, 60 bp for the estimated compared to 161, 110, 60 bp obtained
4 from gel) (Table 3). Brunello et al. (2001) showed that PRA patterns with a more
5 accurate size were obtained using a 10% polyacrylamide gel. However, in practice
6 polyacrylamide gels are more difficult to use than agarose gels as discussed elsewhere
7 (Devallois et al., 1997).

8 The sequence variation present within the mycobacterial *hsp65* gene has made it an
9 attractive locus for species identification and molecular phylogeny (Smole et al.,
10 2002). In spite of the difficulty in amplifying the two isolates of *M. chelonae*, their
11 patterns were identical and very similar with *BstEII* and *HaeIII* restriction,
12 respectively. Nevertheless, intraspecies polymorphism in the 441 bp target region
13 observed in individual *M. chelonae* complex species, such as *M. chelonae*, was shown
14 to cause different restriction patterns for individual isolates of the same species,
15 thereby increasing the ambiguity of the method (Wilson et al., 2001). As reported by
16 other workers, differences between calculated sizes of restriction fragments and
17 published patterns because of running conditions, type of agarose used and the
18 computer programme used, as well as confusion in interpretation of patterns due to
19 lack of standardisation are major drawbacks of PRA (Devallois et al., 1997; Hafner et
20 al., 2004; Rocha et al., 1999). Another difficulty with PRA is that a single base
21 change may lead to the appearance or disappearance of a restriction site (Ringuet et
22 al., 1999). For instance, a point mutation in one of the restriction sites had shifted
23 typical PRA pattern of *M. malmoense* to the *M. marinum* specific PRA pattern (Nolte
24 et al., 2005). The existence of the PRA database web site for PRA of *hsp65* provides
25 an easy way of interpreting the patterns (Cheunoy et al., 2005), however, there was no

1 identical match when the actual size of fragments in Table 3 were compared against
2 those in the database. This observation suggests that the PRA web site has been
3 designed based on published patterns rather than the real size of restricted fragments
4 obtained by sequence analysis.

5 Application of at least two restriction endonuclease enzymes is required in PRA
6 methods used for the identification of mycobacteria to species level. In this regard,
7 *Hae* III in combination with *BstE* II have been traditionally employed in the PRA of
8 441 bp *hsp65* gene. However, in this study the use of the *BstE* II enzyme resulted in
9 the production of identical fragment patterns for different species of mycobacteria,
10 e.g. *M. peregrinum* type II, *M. marnium* and *M. gordonae* type V. Thus, replacement
11 of this restriction enzyme by a more discriminative restriction enzyme seems
12 necessary.

13 In contrast, the nucleic acid sequencing of *hsp65* provided unambiguous results. Two
14 isolates which were previously described as *M. fortuitum* by conventional PCR and
15 identified as *M. chitae* type I by means of PRA, were clearly assigned as *M.*
16 *mucogenicum* (99.3% similarity). The lack of overlap between *hsp65* sequences from
17 *M. fortuitum* and *M. conceptionense* was particularly valuable for the accurate
18 identification of these species. The novel patterns in Table 2 obtained by PRA were
19 confirmed by sequence analysis. Fish strain 277/3/01 was a 99.5% match with *M.*
20 *arupense*, a novel species of mycobacteria that has recently been described (Cloud et
21 al., 2006). Masaki et al. (2006) have also recently reported two human isolates of this
22 species, in Japan. However, strain 277/3/01, isolated from fish in Slovenia and using
23 the GenoType *Mycobacterium* assay (Hain, Lifescience) had been identified as *M.*
24 *gordonae* (Pate et al., 2005). This finding may highlight the zoonotic significance of

1 this newly emerging *Mycobacterium* species. Alternatively, with the assumption of
2 environmental contamination for the isolate 277/3/01, this may provide evidence for
3 an environmental source of *M. arupense*.

4 All field and reference strains of *M. fortuitum* had an identical *hsp65* sequence;
5 whilst, the ITS sequences of isolates S7 and 277/2/01 were only partially matched
6 with this species and the sequence of the type strains could not be determined by
7 direct sequencing of PCR amplicons (Pourahmad, 2008). The snakehead fish isolate,
8 S7, has been previously identified as *M. poriferae* by means of HPLC and
9 biochemical tests (Tortoli et al., 1996). However, using PCR-RCBH, Puttinaowarat et
10 al. (Puttinaowarat et al., 2002) re-characterised this isolate as *M. fortuitum*. The
11 results of sequencing of the *hsp65* gene in this study were in agreement with this
12 group. This observation may emphasise the intraspecies variability of ITS sequences
13 that have been reported, elsewhere (Gürtler et al., 2006; Roth et al., 1998; Yoon et al.,
14 1997). Nevertheless, sequence analysis of other phylogenetically important genes, i.e.,
15 the 16S rRNA and the *rpoB* genes warranted precise identification of the
16 aforementioned species.

17 The current *hsp65*-based protocol (441 bp fragment) yields inefficient amplification in
18 *M. chelonae* complex due to sequence variability in the primer-binding region (Khan
19 et al, 2005). Likewise, in this study the difficulty in amplification of presumed *M.*
20 *chelonae* (NCIMB 13533) and *M. chelonae* MT1900 which had a better match with
21 *M. salmoniphilum* indicated the drawback of the Telenti et al. fragment. This problem
22 as well as the uncertainty in differentiation of field and type strains of *M. marinum*
23 and *M. pseudoshottsii* may highlight the need to develop methods based on alternative
24 genetic targets with better resolution for differentiation of members of the *M.*

1 *chelonae* complex and *M. marinum* which are frequently isolated from fish (Bruno et
2 al., 1998; dos Santos et al., 2002; Kent et al., 2004).

3 The construction of a phylogenetic tree (Fig. 3) was useful for indicating when
4 isolates were either rapid or slow-growing mycobacteria. *Mycobacterium arupense*,
5 for instance, which has been characterised as an intermediate species between fast and
6 slowly growing mycobacteria (Cloud et al., 2006), though grouped with rapidly
7 growing mycobacteria, was located between these two major groups. Furthermore,
8 when an isolate could not be matched as a certain species, the phylogenetic tree
9 showed the relatedness of that isolate to a known species or group of mycobacteria.

10 In conclusion, PRA is simple, rapid and inexpensive method for the identification of
11 mycobacteria. However, sequence analysis of the *hsp65* gene has proven a useful
12 method to integrate into species identification to indicate both slowly- and rapidly-
13 growing species of aquatic mycobacteria. As sequencing facilities are becoming more
14 accessible, sequencing of this 441 bp fragment may increasingly be used instead of
15 PRA as a means of molecular identification of *Mycobacterium* species.

References

Alcaide, F., Richter, I., Bernasconi, C., Springer, B., Hagenau, C., Schulze-Robbeke, R., Tortoli, E., Martin R., Bottger, E.C. Telenti, A., 1997 Heterogeneity and clonality among isolates of *Mycobacterium kansasii*: implications for epidemiological and pathogenicity studies. J. Clin. Microbiol. 35, 1959-1964.

Brunello, F., Ligozzi, M., Cristelli, E., Bonora, S., Tortoli, E., Fontana, R., 2001 Identification of 54 Mycobacterial Species by PCR-Restriction Fragment Length Polymorphism Analysis of the *hsp65* Gene. J. Clin. Microbiol. 39, 2799-2806.

Bruno, D.W., Griffiths, J., Mitchell, C.G., Wood, B.P., Fletcher, Z.J., Drobniowski, F.A., Hastings T.S., 1998 Pathology attributed to *Mycobacterium chelonae* infection among farmed and laboratory-infected Atlantic salmon *Salmo salar*. Dis. Aquat. Org. 33, 101-109.

Cheunoy, W., Prammananan, T., Chaiprasert, A., Foongladda, S., 2005 Comparative evaluation of polymerase chain reaction and restriction enzyme analysis: Two amplified targets, *hsp65* and *rpoB*, for identification of cultured mycobacteria. Diag. Microbiol. Infect. Dis. 51, 165-171.

Chinabut, S., 1999 Mycobacteriosis and Nocardiosis. In *Fish Diseases and Disorders* (Woo P.T.K. & Bruno D.W., eds.), CAB International, New York, NY, pp 319-340.

Cloud, J.L., Meyer, J.J., Pounder, J.I., Jost, K.C., Jr., Sweeney, A., Carroll, K.C., Woods, G.L., 2006 *Mycobacterium arupense* sp. nov., a non-chromogenic bacterium isolated from clinical specimens. Int. J. Syst. Evol. Microbiol. 56, 1413-1418.

Devallois, A., Goh, K., Rastogi, N., 1997 Rapid identification of mycobacteria to species level by PCR- restriction fragment length polymorphism analysis of the *hsp65*

1 gene and proposition of an algorithm to differentiate 34 mycobacterial species. J.
2 Clin. Microbiol. 35, 2969-2973.

3 dos Santos, N.M.S., do Vale, A., Sousa, M.J., Silva, M.T., 2002 Mycobacterial
4 infection in farmed turbot *Scophthalmus maximus*. Dis.Aquat. Org. 52, 87-91.

5 Gürtler, V., Harford, C., Bywater, J., Mayall, B.C., 2006 Direct identification of
6 slowly growing *Mycobacterium* species by analysis of the intergenic 16S-23S rDNA
7 spacer region (ISR) using a GelCompar II database containing sequence based
8 optimization for restriction fragment site polymorphisms (RFLPs) for 12 enzymes. J.
9 Microbiol. Methods 64, 185-199.

10 Hafner, B., Haag, H., Geiss, H.K., Nolte, O., 2004 Different molecular methods for
11 the identification of rarely isolated non-tuberculous mycobacteria and description of
12 new *hsp65* restriction fragment length polymorphism patterns. Molecular. Cellular.
13 Probe. 18, 59-65.

14 Kent, M.L., Whipps, C.M., Matthews, J.L., Florio, D., Watral V., Bishop-Stewart,
15 J.K., Poort, M., Bermudez L., 2004 Mycobacteriosis in zebrafish (*Danio rerio*)
16 research facilities. Compar. Biochemist. Physiol. Part C: Toxicol. Pharmacol.138,
17 383-390.

18 Khan, I.U.H., Selvaraju, S.B., Yadav, J.S., 2005 Occurrence and characterization of
19 multiple novel genotypes of *Mycobacterium immunogenum* and *Mycobacterium*
20 *chelonae* in metalworking fluids. FEMS Microbiol. Ecol. 54, 329-338.

21 Kim, B.J., Lee, S.H., Lyu, M.A., Kim, S.J., Bai, G.H., Kim, S.J., Chae, G.T., Kim,
22 E.C., Cha, C.Y., Kook, Y.H. 1999 Identification of Mycobacterial Species by
23 Comparative Sequence Analysis of the RNA Polymerase Gene (*rpoB*). J. Clin.
24 Microbiol. 37, 1714-1720.

1 Kirschner, P., Springer, B., Vogel, U., Meier, A., Wrede, A., Kiekenbeck, M., Bange,
2 F., Bottger, E., 1993 Genotypic identification of mycobacteria by nucleic acid
3 sequence determination: report of a 2-year experience in a clinical laboratory.
4 J.Clin.Microbiol. 31, 2882-2889.

5
6 Kumar, S., Tamura, K. & Nei, M., 2004 MEGA3: Integrated software for Molecular
7 Evolutionary Genetics Analysis and sequence alignment. Brief. Bioinform. 5, 150-
8 163.

9
10 Li, X.J., Wu, Q.X., Zeng, X.S., 2003 Nontuberculous mycobacterial cutaneous
11 infection confirmed by biochemical tests, polymerase chain reaction-restriction
12 fragment length polymorphism analysis and sequencing of *hsp65* gene. Brit. J.
13 Dermatol. 149, 642-646.

14
15 Masaki, T., Ohkusu, K., Hata, H., Fujiwara, N., Iihara, H., Yamada-Noda, M., Nhung,
16 P.H., Hayashi, M., Asano, Y., Kawamura, Y., Ezaki, T., 2006 *Mycobacterium*
17 *kumamotonense* sp. nov. Recovered from Clinical Specimen and the First Isolation
18 Report of *Mycobacterium arupense* in Japan: Novel Slowly Growing,
19 Nonchromogenic Clinical Isolates Related to *Mycobacterium terrae* Complex.
20 Microbiol. Immunol. 50, 889-897.

21
22 McNabb, A., Eisler, D., Adie, K., Amos, M., Rodrigues, M., Stephens, G., Black,
23 W.A., Isaac-Renton, J., 2004 Assessment of Partial Sequencing of the 65-Kilodalton
24 Heat Shock Protein Gene (*hsp65*) for Routine Identification of *Mycobacterium*
25 Species Isolated from Clinical Sources. J. Clin. Microbiol. 42, 3000-3011.

26
27 Murcia, M.I., Tortoli, E., Menendez, M.C., Palenque, E., Garcia, M.J., 2006
28 *Mycobacterium colombiense* sp. nov., a novel member of the *Mycobacterium avium*

complex and description of MAC-X as a new ITS genetic variant. Int. J. Syst. Evol. Microbiol. 56, 2049-2054.

Nolte, O., Haag, H., Hafner, B., 2005 A mutation in the 65,000 Dalton heat shock protein gene, commonly used for molecular identification of non-tuberculous mycobacteria, leads to the misidentification of *Mycobacterium malmoense* as *Mycobacterium marinum*. Molecular. Cellular. Probe. 19, 275-277.

Pate, M., Jencic, V., Zolnir-Dovc, M., Ocepek, M., 2005 Detection of mycobacteria in aquarium fish in Slovenia by culture and molecular methods. Dis. Aquat. Org. 64, 29-35.

Puttinaowarat, S., Thompson, K.D., Kolk, A., Adams, A., 2002 Identification of *Mycobacterium* spp. isolated from snakehead, *Channa striata* (Fowler), and Siamese fighting fish, *Betta splendens* (Regan), using polymerase chain reaction-reverse cross blot hybridization (PCR-RCBH). J. Fish Dis. 25, 235-243.

Pourahmad, F., 2008 Molecular Detection and Identification of Aquatic Mycobacteria Ph.D. Thesis edn., University of Stirling, Stirling, UK.

Pourahmad, F., Thompson, K.T., Taggart, J. B., Adams, A., Richards, R. H., 2008 *Mycobacterium stomatepiae* sp. nov., a slowly growing non-chromogenic species isolated from fish. Int. J. Syst. Evol. Microbiol. In Press.

Rhodes, M.W., Kator, H., McNabb, A., Deshayes, C., Reyrat, J.M., Brown-Elliott, B.A., Wallace, R., Jr., Trott, K.A., Parker, J.M., Lifland, B., Osterhout, G., Kaattari, I., Reece, K., Vogelbein, W., Ottinger, C.A., 2005 *Mycobacterium pseudoshottsii* sp. nov., a slowly growing chromogenic species isolated from Chesapeake Bay striped bass (*Morone saxatilis*). Int. J. Syst. Evol. Microbiol. 55, 1139-1147.

1 Ringuet, H., koua-Koffi, C., Honore, S., Varnerot, A., Vincent, V., Berche, P.,
2 Gaillard, J.L., Pierre-Audigier, C., 1999 *hsp65* Sequencing for Identification of
3 Rapidly Growing Mycobacteria. J. Clin. Microbiol. 37, 852-857.

4 Rocha, A.D., Leite, C.D., Torres, H.M., de Miranda, A.B., Lopes, M.Q.P., Degrave,
5 W.M., Suffys, P.N., 1999 Use of PCR-restriction fragment length polymorphism
6 analysis of the *hsp65* gene for rapid identification of mycobacteria in Brazil. J.
7 Microbiol. Methods 37, 223-229.

8 Roth, A., Fischer, M., Hamid, M.E., Michalke, S., Ludwig, W., Mauch, H., 1998
9 Differentiation of Phylogenetically Related Slowly Growing Mycobacteria Based on
10 16S-23S rRNA Gene Internal Transcribed Spacer Sequences. J. Clin. Microbiol. 36,
11 139-147.

12 Smole, S.C., McAleese, F., Ngampasutadol, J., von Reyn, C.F., Arbeit, R.D., 2002
13 Clinical and Epidemiological Correlates of Genotypes within the *Mycobacterium*
14 *avium* Complex Defined by Restriction and Sequence Analysis of *hsp65*. J. Clin.
15 Microbiol. 40, 3374-3380.

16 Telenti, A., Marchesi, F., Balz, M., Bally, F., Bottger, E.C., Bodmer, T., 1993 Rapid
17 identification of mycobacteria to the species level by polymerase chain reaction and
18 restriction enzyme analysis. J. Clin. Microbiol. 31, 175-178.

19 Tortoli, E., Bartoloni, A., Bozzetta, E., Burrini, C., Lacchini, C., Mantella, A., Penati,
20 V., Simonetti, M.T., Ghittino, C., 1996 Identification of the newly described
21 *Mycobacterium poriferae* from tuberculous lesions of snakehead fish (*Channa*
22 *striatus*). Compar. Immunol. Microbiol. Infect. Dis. 19, 25-29.

1 Turenne, C.Y., Semret, M., Cousins, D.V., Collins, D.M., Behr, M.A., 2006
2 Sequencing of *hsp65* Distinguishes among Subsets of the *Mycobacterium avium*
3 Complex. J. Clin. Microbiol. 44, 433-440.

4 Turenne, C.Y., Tschetter, L., Wolfe, J., Kabani, A., 2001 Necessity of Quality-
5 Controlled 16S rRNA Gene Sequence Databases: Identifying Nontuberculous
6 *Mycobacterium* Species. J. Clin. Microbiol. 39, 3637-3648.

7 Ucko, M., Coloni, A., Kvitt, H., Diamant, A., Zlotkin, A., Knibb, W.R., 2002 Strain
8 Variation in *Mycobacterium marinum* Fish Isolates. Appl. Environ. Microbiol. 68,
9 5281-5287.

10 Whipps, C.M., Butler, W.R., Pourahmad, F., Watral, V.G., Kent, M.L., 2007
11 Molecular systematics support the revival of *Mycobacterium salmoniphilum* (ex Ross
12 1960) sp. nov., nom. rev., a species closely related to *Mycobacterium chelonae*. Int. J.
13 Syst. Evol. Microbiol. 57, 2525-2531.

14 Wilson, R.W., Steingrube, V.A., Bottger, E.C., Springer, B., Brown-Elliott, B.A.,
15 Vincent, V., Jost, K.C., Jr., Zhang, Y., Garcia, M.J., Chiu, S.H., Onyi, G.O.,
16 Rossmore, H., Nash, D.R., Wallace, R.J. Jr., 2001 *Mycobacterium immunogenum* sp.
17 nov., a novel species related to *Mycobacterium abscessus* and associated with clinical
18 disease, pseudo-outbreaks and contaminated metalworking fluids: an international
19 cooperative study on mycobacterial taxonomy. Int. J. Syst. Evol. Microbiol. 51, 1751-
20 1764.

21 Yoon, J.H., Lee, S.T., Kim, S.B., Goodfellow, M., Park, Y.H., 1997 Inter- and
22 intraspecific genetic analysis of the genus *Saccharomonospora* with 16S to 23S
23 ribosomal DNA (rDNA) and 23S to 5S rDNA internally transcribed spacer sequences.
24 Int. J. Syst. Evol. Microbiol. 47, 661-669.




Fig.1: A representative selection of PRA profiles of *hsp65* gene PCR products using *Bst*EII enzyme

Lanes: 1) *M. marinum* (NCIMB1297); 2) *M. fortuitum* (NCIMB 1294); 3) *M. gordonae* 79/02; 4) *M. fortuitum* 277/2/01; 5) '*M. angelicum*' (DSM 45057^T); 6) *M. gordonae* 126/1/03; 7) *Mycobacterium* sp. 32/02; 8) *M. marinum* 7/04; 9) *M. marinum* 131/03; 10) *M. fortuitum* 276/3/01; 11) *M. marinum* 5/9/03; 12) *M. arupense* 277/3/01; 13) *M. conceptionense* 42/04; 14) '*M. aemonae*' (DSM 45058^T); 15) *Mycobacterium* sp. 50/04, 16) *M. conceptionense* 11/02; 17) *M. marinum* 9/04; 18) *M. gordonae* 49/21/03




Fig.2: A representative selection of PRA profiles of *hsp65* gene PCR products using *HaeIII* enzyme

Lanes 1) *M. fortuitum* (NCIMB 1295); 2) *M. fortuitum* (NCIMB 1294); 3) *M. conceptionense* IoA1; 4) *M. conceptionense* IoA2; 5) *M. conceptionense* IoA3; 6) *M. conceptionense* IoA4; 7) *M. conceptionense* IoA6; 8) *M. conceptionense* IoA7; 9) *M. mucogenicum* S11; 10) *M. mucogenicum* S12; 11) *M. peregrinum* S13; 12) *M. conceptionense* S14




Fig.3: Phylogenetic tree of the *hsp65* gene of aquatic mycobacteria prepared by using the neighbour-joining method and Kimura's two-parameter distance correction model. The support of each branch, as determined from 1,000 bootstrap samples, is indicated by the value at each node (as a percentage). Bootstrap values less than 70% are not shown. *Mycobacterium* sp. DL049 was used as the out-group. The scale bar indicates a 1% difference in nucleotide sequences.

Table 1: Species of mycobacteria, source, and origin used in this study

Species*	Strain	Source	Origin
<i>M. conceptionense</i>	CIP 108544 ^{Tc}	Human	France
<i>M. pseudoshottsii</i>	NCTC 13318 ^T	Striped bass (<i>Morone saxatilis</i>)	USA
<i>M. shottsii</i>	NCTC 13215 ^T	Striped bass (<i>Morone saxatilis</i>)	USA
<i>M. stomatepiae</i>	DSM 45059 ^T	Barombi Mbo Cichlid (<i>Stomatepia mariae</i>)	UK
<i>M. chelonae</i>	NCIMB 13533	Atlantic salmon (<i>Salmo salar</i>)	UK
<i>M. fortuitum</i>	NCIMB 1294	Chinook salmon (<i>Oncorhynchus tshawytscha</i>)	Unknown
<i>M. fortuitum</i>	NCIMB 1295	Unknown fish	Unknown
<i>M. marinum</i>	NCIMB 1297 ^a	Unspeciated marine fish	USA
<i>M. peregrinum</i>	NCTC 10264 ^b	Human	Mexico
<i>M. senegalense</i>	NCTC 10956	Bovine lymph node	France
<i>M. marinum</i>	DL921110 ^d	Sea bass (<i>Dicentrarchus labrax</i>)	Denmark
<i>M. marinum</i>	DL150191	Sea bass (<i>Dicentrarchus labrax</i>)	Israel
<i>M. marinum</i>	SO020195	Red Drum (<i>Scianops ocellatus</i>)	Israel
<i>M. marinum</i>	SR180194	Rabbit fish (<i>Siganus rivulatus</i>)	Israel
<i>M. marinum</i>	SR250195-B	Rabbitfish (<i>Siganus rivulatus</i>)	Israel
<i>M. marinum</i>	DL180892	Sea bass (<i>Dicentrarchus labrax</i>)	Israel
<i>M. marinum</i>	DL240490	Sea bass (<i>Dicentrarchus labrax</i>)	Israel
<i>M. marinum</i>	“Hernandez”	Unknown fish	Germany
<i>M. marinum</i>	1521/68	Unknown	Germany
<i>M. marinum</i>	DL045	Thalassa (<i>Dicentrarchus labrax</i>)	Greece
<i>Mycobacterium</i> sp.	DL049	Thalassa (<i>Dicentrarchus labrax</i>)	Greece
<i>M. marinum</i>	L.L. ATCC	Unknown	USA
<i>M. chelonae</i>	29/02 ^e	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>M. fortuitum</i>	277/2/01	Three-spot gourami (<i>Trichogaster trichopterus</i>)	Slovenia
<i>M. fortuitum</i>	55/02	Sterlet (<i>Acipenser ruthenus</i>)	Slovenia
<i>M. fortuitum</i>	32/02	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>M. fortuitum</i>	276/3/01	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>M. fortuitum</i>	50/04	Aquarium water	Slovenia
<i>M. fortuitum</i>	11/02	Dwarf gourami (<i>Colisa lalia</i>)	Slovenia
<i>M. fortuitum</i>	276/5/01	Three-spot gourami (<i>Trichogaster trichopterus</i>)	Slovenia
<i>M. gordonae</i>	277/3/01	Guppy (<i>Poecilia reticulata</i>)	Slovenia
<i>M. gordonae</i>	79/02	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>M. gordonae</i>	126/1/03	Freshwater angelfish (<i>Pterophyllum scalare</i>)	Slovenia
<i>M. gordonae</i>	49/21/03	Tap water	Slovenia
<i>M. marinum</i>	5/9/03	Human	Slovenia
<i>M. marinum</i>	7/04	Catfish (<i>Corydoras</i> sp.)	Slovenia

Table 1 (cont.): Species of mycobacteria, source, and origin used in the project

Species*	Strain	Source	Origin
<i>M. marinum</i>	131/03	Three-spot gourami (<i>Trichogaster trichopterus</i>)	Slovenia
<i>M. marinum</i>	09/04	Platyfish (<i>Xiphophorus maculatus</i>)	Slovenia
<i>Mycobacterium</i> sp.	42/04	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>Mycobacterium</i> sp.	126/5/01	Freshwater angelfish (<i>Pterophyllum scalare</i>)	Slovenia
<i>Mycobacterium</i> sp.	276/7/01	Guppy (<i>Poecilia reticulata</i>)	Slovenia
<i>M. szulgai</i> / <i>M. intermedium</i>	54/02	Goldfish (<i>Carassius auratus</i>)	Slovenia
<i>M. fortuitum</i>	TB1 ^f	Siamese fighting fish (<i>Betta splendens</i>)	Thailand
<i>M. fortuitum</i>	TB40	Siamese fighting fish (<i>Betta splendens</i>)	Thailand
<i>M. fortuitum</i>	S7	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S11	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S12	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S13	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S14	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S16	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. fortuitum</i>	S18	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. marinum</i>	S4	Snakehead fish (<i>Channa striata</i>)	Thailand
<i>M. marinum</i>	2B	Unknown fish	UAE ^g
<i>M. marinum</i>	3A	Unknown fish	UAE
<i>M. fortuitum</i>	IoA1	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA2	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA3	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA4	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA5	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA6	Unknown fish	Unknown
<i>M. fortuitum</i>	IoA7	Unknown fish	Unknown
<i>M. chelonae</i>	MT1900	Atlantic salmon (<i>Salmo salar</i>)	UK
<i>M. chelonae</i>	T5-2 ^h	Rosy barb (<i>Puntius conchonius</i>)	UK
<i>M. chelonae</i>	T9	Lumpsucker (<i>Cyclopterus lumpus</i>)	UK
<i>M. chelonae</i>	T14	Yellow seahorse (<i>Hippocampus kuda</i>)	UK
<i>M. marinum</i>	T1-1	Lumpsucker (<i>Cyclopterus lumpus</i>)	UK
<i>M. marinum</i>	T1-2	Lumpsucker (<i>Cyclopterus lumpus</i>)	UK
<i>M. marinum</i>	T5-1	Rosy barb (<i>Puntius conchonius</i>)	UK
<i>M. marinum</i>	T16	Otjikoto tilapia (<i>Tilapia guinasana</i>)	UK
<i>M. stomatepiae</i>	T3	Barombi Mbo Cichlid (<i>Stomatepia mariae</i>)	UK
<i>M. stomatepiae</i> ⁱ	T4	Barombi Mbo Cichlid (<i>Stomatepia mariae</i>)	UK

^{*}, Names as received

^a, National Collection of Industrial and Marine Bacteria, Aberdeen, Scotland, UK

^b, National Collection of Type Cultures, UK

^c, Collection de l'Institut Pasteur

^d, Isolates from DL921110 to L.L. ATCC were kindly provided by Dr. Angelo Colorni from Israel Oceanographic and Limnological Research Ltd., National Centre for Mariculture, Eilat, Israel.

^e, Mycobacterial isolates from Slovenia were kindly provided by Dr. Mateja Pate from Veterinary Faculty Ljubljana, University of Ljubljana, Gerbiceva 60, 1115 Ljubljana, Slovenia

^f, Strains obtained from different countries and held at -70°C in the Institute of Aquaculture (IoA), University of Stirling

^g, United Arab Emirates

^h, Mycobacterial strains isolated from London Zoo Aquarium during this study

ⁱ, Pourahmad et al.(2008), article in press

The superscript letter "T" denotes Type strain.

Table 2: Summary of patterns obtained by PRA of *hsp65* for aquatic *Mycobacterium* spp.

Species* and/or isolate name	No. of isolates	PRA pattern (bp)		Results
		<i>HaeIII</i>	<i>BstEII</i>	
<i>M. conceptionense</i> CIP 108544 ^T	1	145, 120, 50	240, 120, 85	<i>M. fortuitum</i>
<i>M. pseudoshottsii</i> . NCTC 13318 ^T	1	145, 105, 80	240, 210	<i>M. marinum</i>
<i>M. shottsii</i> NCTC 13215 ^T	1	145, 105, 80	240, 210	<i>M. marinum</i>
<i>M. stomatepiae</i> DSM 45059 ^T	1	145, 130	330, 115	New pattern
<i>M. chelonae</i> NCIMB 13533	1	200, 60	310, 130	<i>M. chelonae</i>
<i>M. fortuitum</i> NCIMB 1294	1	145, 120, 50	240, 120, 85	<i>M. fortuitum</i>
<i>M. fortuitum</i> NCIMB 1294	1	145, 120, 50	240, 120, 85	<i>M. fortuitum</i>
<i>M. marinum</i> NCIMB 1297	1	145, 105, 80	240, 210	<i>M. marinum</i>
<i>M. peregrinum</i> NCTC 10264	1	140, 120, 100, 60	240, 210	<i>M. peregrinum</i> II ^b
<i>M. senegalense</i> NCTC 10956	1	145, 120, 50	240, 120, 85	<i>M. fortuitum</i>
<i>M. chelonae</i>	5	200, 60	310, 130	<i>M. chelonae</i>
<i>M. fortuitum</i>	18	145, 120, 50	240, 120, 85	<i>M. fortuitum</i>
<i>M. fortuitum</i> IoA5, S13	2	140, 120, 100, 60	240, 210	<i>M. peregrinum</i> II ^b
<i>M. fortuitum</i>	3	multiple	multiple	Mixed culture
<i>M. fortuitum</i> S11, S12	2	140, 85, 60	320, 115	<i>M. chitae</i> I
<i>M. gordonae</i> 79/02	1	130, 110	240, 210	<i>M. gordonae</i> V ^a
<i>M. gordonae</i> 126/1/03	1	155, 110, 60	240, 120, 105	<i>M. gordonae</i> I ^b
<i>M. gordonae</i> 49/21/03	1	200, 60	325, 130	<i>M. gordonae</i> IV ^c
<i>M. gordonae</i> 277/3/01	1	145, 60	330, 115	New pattern
<i>M. marinum</i>	22	145, 105, 80	240, 210	<i>M. marinum</i>
<i>M. stomatepiae</i>	2	145, 130	330, 115	New pattern
<i>M. szulgai</i> / <i>M. intermedium</i>	1	145, 130, 60	240, 130, 80	New pattern
<i>Mycobacterium</i> sp. 126/5/01	1	130, 10	Not digested	New pattern
<i>Mycobacterium</i> sp. DL049	1	150, 80, 65, 55	240, 210	New pattern

*, Names as received

^a, Published pattern of Telenti et al. (1993)

^b, Published pattern of Brunello et al. (2001)

^c, Published pattern of Devallois et al. (1997)

Table 3: Mycobacteria identified by sequencing of the *hsp65* PCR products and comparison of fragment length after digestion by *HaeIII* and *BstEII* with fragments deduced from sequence analysis

Sequencing (% of best match)	No. of isolates	PRA results	PRA pattern (bp)*		Sequence (bp)**	
			<i>HaeIII</i>	<i>BstEII</i>	<i>HaeIII</i>	<i>BstEII</i>
<i>M. marinum</i>	18	<i>M. marinum</i>	145, 105, 80	240, 210	145, 106, 78	231, 210
<i>M. shottsii</i> (100)	1	<i>M. marinum</i>	145, 105, 80	240, 210	145, 106, 78	231, 210
<i>M. pseudoshottsii</i> (100)	6	<i>M. marinum</i>	145, 105, 80	240, 210	145, 106, 78	231, 210
<i>M. fortuitum</i> (100)	5	<i>M. fortuitum</i>	145, 120, 50	240, 120, 85	145, 123, 58	231, 115, 80
<i>M. peregrinum</i> (100)	3	<i>M. peregrinum</i> II	140, 120, 100, 60	240, 210	145, 139, 98, 53	231, 210
ND	3	Multiple fragments	Multiple fragments	Multiple fragments	NS	NS
<i>M. conceptionense</i> (100)	16	<i>M. fortuitum</i> ***	145, 120, 50	240, 120, 85	139, 123, 58	231, 115, 80
<i>M. senegalense</i> (100)	1	<i>M. fortuitum</i> ***	145, 120, 50	240, 120, 85	139, 123, 58	231, 115, 80
<i>M. mucogenicum</i> (99.3)	2	<i>M. chitae</i> I	140, 85, 60	320, 115	139, 87, 58	325, 116
<i>M. gordonae</i> (99.7)	1	<i>M. gordonae</i> V	130, 110	240, 210	127, 110	231, 210
<i>M. gordonae</i> (99.8)	1	<i>M. gordonae</i> I	155, 110, 60	240, 120, 105	161, 110, 60	23, 115, 80
<i>M. gordonae</i> (99.5)	1	<i>M. gordonae</i> IV	200, 60	325, 130	161, 110, 60	325, 116
<i>M. chelonae</i> (100)	4	<i>M. chelonae</i>	200, 60	310, 130	197, 60, 58, 54	310, 131
<i>M. salmoniphilum</i> (100)	2	<i>M. chelonae</i>	200, 60	310, 130	197, 60, 58, 54	310, 131
New patterns for:						
<i>M. stomatepiae</i> (100)	3	DSM 45059 ^T , T3, T4	145, 130	330, 115	145, 127, 50	325, 116
<i>M. avium</i> (96.8)	1	<i>Mycobacterium</i> sp.126/5/03	130, 110	Not digested	127, 112	Not digested
<i>M. arupense</i> (99.5)	1	<i>M. gordonae</i> 277/3/01	145, 60	330, 115	145, 60	325, 116
<i>M. gordonae</i> (95.5)	1	<i>M. szulgai</i> // <i>M. intermedium</i>	145, 130, 60	240, 130, 80	145, 127, 60	231, 130, 80
<i>M. poriferae</i> (94)	1	<i>Mycobacterium</i> sp. DL049	145, 80, 65, 55	240, 210	140, 80, 58, 50	231, 210

ND, Not done;

NS, Not stated;

*, Estimation of fragment size obtained by agarose gel;

**, Estimation of fragment size deduced from sequences;

***, The type strains of *M. senegalense* and *M. conceptionense* have also had PRA patterns very similar to *M. fortuitum*.