

REVIEW

Open Access



Does plastic waste present an infection hazard? Current evidence, surveillance challenges and research priorities

Patrick Ken Kalonde^{1,2,3*}, Kestern Mkoola¹, Michael Ormsby⁴, Taonga Mwapasa⁵, Precious Innocent Mastala^{1,6}, Madalitso Mphasa¹, Katalina Tonthola¹, Fanuel Meckson Bickton^{1,7,8}, Peter Nambala¹, Tony Robertson⁹, Richard S. Quilliam¹⁰, Kondwani Chidziwisano⁵, Tracy Morse¹¹, Christopher M. Jones², Michelle C. Stanton², Peter M. Atkinson³, Marc Y.R. Henrion^{1,2} and Nicholas Feasey^{1,12}

Abstract

Unmanaged plastic waste is widespread in settings with limited waste collection, yet its contribution to infectious disease risk remains poorly understood. In this review, we synthesise current evidence on the human health implications of plastic waste, with a specific focus on environmentally persistent bacteria that are transmitted to humans through the faecal oral route. We examine studies documenting microbial colonisation of plastic surfaces, including pathogenic and antimicrobial-resistant bacteria, and assess how well existing human and environmental surveillance systems capture these risks. We also review recent developments in remote sensing for detecting waste accumulation and evaluate spatial and statistical approaches that could link mapped waste to human exposure or infection data. Drawing on this synthesis, we identify major gaps that limit assessment of plastic-associated disease risk and outline key research priorities: improving methods for mapping waste using remotely sensed imagery, characterising spatial and temporal patterns of waste accumulation, and developing exposure metrics that incorporate environmental transport pathways. This review provides a foundation for advancing One Health research on plastic waste and its potential role in disease transmission.

Keywords Plastic waste, Antimicrobial resistance, One Health, Environmental surveillance, Remote sensing

*Correspondence:

Patrick Ken Kalonde
pkalonde@mlw.mw

¹Malawi-Liverpool-Wellcome Programme, Kamuzu University of Health Sciences, Blantyre, Malawi

²Liverpool School of Tropical Medicine, Liverpool, UK

³Faculty of Science and Technology, Lancaster University, Lancaster LA1 4YR, UK

⁴School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow, UK

⁵Centre for Water, Sanitation, Health, and Appropriate Technology Development (WASHTED), University of Business and Applied Sciences, Blantyre, Malawi

⁶University of Arizona, Arizona, USA

⁷Department of Rehabilitation Sciences, School of Life Sciences and Allied Health Professions, Kamuzu University of Health Sciences, Blantyre, Malawi

⁸Department of Physiotherapy, School of Therapeutic Sciences, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

⁹School of Health and Wellbeing, University of Glasgow, Glasgow G12 8TB, UK

¹⁰Biological & Environmental Sciences, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA, UK

¹¹Department of Civil and Environmental Engineering, University of Strathclyde, Glasgow, UK

¹²The School of Medicine, University of St. Andrews, St. Andrews, UK



© The Author(s) 2026. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Background

In many low-income countries, limited capacity for waste collection and safe waste disposal remains a major challenge [1, 2]. This problem is acute in rapidly growing informal or unplanned settlements where communities often fall outside the reach of formal waste management systems, leaving residents with few alternatives but to discard waste into their surrounding environment [3]. As a result, unmanaged waste often accumulates in open spaces, especially along roads, near homes, schools, markets and drainage networks, ultimately forming persistent and unmanaged waste piles [4, 5]. These piles create potentially high-risk zones where human, animal, and environmental health intersect. This issue, thus, represents a One Health challenge, requiring an integrated approach [6, 7]. Yet, despite their scale and persistence, such waste sites remain largely overlooked in urban planning and public health discourse.

Plastic waste is often the most visible and persistent component of unregulated waste piles (Fig. 1), accounting for approximately 50% of the total surface area in many settings [8]. This dominance reflects a broader global trend: plastic production has surged to an estimate of 459 million tonnes annually (as of 2019), an increase of approximately 2,195% since 1950, when global production was around 20 million tonnes [9]. Much of this plastic comes from single-use packaging, agricultural films (thin plastic sheets used to cover soil or crops to retain moisture and control weeds), food wrappers,

personal care containers, and medical disposables, and is typically composed of polymers such as polyethylene (PE), polypropylene (PP), and polyethylene terephthalate (PET). These materials are lightweight, durable, chemically stable, hygienic, waterproof, versatile and relatively inexpensive to produce, qualities that have made plastic an indispensable part of modern life. From households to hospitals, farms to factories, plastics have become deeply embedded in the fabric of everyday living. The problem, however, lies in its afterlife. Plastics are poorly recycled, particularly in low- and middle-income countries (LMICs) where formal recovery systems are limited or absent. In Africa, for example, only about 4% of waste is recovered by informal recyclers [10]. Globally, it is estimated that 79% of plastic waste ends up in the environment after use [11]. Once discarded, plastic undergoes slow degradation through exposure to sunlight and oxygen, breaking down into smaller fragments that persist in soils, water bodies, and even the air, for decades [12, 13]. Its persistence, coupled with its ubiquity, makes plastic waste a growing concern for both environmental and public health.

Although there is growing evidence that plastic waste accumulates in areas with a high burden of infectious disease [14, 15], whether and how this waste contributes to transmission is not well understood, and this review addresses the question by pursuing two connected aims. First, we review the evidence linking plastic waste to human health, considering several proposed pathways,



Fig. 1 A typical waste-pile in a setting where waste disposal in the environment is practiced. The waste pile includes a mix of waste including plastics (most of them blue plastics which are given free of charge in the country shown, Malawi), food waste, organic waste and numerous other materials. Photograph taken by the lead author

including toxic emissions from open burning, chemical leaching, and the creation of mosquito breeding habitat, before concentrating on bacterial colonisation of plastic surfaces, where evidence that plastic waste can act as a reservoir for pathogenic bacteria is consistently reported. Second, we ask how bacteria-related risks from waste can be identified and tracked, assessing existing human and environmental surveillance methods. We specifically examined a unique role of remote sensing for mapping waste, and spatial statistical methods for linking it to human exposure. The first part of the review therefore sets out why bacterial risk should be prioritised, and the second part builds directly on this by considering how waste mapping and its linkage to human exposure have been approached to date, before setting out a clear research direction for addressing the gaps that remain. Although this review synthesises evidence from a range of settings, several of the observations and examples highlighted are informed by the authors' experience conducting clinical, environmental, and spatial surveillance research in Malawi.

Human health risks associated with plastic waste

Plastics are associated with a range of human health risks, but in many low- and middle-income countries the most immediate concerns stem from what happens after plastics are discarded (See Additional file 1: Table S1 for a summary of disposal practices, associated hazards, and possible health outcomes). Open burning remains common where waste services are limited [16], producing $PM_{2.5}$, dioxins, and other toxic by-products [17, 18]. These emissions contribute to respiratory inflammation and chronic disease, including asthma, chronic bronchitis and cardiovascular complications [19]. Such exposures typically occur in crowded environments where burning is used to reduce waste volume or supplement household fuel, bringing pollutants into direct contact with the most vulnerable residents.

When plastics accumulate rather than burn, they present a different set of hazards. Additives such as bisphenol A and phthalates [20, 21] leach into surrounding soils and water sources [22], contaminating shallow wells and surface water used for drinking, cooking and washing. This contamination intensifies during the rainy season, when runoff mobilises leachates and transports them into households and shared water points [23, 24]. These chemical risks are well recognised, but they represent only part of the health burden associated with the persistence of discarded plastics in the environments where people live.

Plastic items such as bottles, bags and containers often trap rainwater and create small but persistent aquatic microhabitats capable of supporting *Aedes* and *Anopheles* mosquitoes [25]. Vector-borne diseases account for

a substantial proportion of the global infectious disease burden [26], and mosquitoes require water for larval development at every stage [27]. Evidence from diverse settings links discarded plastics to enhanced mosquito breeding. During the 2000 dengue outbreak in El Salvador, households with water-holding plastics colonised by mosquito larvae experienced higher dengue incidence [28]. In many other locations, studies have simply documented immature *Aedes* stages developing in rainwater accumulated within discarded plastic containers, including reports from Bangladesh [29], Sri Lanka [30], India [31–33], Tanzania [34, 35], Brazil [36], Canada [37] and the United States [38]. Plastics are one contributor among many artificial containers, and mosquitoes also exploit water storage vessels and other receptacles that may provide more stable breeding conditions [39].

While some plastic-associated chemicals have also been shown experimentally to affect mosquito development, including shortened emergence times under specific laboratory conditions [40], it is microplastics that have attracted more attention as a potential modifier of mosquito biology. Microplastics (MPs), small plastic fragments now widespread in aquatic environments, are readily ingested by larvae, and laboratory studies have explored their effects on survival [41–43], body weight [41, 43, 44], wing morphology [43, 45], gut microbiome composition [44, 46] and insecticide susceptibility [44]. Most studies report minimal or inconsistent effects on survival and morphology, and it is important to note that many of these effects have been observed at MP concentrations far higher than those measured in typical field settings [45]. MPs can, however, alter larval gut microbiota with plausible consequences for digestion, immunity and pathogen susceptibility [46] and may adsorb insecticidal compounds, reducing their bioavailable concentration [44]. These mechanisms remain under active investigation [47] and, although relevant to the broader ecological context of plastic pollution, they are not considered further in this review.

The most consistent and concerning body of evidence relates to the capacity of discarded plastics to act as microbial habitats [48]. Plastic surfaces readily support multispecies biofilms, often termed the *plastisphere*, providing stable substrates on which pathogenic bacteria can persist [49]. Pathogens have been identified on plastics collected from peri urban environments [50], estuarine systems [51] and waste piles in informal settlements [52], and some retain virulence after prolonged environmental exposure [53]. Plastics recovered from sewage, soils, beach debris, river systems and dumpsites frequently carry organisms with antimicrobial resistance (AMR) genes [50–52, 54]. Evidence from Ndirande in Blantyre demonstrates that waste pile plastics can harbour antimicrobial resistant bacteria at higher levels during the

wet season, when moisture promotes biofilm growth and microbial survival [52]. Horizontal gene transfer within biofilms provides a plausible mechanism for the accumulation of resistance determinants in these communities [55]. These observations indicate that plastics function as persistent environments in which pathogenic and antimicrobial resistant bacteria can persist, interact and amplify, particularly in settings where waste accumulates close to where people live.

Enteric disease remains a leading cause of death worldwide, with a disproportionate impact on children under five [56, 57], and antimicrobial resistance in enteric pathogens is an increasing contributor to this burden, with drug-resistant *Salmonella Typhi* now endemic in several African and South Asian settings [58, 59]. Against this background, the persistence of the plastisphere warrants scrutiny. Current evidence does not suggest that biofilm formation is specific to particular plastic polymers [60–64], and biodegradable plastics also support extensive microbial colonisation [65, 66]. Microbes colonise a wide range of discarded materials, including fabrics and organic waste [52], indicating that material type is not the determining factor. Plastics differ, however, in that they degrade extremely slowly, potentially sustaining microbial communities, including pathogens and AMR genes, for long periods and even as they fragment into microplastics [67]. For this reason, and because microbes show no preference for one substrate over another, large and unmanaged waste piles that contain substantial quantities of plastic should be viewed with particular concern. They create durable and accessible microbial reservoirs that can sustain pathogens, facilitate resistance exchange and support environmental dissemination in settings where waste management is limited. Of the hazards associated with plastic waste, bacterial colonisation therefore presents the clearest and most consistently documented risk to human health: the evidence linking plastics to bacterial persistence and AMR is more extensive than that available for other pathways, and the mortality burden of enteric disease specifically makes this risk pathway a priority for further investigation. The remainder of this review accordingly focuses on bacteria, and enteric bacteria in particular.

Monitoring and assessing health risk

Plastic waste has been found to harbour enteric and opportunistic bacterial pathogens including *Salmonella* spp., *Klebsiella pneumoniae*, *E. coli*, and potentially pathogenic *Vibrio* spp [50–52, 54, 60, 63, 64, 67–74]. Several isolates recovered from plastic waste exhibited antimicrobial resistance, although not all of members of these genera are pathogenic or AMR associated. Infections caused by these bacteria are responsible for major disease burden in settings with limited sanitation and

hygiene, with the majority of cases occurring in South Asia and Africa [75]. Outbreaks linked to multidrug resistant (MDR) bacterial strains have been reported in these regions [76]. Despite growing evidence that plastics harbour pathogenic and antimicrobial-resistant bacteria, the extent to which waste accumulation influences human exposure remains poorly characterised [77]. Investigation of the microbes associated with waste piles in the environment and exploring their links to humans can help clarify how plastic waste contributes to disease burden and how improvements in environmental conditions might translate into improvements in human health. The following section reviews what is known about the presence of human pathogens in the environment and approaches to human and environmental surveillance (see Additional file 1: Table S1).

Surveillance of pathogenic bacterial infections in humans

Surveillance of pathogenic bacteria and associated disease in human populations involves continuous and systematic collection, analysis and interpretation of health-related data [78]. Surveillance can be implemented in healthcare facilities or in the community, and is typically carried out either by Ministry of Health clinical and laboratory staff as part of routine care, or by research institutions running dedicated studies. Long-term, systematic bacteriological surveillance of the kind carried out by research institutions remains uncommon in low-income countries, and where it exists its sustainability and coverage depend on continued research funding rather than routine national systems. Where present, such programmes generate valuable longitudinal data for studying trends in bacterial infections [76]. Surveillance programmes, even in well-resourced healthcare settings typically prioritize monitoring overall prevalence over capturing their geographic distribution at a fine scale. Linking cases to specific places is straightforward only in settings with well-defined postal or addressing systems. Across much of the LMICs, formal address systems remain incomplete; in sub-Saharan Africa specifically, this has been linked to gaps in emergency service delivery and access to basic infrastructure [79]. Although substantial variation is likely to exist between countries and between urban and rural settings within the same country, health facilities often lack standardized conventions regarding the granularity at which patient locations are documented [80, 81]. In the absence of a formal addressing system, an alternative approach involves asking patients to identify their home locations directly on digital maps, an approach implemented by some initiatives [82, 83]. Research programmes can link facility-based records to residential locations or administrative units, enabling mapping of disease patterns across space [84–87]. Such linkages have been used to explore spatial

clustering and identify areas of elevated disease burden [88]. However, fine-scale geographical tracking of bacterial infections remains uncommon outside of dedicated research programmes.

A shortcoming of passive surveillance systems is their dependence on healthcare facility records that only capture those who seek care, by definition, they overlook asymptomatic carriers or paucisymptomatic individuals who do not seek health care [89–92]. This limitation is compounded by low rates of care-seeking itself. Across low- and middle-income countries, only around a third of diarrhoeal cases result in a visit to a hospital or clinic [93]. Barriers including distance to facilities, transport costs, and limited trust in or availability of care have been identified consistently across settings [94]. Passive surveillance therefore substantially undercounts disease burden, particularly among populations facing the greatest access barriers, such as those in peri-urban informal settlements. Such individuals may nonetheless act as reservoirs and contribute to ongoing transmission and, therefore, represent important targets for public health intervention, particularly during outbreaks. Surveillance can also be extended to apparently healthy individuals through community-based studies, however these “active” surveillance systems are more expensive [90]. These studies collect biological samples such as blood, stool, or urine [95–97], and in some cases, record point-level information such as GPS coordinates of the participants’ households [96]. These samples may be analysed using the same tests as those collected in healthcare facilities. Alternatively, serological assays can be employed to detect pathogen-specific antibodies, potentially (subject to the test parameters) capturing clinical, subclinical, and past infections. While the use of serology to quantify exposure in asymptomatic individuals was common during the COVID-19 pandemic, a single blood sample can be used to test for antibodies against multiple pathogens [98–100]. When linked with participant locations, serological data offer an opportunity to map geographical patterns of human exposure to pathogenic bacteria and to explore associations with waste, thereby strengthening surveillance.

Environmental surveillance of pathogenic bacteria

Environmental surveillance of pathogenic bacteria aims to determine where the bacteria occur in the environment, and how it persists under different conditions. It focuses on detecting and quantifying pathogens in environmental sources such as animals [101, 102], waste piles [52], soil [103, 104], surface water [105], wastewater [106] and river water [106–108]. Environmental samples are usually collected and transported to laboratories to isolate target bacteria using culture-based methods [71, 73, 109]. Isolates are then confirmed and characterised

using antibacterial susceptibility testing or molecular techniques such as quantitative PCR [71]. Environmental surveillance is a relatively low-cost approach for identifying circulating pathogens and strains [110], particularly in settings lacking systematic clinical surveillance [111, 112], and it poses fewer ethical and logistical challenges because individual-level consent is generally not required [113]. In contexts marked by inequality, it can capture information from areas poorly served by formal health systems or research teams [114] and has been recommended for integration into broader AMR monitoring programmes [115].

This general approach to environmental surveillance is not new, and has been applied to a wide range of enteric pathogens for decades, including *Vibrio cholerae*, rotavirus, hepatitis A virus, and non-polio enteroviruses, well before it became associated with the COVID-19 pandemic [116]. A systematic review of 100 pre-pandemic studies across 38 countries identified 25 distinct pathogen families under environmental surveillance [116]. Much of the field’s current methodology, however, was shaped specifically by global poliovirus eradication efforts [117–119], where environmental data guided vaccine strategies and assessed intervention impact [116]. More recently, wastewater surveillance gained prominence during the COVID-19 pandemic as a cost-effective method for tracking community infection trends [120–122], and has since expanded further to a wider array of viral, bacterial, and parasitic targets [123]. However, these approaches have largely relied on centralized wastewater infrastructure, which is limited or absent in many low- and middle-income countries, including much of sub-Saharan Africa [124]. In these settings, pit latrines are commonly used instead of sewer systems, and sometimes latrine contents may enter open drains and rivers through direct disposal, flooding, or other events that mobilise waste from containment systems [125, 126].

In settings without formal waste collection, domestic waste is discarded into the surrounding environment [4, 127, 128], creating alternative pathways for pathogen dissemination. Plastics compound this challenge as they are easily transported by wind or rainwater [129], can carry viable bacterial pathogens [68] for extended periods [53, 130], and are increasingly recognized as environmental vectors in their own right [131], yet their dispersal patterns remain poorly quantified. This mobility has significant implications for surveillance. A piece of plastic may carry its associated pathogens well beyond the point at which they were originally acquired, such that surveillance directed at a fixed point of origin is likely to underestimate the associated risk substantially. What is of greater relevance, therefore, is the location at which the plastic, and the pathogens it carries, ultimately accumulate. Spatial epidemiology follows the same principle at



Fig. 2 Waste pile dispersal: (a) waste accumulating into a pile before rainfall; (b) waste pile being dispersed by storm water

the population level. Rather than attempting to reconstruct the transmission history of individual cases, it seeks to determine whether disease occurrence or pathogen exposure exhibits spatial clustering and whether these patterns are associated with environmental hazards, reservoirs, or pathways of dispersal across the landscape [132]. Establishing where waste accumulates, and how this relates to patterns of human exposure, therefore constitutes a research priority. It is in this respect that the spatial sciences, encompassing both remote sensing and spatial statistics, assume particular importance. Remote sensing provides a means of mapping waste accumulation at scale. Fine-resolution aerial imagery, including that acquired by drones, may be used to train pattern-recognition models for the detection of waste [133–135], thereby supporting repeated mapping of waste accumulation and dispersal over time. Spatial statistical methods can subsequently be applied to test the association between mapped waste and human exposure, both of which are addressed in the sections that follow.

Remote sensing for monitoring of waste

Thus far, we have considered waste at the microscopic scale, focusing on pathogens that may pose risks to human health. Waste piles can act as reservoirs of pathogenic microbes, and it is now necessary to “zoom out” and consider the urban and peri-urban environments in which these piles accumulate and move. Identifying where waste builds up, where it is dispersed by storm water, and where it subsequently re-accumulates (Fig. 2) is essential for understanding potential exposure pathways. Remote sensing provides a means to capture these spatial patterns by using sensors mounted on satellites, aircraft, or drones [136] to collect spatially continuous observations of reflected light, enabling the mapping of waste across large areas and offering information that would otherwise be unavailable. This capability is

particularly valuable in settings with limited formal environmental monitoring systems, where ground-based observations are sparse or inconsistent.

Trends in environmental remote sensing

Remote sensing has evolved from black-and-white aerial photography to full-colour digital imagery captured from controlled angles by drones [134, 137, 138], aircraft [139–141], and satellites [142]. Early aerial photograph images required manual interpretation, but digital acquisition and advances in digital image processing and artificial intelligence now enable automated extraction of key information. A substantial body of research has demonstrated the utility of automated approaches for environmental mapping, including applications such as forest loss mapping [143] and tracking of urban expansion [144]. In contrast, environmental waste assessment has traditionally relied on field surveys centred on manual counting and weighing of waste materials [145], and in some cases, on characterizing the microbial composition of waste [52]. More recently, there has been growing interest in using aerial imagery to locate and quantify waste accumulating in the environment [133, 142, 146]. However, to advance environmental surveillance of waste, including plastic waste, the central question is: how can aerial imagery captured through remote sensing best support the mapping of waste, and potentially improve environmental and/or human disease surveillance? The following sections describe the characteristics of remote sensing data previously used for waste mapping, and with a specific focus on spatial and spectral characteristics.

Aerial data: spectral characteristics

Plastic waste can be identified in aerial imagery through its unique spectral properties. While the human eye perceives only visible light (blue: 450–485 nm; green:

500–565 nm; red: 625–750 nm), aerial sensors capture a broader spectral range, including the near- and short-wave-infrared regions, which can reveal materials that are visually indistinct. Laboratory studies show that plastics exhibit distinct reflectance values in the shortwave infrared (SWIR) region [139], and in some cases, the spectra can even differentiate polymer types [147]. Although capturing the unique SWIR region is not yet part of standard aerial imagery, these observations have the potential to guide the design of pragmatic sensors for mapping plastics accumulating in the environment. The thermal properties of plastics have been explored, but their utility remains limited because many surface materials share similar thermal spectra [148]. In contrast, multispectral satellites such as Sentinel-2 have demonstrated the ability to detect plastic waste in coastal environments, for example, in Scotland [149] and Greece [150]. Furthermore, benchmark datasets such as MARIDA have accelerated the development of models for mapping marine litter [151]. Comparable datasets for mapping waste in terrestrial and urban settings are lacking and need to be developed.

Aerial data: spatial resolution Identification of plastic waste from aerial images depends on the spatial resolution of the aerial data [8, 152]. Spatial resolution refers to the smallest discernible unit in an image. Satellite and aerial platforms collect imagery at a wide range of spatial resolutions. Landsat provides imagery at approximately 30 m resolution [153], Sentinel-2 offers a 10 m resolution [154], and Planet Labs' medium-resolution constellations acquire imagery at about 3 m resolution [155]. Commercial systems such as Pleiades [156] and Vantor (formerly Maxar) [157] provide sub-metre detail. Revisit times range from daily for systems such as Planet Labs [155] to every 16 days for Landsat [153]. Access to the imagery also differs. Sentinel-2 and Landsat data are freely available. Planet Labs provides some imagery for free to educational and research programmes, while fine-resolution products from Planet, Pleiades, and Vantor are commercially licensed. Some satellites can also be tasked to acquire imagery on specific dates. Most optical satellite imagery is affected by cloud cover, whereas radar systems such as Sentinel-1 SAR can penetrate clouds. This limitation has encouraged the use of optical sensing via drones, which enable flexible, cloud-free observations.

Drones can capture imagery with sub-centimetre detail and can do so flexibly and repeatedly, with the spatial resolution determined by the sensor and flight altitude [8, 152]. Low-altitude flights at about 10 to 20 m have been used to produce imagery in which individual plastic items were identifiable [158–165]. Although such high-resolution imagery can be visually interpreted, manual interpretation is labour-intensive and difficult to scale [166].

As a result, pattern-recognition methods are increasingly used to automate mapping, with studies showing success in detecting waste along beaches [134, 158, 159, 165, 167, 168] and in open water [138, 140, 146]. These settings have relatively uniform backgrounds, but in urban and peri-urban areas in low- and middle-income countries where waste disposal is common, very low flight altitudes are impractical because of presence of large trees, buildings, power lines and variable terrain. In these areas, waste accumulates in piles, and mapping such piles is important despite limited studies demonstrating how to map them effectively [133].

Since operating drones at very low altitudes might be unsafe and impractical in urban and peri-urban areas, it is important to investigate whether imagery collected at higher altitudes, with correspondingly lower spatial resolution, can still preserve the spatial texture required to reliably distinguish waste piles. Waste piles in informal settlements typically consist of heterogeneous mixtures of materials such as soil, organic matter, and plastics rather than isolated items [169], may exhibit spatial patterns that remain detectable even when individual objects cannot be resolved. Human interpreters may recognise these textural cues intuitively; assessing how consistently they do so is a necessary first step. Cohen's κ is used for agreement between two raters, whereas Krippendorff's α extends to more than two, and better accommodates varying resolutions and unclassified patches [170]. Reliable manual interpretation also provides a reference against which automated methods can be evaluated. Automating waste detection is likely to require models specifically trained to capture these textural patterns. Our own work using machine learning to identify waste based on groups of similar pixels suggests that this approach is promising [133], although further refinement is needed, including exploring deep learning methods for waste mapping. Detecting a waste pile and delineating its boundary are distinct aspects of performance. Precision and recall capture false positive and false negative rates, F1 balances the two, and Intersection over Union measures how closely predicted boundaries match reference ones. Reporting these across resolutions and sensors reveals the scales at which each deteriorates, which need not coincide.

Techniques for automatic mapping of plastics from aerial imagery

Automatic mapping of plastic waste from remotely sensed imagery has been performed using various automated approaches, which can be grouped broadly into three categories: index-based, machine learning, and deep learning techniques. Index-based methods combine spectral bands using mathematical formulae to highlight and identify plastic waste, and these indices have been

developed for uniform environments such as ocean surfaces [142, 171, 172]. Machine learning and deep learning are approaches that use identified examples of waste, allowing the algorithms to learn patterns associated with these examples and optimize the model to maximize the chances of correctly identifying waste. These approaches rely on training data, which are usually generated by manually digitizing waste to indicate its presence or outline [133, 137, 140, 159, 161, 165]. Such trained models reduce reliance on labour-intensive manual mapping [133, 141]. While traditional machine learning methods (such as random forest) require the pre-selection of spectral and textural features [173], deep learning architectures utilize multi-layered neural networks to automatically extract hierarchical patterns directly from the data [174], which might be important for mapping waste. Although machine-learning and deep-learning methods show promise for automating waste-pile mapping, their reliability is limited by the highly variable material composition of waste [133], so future research must develop more robust classifiers that can handle this heterogeneity and generalize effectively across diverse environmental contexts.

Linking environmental waste to human exposure to bacteria

Given the consistent evidence that bacteria readily colonise plastic surfaces, form persistent plastisphere biofilms and carry AMR genes, and in view of the substantial clinical burden posed by drug resistant bacterial infections [175, 176], the bacteria detected on plastics warrant close attention. The organisms repeatedly isolated from discarded plastics include *Vibrio cholerae*, *Salmonella enterica*, *Escherichia coli* and *Klebsiella* species. These pathogens are carried in the gastrointestinal tract and share a faecal-oral transmission pathway. At present, the strongest biological rationale for a link between environmental plastics and human health lies in the faecal-oral transmission of these enteric bacteria that colonise plastic surfaces, with the oral component reflecting human ingestion of organisms mobilised from contaminated waste. Although rodents and other pests on these waste piles can carry these pathogens to humans [177–179], the more likely and direct routes of exposure in informal urban settings involve contact with contaminated surfaces or the ingestion of water that mobilises bacteria from waste piles. At a basic level, therefore, it is important to determine whether waste sites or waste piles are associated with exposure to these pathogens in a way that allows assessment of whether the burden of enteric bacterial disease is linked to environmental plastics. This remains a notable gap. Establishing such a relationship would clarify whether reducing waste accumulation could contribute to measurable reductions in infection

burden. The following section outlines how such linkages can be examined and how spatial hotspots associated with waste can be identified, analysed and interpreted.

Conceptual need for linking environmental sources and human exposure

Historically, bacteria have been identified at species level, which limits the information available for understanding transmission dynamics. The advent of whole genome sequencing offers vastly increased resolution to define whether two bacterial isolates are genetically distinct or nearly identical. Ideally, the link between environmental waste and human infections should be established by comparing the genome sequence of pathogens detected in environmental samples with those found in clinical isolates or among asymptomatic individuals [180]. When human strains closely match environmental strains, it provides strong evidence of a flux between human and environmental compartments, although inferring the direction of transmission is yet more complicated [180–183]. Genomic analyses have been used to infer transmission networks by identifying highly similar *E. coli* strains and antimicrobial resistance determinants shared across humans, animals, and household environments, revealing hundreds of cross-compartment transmission events [184]. Apart from genomic similarity, these strains can also be examined in relation to geography; previous research in Blantyre showing that typhoid cases occurring close to one another and within the same river catchment often shared highly similar strains [88], suggesting that river systems may act as conduits for transmission. Nonetheless, mapping exposure populations alongside environmental features such as waste piles or river networks can reveal exposure hotspots and generate hypotheses about transmission routes, enabling targeted interventions to reduce risk. The relationships underlying such mapping are unlikely to hold uniformly across settings. The presence of waste piles, bacterial colonisation and its pathogenicity, the potential for water contamination, and the potential for human contact each depend on local geography, stormwater management, seasonality, and behaviour, and need not generalise beyond the setting in which they were derived. Transferability to other contexts should not be assumed, unless the model has been specifically developed to be sensitive to such geographic variation.

Methods for exploring associations across space

Matching strains in environmental and human samples indicates connectivity but not direction; contamination can flow from people to environment (shedding), from environment to people (exposure), or both. For exposed populations, it is important to recognize that not all exposures lead to illness [185], meaning many exposed

individuals are never captured in clinical case data. Geo-tagged disease cases can be aggregated at area level and aerial level models have been used previously to assess whether a high case-load relates to a high density of waste [86]. Whether exposure progresses to disease depends on factors such as pathogen dose and host immunity [186, 187]. Geotagged serological datasets may help bridge this gap by detecting markers of recent (IgM) or past (IgG) exposure, and they hold the potential to provide a more complete picture of where exposure occurs. It should be remembered that bacterial serological markers are intrinsically complex and this science continues to evolve.

On the environmental side, sources of pathogens may include waste piles [52], and their locations can be mapped [133]. When these datasets are combined, several point-level approaches can be used to test whether patterns of human exposure relate to mapped waste accumulation. These include statistical measures of association, where distance to waste is treated as a covariate, spatial clustering techniques, point-process models and geostatistical approaches.

Distance-based statistical associations

Statistical measures of association use distance to waste as a way to approximate the likelihood of exposure by assuming that interaction with waste is linked, in some way, to how close an individual is to a waste pile; for example, individuals who live nearer to waste piles are more likely to interact with them and are therefore more likely to be exposed. Because direct observational data on human-waste interactions are rarely available at scale, proximity might serve as a pragmatic proxy for potential exposure. In practice, this involves calculating the straight-line distance between each individual and the nearest waste pile [86], and incorporating this value into statistical models to examine whether the probability or intensity of exposure-related outcomes, such as geo-tagged serological markers of bacterial exposure, varies with distance. Continuous outcomes, including antibody concentrations, can be analysed using linear regression, generalized linear models [188], or random-effects models for repeated measurements [189], while binary outcomes such as seropositivity can be evaluated using logistic regression or mixed-effects logistic models. This distance-based approach provides a simple initial approach for testing whether closer proximity, and therefore greater opportunity for interaction with waste piles, is associated with bacterial exposure, and it can help generate picture to inform targeted interventions. This approach has several limitations, however. Straight-line distance assumes exposure risk declines uniformly in all directions from a waste pile, but terrain, drainage, and the routes people actually walk to reach water, markets, or schools are not accounted for. Waste piles are

also treated as static points, when in reality they grow, shrink, and disperse with rainfall and human activity, so a distance measured at one time point may not reflect exposure accumulated over the study period. Results are also sensitive to how distance is modelled and which covariates are included. Most importantly, association is not causation, and a significant distance effect does not establish that waste itself is the source of exposure, given the multiple, overlapping sources of contamination discussed earlier in this review.

Spatial clustering approaches

While a variety of spatial clustering methods exist, only a few are suitable for exploring the relationship between population exposure to pathogenic bacteria and environmental waste. Most clustering approaches, such as density-based clustering [190] and Kernel Density Estimation (KDE) [191, 192], are useful primarily for detecting spatial clusters but do not allow formal testing of associations with environmental drivers. In contrast, Kulldorff's Spatial Scan statistic searches for clusters by sliding circles of different sizes across the study area and evaluating each of these for unusually high case concentration [193]. This approach allows formal hypothesis testing by including covariates such as distance to waste in the model. Local Moran's I [194], and Getis-Ord G_i^* [195, 196] detect statistically significant spatial hotspots, providing confidence that observed clusters are unlikely to have arisen by chance. Although they do not provide a formal link with waste, a corresponding hotspot map of waste points can be generated, and the relationship between pathogen exposure and waste can then be assessed by examining spatial correlation or the degree of hotspot overlap. This overlap does not prove that waste is the direct cause of exposure. Waste and disease may both be linked to the same underlying factors, such as poverty, population density, or poor access to sanitation and water, and these shared factors alone can produce a spatial correlation between waste and disease even without waste itself being the source of exposure. Such analyses can therefore only suggest possible exposure routes, not confirm them; doing so would require accounting for these other factors or gathering exposure data directly from individuals.

Point-process models

Spatial point process models provide a robust framework for testing whether human exposure to pathogens is associated with waste locations. In this context, individuals with serological markers of recent or past exposure (e.g., IgM or IgG) or binary infection status are treated as marks on spatial points (i.e., individuals), while waste piles are represented as fixed environmental features. The intensity of exposed individuals can be modelled

as a function of proximity to waste, with distance-to-waste included as a covariate in inhomogeneous Poisson [197] or Cox process models [198]. This allows for testing whether exposure is more likely near waste sites than would be expected by chance, while accounting for the underlying spatial heterogeneity. When individual-level outcomes are continuous (antibody concentrations), marked point process models can incorporate these “marks” to assess how exposure levels vary with distance to waste. Additionally, second-order summary statistics, such as Cross- K functions or pair correlation functions, can detect clustering of exposed individuals around waste points beyond random expectation.

Geostatistical approaches

Exploration of associations using Geostatistics is similar to the statistical approaches mentioned previously, where exposure risk can be modelled as a function of proximity to waste included as a covariate. The key difference is that geostatistical models explicitly account for spatial dependence by incorporating a spatial covariance structure and spatially correlated random effects [199]. This allows adjustment for autocorrelation in the residuals among nearby individuals and enables predictions at unsampled locations through techniques such as Kriging [200]. These models can integrate continuous environmental surfaces and provide uncertainty estimates for predicted risk, offering a more nuanced understanding of the relationship between exposure and environmental hazards.

Limitations of proximity and the need for improved exposure metrics

Understanding human exposure to bacteria associated with waste is complex, and simple proximity measures do not fully capture how exposure occurs in real environments. While Euclidean distance offers a convenient way to represent how close individuals are to waste [86], it assumes that those who are nearer are proportionally more likely to be exposed. However, emerging evidence shows that pathogenic bacteria on waste materials can be transported by rainwater and surface runoff, with clustering of infections documented in downstream areas and, in some cases, directly linked to hydrological dispersal of contaminated waste [201]. These observations suggest that proximity alone may inadequately describe exposure risk.

Building on this, it is increasingly evident that exposure is shaped not only by how close people live to waste but also by the environmental pathways that move bacteria through the landscape. Direct interactions with waste piles certainly occur in many settings [202], yet hydrological processes can create additional exposure routes that operate independently of these behaviours. Downstream households, in particular, may be exposed through

storm-water driven transport of waste or through contamination of river water with bacteria shed from plastics [203]. In areas where piped water supplies are unreliable, river water is often used for domestic purposes [204], and such use has been linked with bacterial infection risk [205]. These hydrological pathways can amplify or redistribute exposure in ways that simple Euclidean cannot represent. For this reason, exposure metrics need to account not only for proximity but also for environmental transport processes such as terrain and drainage patterns, which influence where contaminated runoff is likely to accumulate. Incorporating these features provides a more realistic representation of potential exposure pathways and strengthens the basis for subsequent spatial analyses.

Both distance-based and hydrologically-informed exposure metrics remain silent on how people actually interact with waste. In many communities, residents actively sort through waste piles to recover materials for construction or resale; children are commonly observed playing in and around waste piles; and waste piles are frequently crossed as part of routine daily movement, for example to reach water points or markets. None of these behaviours would be captured by distance to the nearest waste pile, nor by models of hydrological transport, both of which treat exposure as a function of location rather than of what people actually do at that location. Capturing this would require direct behavioural or social science data, for example through structured observation or time-activity studies. Such methods carry their own limitations, including observer effects and the difficulty of capturing infrequent but high-risk interactions, but they offer a route to distinguishing active contact with waste from passive proximity to it, and represent an important complement to the spatial approaches discussed above.

Future research

The scientific community has established that discarded plastics readily develop a “plastisphere” [49], a biofilm capable of harbouring pathogenic and antimicrobial-resistant bacteria [60]. Earlier discussions of waste-related health risks also considered the possibility that plastics might influence disease transmission through vector-borne pathways. However, as summarised in this review, the evidence for mosquito-related mechanisms is inconsistent, highly context dependent, and often derived from controlled experimental studies whose real-world relevance is uncertain. In contrast, evidence that plastics act as stable substrates for pathogenic bacteria [52, 53, 74, 130] and some with antimicrobial-resistant genes [50–52, 54]. This evidence is stronger and more consistent. Much as we know that waste function as an environmental reservoir for harmful microbes, we still lack a clear understanding of how it accumulates

in the environment, how it persists and moves across space and time, and how these patterns relate to human exposure. Given the widespread presence of unmanaged waste, particularly in low- and middle-income countries [206, 207], and discovery of drug resistance pathogenic bacteria within waste piles accumulating in these settings [52], an important question arises: should waste accumulation be considered not only an environmental management issue but also a public health concern? Addressing this question requires first understanding how waste in the environment contributes to human exposure to pathogenic bacteria. Progress in this area will depend on detailed examination of where waste builds up, how it changes over time, and how these spatial and temporal patterns intersect with human activity. A clearer understanding of these dynamics will create opportunities to investigate exposure pathways and support the development of interventions that interrupt transmission. To advance this work, research should focus on three priorities:

Develop robust methods for mapping waste using remotely sensed imagery

Existing pattern-recognition methods have been validated on beaches and open water, where backgrounds are relatively uniform; no validated method exists for the heterogeneous, mixed-material waste piles typical of informal settlements. Advances in remote sensing have made it possible to acquire fine-resolution aerial imagery, and parallel developments in pattern recognition offer algorithms capable of detecting visual patterns in complex environments. However, waste piles often contain a mixture of plastics, organic matter, and other debris, and we still do not know how to have pattern recognition models for mapping these waste piles from remote sensing imagery. Research is needed to adapt existing pattern recognition algorithms and train and evaluate models for mapping waste. This work will provide foundational tools for systematically mapping waste from remotely sensed images and enabling establishment of spatial distribution of waste accumulation.

Observe and explain spatial and temporal patterns of waste accumulation

No study has tracked how waste piles form, persist or disperse over time in these settings, in contrast to the established literature on beach and marine litter dynamics. Although many studies describe the presence of unmanaged waste in developing-country settings, little is known about how waste accumulates, persists, and disperses across space and time. Waste may be redistributed by human activity, rainfall, flooding, blocked drainage, or informal burning, yet the mechanisms shaping these patterns remain poorly understood. Research should

characterise patterns of waste accumulation, and determine which areas are most likely to develop persistent accumulations that may serve as reservoirs for microbial communities. Understanding these patterns is essential for identifying where risks may be concentrated and how patterns of waste accumulation may influence potential exposure.

Develop novel methods for assessing human exposure to bacteria associated with waste

Distance measures capture neither hydrological transport nor active human contact with waste, as set out above; this priority addresses both. Existing spatial methods for analysing exposure are well established, but they were not designed to capture the epidemiological pathways that are most relevant for waste-associated bacteria. Simple proximity measures, such as straight-line distance, overlook the fact that waste and its associated pathogens can move through the environment. Hydrological processes, including storm water runoff, can transport waste materials and pathogenic bacteria downstream and far from their original location [208], creating exposure pathways that do not follow Euclidean distance. Evidence of infection clustering near rivers [88] and in downstream communities [201] further demonstrates the importance of these processes. Exposure is also shaped by how people actively engage with these water sources. Activities such as bathing, washing clothes, and collecting water for household use bring people into direct contact with rivers, drains, and other waste-affected water bodies, and the frequency of such contact varies with age and household role, with young children and women, who more often carry out water collection and washing tasks, potentially facing more frequent and direct exposure. There is therefore a need to evaluate how well current spatial approaches represent real environmental transport mechanisms and to develop new metrics that incorporate hydrological connectivity, and behavioural data on human-water and human-waste interaction, stratified by age and activity, and associated dispersal of pathogens. These approaches will help improve our understanding of spatio-temporal patterns of human exposure to pathogenic bacteria, and guide the design of more targeted One Health interventions.

Concluding remarks

Unmanaged waste, particularly plastics, is an emerging public health concern that has received limited attention in terms of the human infection risk they present. Plastics provide surfaces for microbial growth and may support the persistence and spread of antimicrobial resistance, yet their role in human infection is not well understood. Although pathogenic bacteria on waste are increasingly considered in environmental surveillance,

there are no established methods to assess the health risks they pose. Remote sensing offers a promising way to map waste accumulation, but most applications have focused on beaches and marine environments rather than settings with inadequate waste management such as informal settlements. This review makes three contributions in response: it brings evidence on plastic-associated bacterial risk, the state of surveillance practice, and advances in remote sensing together for the first time within a single One Health framework; in doing so, it identifies a specific gap between the ability to map waste and the ability to link it to human exposure; and it sets out research priorities intended to apply beyond the specific settings from which much of the evidence is drawn. The conditions we describe, constrained waste collection, unregulated latrine emptying, and the resulting mixing of pathogens from multiple environmental sources, are documented broadly across informal settlements in the LMICs, and the priorities identified are intended to generalise accordingly. Developing automated detection tools, studying patterns of waste dispersal, and identifying factors that influence these changes are essential to mapping and characterizing the patterns of waste accumulation and dispersal. Once this knowledge is established, new approaches to link human exposure to pathogens on waste through long-cycle transmission pathways should be developed to strengthen environmental surveillance and guide targeted interventions in resource-limited contexts.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s42522-026-00237-0>.

Supplementary Material 1

Acknowledgements

We acknowledge the support provided by the Liverpool School of Tropical Medicine, especially that provided by Mrs. Alison Derbyshire from the LSTM Library.

Author contributions

PKK conceptualised the study, led the literature review, synthesised the findings and drafted the initial manuscript. KN assisted with the literature review and contributed to writing the initial draft. MO provided early feedback and contributed to drafting and substantive editing. TM, MM, PIM, KT, FMB, PN, TR, and PMA reviewed the manuscript and provided critical edits. MCS, PMA, MYRH, and NF supervised the research and provided strategic guidance throughout the study. TR, RQ, KC, TM, CMJ, MCS, PMA, MYRH, and NF secured funding for the PhD studentship that supported this work. All authors read and approved the final manuscript and agreed to be accountable for all aspects of the research.

Funding

This research was funded in part by the Wellcome Trust through a strategic award to the Malawi Liverpool Wellcome Programme [grant number 206545/Z/17/Z]. For the purpose of open access, the author has applied for a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. The PhD studentship that led to this work was

supported by the UK Natural Environment Research Council (NERC) as part of the GCRF SPACES project [grant number NE/V005847/1].

Data availability

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 14 April 2026 / Accepted: 10 August 2026

Published online: 13 August 2026

References

1. Medina M. Solid wastes, poverty and the environment in developing country cities: Challenges and opportunities [WIDER Working Paper] [Internet]. Helsinki: The United Nations University World Institute for Development Economics Research (UNU-WIDER); 2010. WIDER Working Paper no.: 2010/23. Available from: <https://hdl.handle.net/10419/54107>
2. Bundhoo ZMA. Solid waste management in least developed countries: current status and challenges faced. *J Mater Cycles Waste Manag*. 2018;20(3):1867–77. <https://doi.org/10.1007/s10163-018-0728-3>.
3. Kekana HN, Ruhiiga TM, Ndou NN, Palamuleni LG. Environmental justice in South Africa: the dilemma of informal settlement residents. *GeoJournal*. 2023;88(4):3709–25. <https://doi.org/10.1007/s10708-022-10808-z>.
4. Ferronato N, Torretta V. Waste Mismanagement in Developing Countries: A Review of Global Issues. *Int J Environ Res Public Health*. 2019;16(6):1060. :10.3390/ijerph16061060 PubMed PMID: 30909625; PubMed Central PMCID: PMC6466021.
5. Vanapalli KR, Sharma HB, Ranjan VP, Samal B, Bhattacharya J, Dubey BK, et al. Challenges and strategies for effective plastic waste management during and post COVID-19 pandemic. *Sci Total Environ*. 2021;750:141514. <https://doi.org/10.1016/j.scitotenv.2020.141514> PubMed PMID: 32835961; PubMed Central PMCID: PMC7399665.
6. Panel (OHHLEP), OHHLE, Adisasmito WB, Almuhairei S, Behravesh CB, Bilivogui P, Bukachi SA, et al. One Health: A new definition for a sustainable and healthy future. *PLoS Pathog*. 2022;18(6):e1010537. <https://doi.org/10.1371/journal.ppat.1010537>.
7. Lerner H, Berg C. The concept of health in One Health and some practical implications for research and education: what is One Health? *Infect Ecol Epidemiol*. 2015;5(1):25300. <https://doi.org/10.3402/iee.v5.25300>. PubMed PMID: 25660757.
8. Kalonde P. Geospatial Methods for Mapping Domestic Waste Piles and Macro Plastics. Culminating Projects in Geography and Planning [Internet]. 2022 Jun 1. Available from: https://repository.stcloudstate.edu/gp_etds/13
9. Ritchie H, Roser M, Plastic, Pollution. Our World in Data [Internet]. 2018 Sep 1 [cited 2021 Oct 8]. Available from: <https://ourworldindata.org/plastic-pollution>.
10. Godfrey L, Ahmed MT, Gebremedhin KG, Katima JHY, Oelofse S, Osibanjo O et al. Solid waste management in Africa: governance failure or development opportunity? In: Regional development in Africa [Internet]. IntechOpen; 2019 [cited 2023 Nov 21]. Available from: <https://doi.org/10.5772/intechopen.86974>.
11. Geyer R, Jambeck JR, Law KL. Production, use, and fate of all plastics ever made. *Sci Adv*. 2017;3(7):e1700782. <https://doi.org/10.1126/sciadv.1700782>. PubMed PMID: 28776036; PubMed Central PMCID: PMC5517107.
12. Andrady AL. Persistence of plastic litter in the oceans. *Mar anthropogenic litter*. 2015;57–72.
13. Gewert B, Plassmann MM, MacLeod M. Pathways for degradation of plastic polymers floating in the marine environment. *Environ Sci: Processes Impacts*. 2015;17(9):1513–21. <https://doi.org/10.1039/C5EM00207A>.

14. Jambeck JR, Geyer R, Wilcox C, Siegler TR, Perryman M, Andrady A, et al. Plastic waste inputs from land into the ocean. *Science*. 2015;347(6223):768–71.
15. Law KL, Thompson RC. Microplastics in the seas. *Science*. 2014;345(6193):144–5. <https://doi.org/10.1126/science.1254065>.
16. Velis CA, Cook E. Mismanagement of Plastic Waste through Open Burning with Emphasis on the Global South: A Systematic Review of Risks to Occupational and Public Health. *Environ Sci Technol*. 2021;55(11):7186–207. <https://doi.org/10.1021/acs.est.0c08536>.
17. Wu D, Li Q, Shang X, Liang Y, Ding X, Sun H, et al. Commodity plastic burning as a source of inhaled toxic aerosols. *J Hazard Mater*. 2021;416:125820. <https://doi.org/10.1016/j.jhazmat.2021.125820>.
18. Islam MR, Welker J, Salam A, Stone EA. Plastic Burning Impacts on Atmospheric Fine Particulate Matter at Urban and Rural Sites in the USA and Bangladesh. *ACS Environ Au*. 2022;2(5):409–17. <https://doi.org/10.1021/acsenvironau.1c00054>.
19. Saha SC, Saha G. Effect of microplastics deposition on human lung airways: a review with computational benefits and challenges. *Heliyon*. 2024;10(2). <https://doi.org/10.1016/j.heliyon.2024.e24355>.
20. Hahladakis JN, Iacovidou E, Gerassimidou S. An overview of the occurrence, fate, and human risks of the bisphenol-A present in plastic materials, components, and products. *Integr Environ Assess Manag*. 2023;19(1):45–62.
21. Verma R, Vinoda KS, Papireddy M, Gowda ANS. Toxic Pollutants from Plastic Waste—A Review. *Procedia Environ Sci* 2016;Waste Manage Resource Utilisation35:701–8. <https://doi.org/10.1016/j.proenv.2016.07.069>.
22. Meng W, Sun H, Su G. Plastic packaging-associated chemicals and their hazards – An overview of reviews. *Chemosphere*. 2023;331:138795. <https://doi.org/10.1016/j.chemosphere.2023.138795>.
23. Chounlamany V, Tanchuling MA, Inoue T. Water quality and pollution loading of a river segment affected by landfill leachate and domestic waste. *Int J Environ Stud*. 2019;76(3):379–95.
24. Kjeldsen P, Barlaz MA, Rooker AP, Baun A, Ledin A, Christensen TH. Present and Long-Term Composition of MSW Landfill Leachate: A Review. *Crit Rev Environ Sci Technol*. 2002;32(4):297–336. <https://doi.org/10.1080/10643380290813462>.
25. Abdullah S, Jahidul Islam M, Hasan D, Sarkar MS, Rana SS, Das, et al. Impact of waste management on infectious disease control: evaluating strategies to mitigate dengue transmission and mosquito breeding sites – A systematic review. *J Angiotherapy*. 2024;8(8). <https://doi.org/10.25163/angiotherapy.889850>.
26. WHO. Vector-borne diseases [Internet]. 2024 [cited 2025 Mar 26]. Available from: <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>
27. Barrera R, Medialdea V. Development time and resistance to starvation of mosquito larvae. *J Nat Hist*. 1996;30(3):447–58.
28. Hayes JM, García-Rivera E, Flores-Reyna R, Suárez-Rangel G, Rodríguez-Mata T, Coto-Portillo R, et al. Risk factors for infection during a severe dengue outbreak in El Salvador in 2000. *Am J Trop Med Hyg*. 2003;69(6):629–33.
29. Paul KK, Dhar-Chowdhury P, Haque CE, Al-Amin HM, Goswami DR, Kafi MAH, et al. Risk factors for the presence of dengue vector mosquitoes, and determinants of their prevalence and larval site selection in Dhaka, Bangladesh. *PLoS ONE*. 2018;13(6):e0199457.
30. Surendran SN, Jayadas TT, Thiruchenthooan V, Raveendran S, Tharsan A, Santhirasegaram S, et al. Aedes larval bionomics and implications for dengue control in the paradigmatic Jaffna peninsula, northern Sri Lanka. *Parasites vectors*. 2021;14(1):1–13.
31. Banerjee S, Aditya G, Saha GK. Household Wastes as Larval Habitats of Dengue Vectors: Comparison between Urban and Rural Areas of Kolkata, India. *PLoS ONE*. 2015;10(10):e0138082. <https://doi.org/10.1371/journal.pone.0138082>.
32. Shukla A, Rajalakshmi A, Subash K, Jayakumar S, Arul N, Srivastava PK, et al. Seasonal variations of dengue vector mosquitoes in rural settings of Thiruvallur district in Tamil Nadu, India. *J Vector Borne Dis*. 2020;57(1):63–70.
33. Dasgupta S, Krishnan J, Ishtiaq F. Island biogeography and human practices drive ecological connectivity in mosquito species richness in the Lakshadweep Archipelago. *Sci Rep*. 2022;12(1):1–14.
34. Saleh F, Kitau J, Konradsen F, Alifrangis M, Lin CH, Juma S, et al. Habitat characteristics for immature stages of *Aedes aegypti* in Zanzibar city. *Tanzan J Am Mosq Control Association*. 2018;34(3):190–200.
35. Kampango A, Furu P, Sarath DL, Haji KA, Konradsen F, Schiøler KL, et al. Risk factors for occurrence and abundance of *Aedes aegypti* and *Aedes bromeliae* at hotel compounds in Zanzibar. *Parasites Vectors*. 2021;14(1):544. <https://doi.org/10.1186/s13071-021-05005-9>.
36. David MR, Lourenço-de-Oliveira R, de Freitas RM. Container productivity, daily survival rates and dispersal of *Aedes aegypti* mosquitoes in a high income dengue epidemic neighbourhood of Rio de Janeiro: presumed influence of differential urban structure on mosquito biology. *Memórias do Instituto Oswaldo Cruz*. 2009;104:927–32.
37. Giordano B, Gasparotto A, Liang P, Nelder M, Russell C, Hunter F. Discovery of an *Aedes* (*Stegomyia*) albopictus population and first records of *Aedes* (*Stegomyia*) *aegypti* in Canada. *Med Vet Entomol*. 2020;34(1):10–6.
38. Yitbarek S, Chen K, Celestin M, McCary M. Urban mosquito distributions are modulated by socioeconomic status and environmental traits in the USA. *Ecol Appl*. 2023;33(5):e2869.
39. Aziz AT, Dieng H, Ahmad AH, Mahyoub JA, Turkistani AM, Mesed H, et al. Household survey of container-breeding mosquitoes and climatic factors influencing the prevalence of *Aedes aegypti* (Diptera: Culicidae) in Makkah City, Saudi Arabia. *Asian Pac J Trop Biomed*. 2012;2(11):849–57. [https://doi.org/10.1016/S2221-1691\(12\)60242-1](https://doi.org/10.1016/S2221-1691(12)60242-1).
40. Valsala AGR, Asirvadani ED. Bisphenol A acts as developmental agonist in *Culex quinquefasciatus* Say. *Environ Sci Pollut Res*. 2022;29(49):74428–41. <https://doi.org/10.1007/s11356-022-21001-7>.
41. Al-Jaibachi R, Cuthbert RN, Callaghan A. Examining effects of ontogenic microplastic transference on *Culex* mosquito mortality and adult weight. *Sci Total Environ*. 2019;651:871–6.
42. McConnel G, Lawson J, Cañas-Carrell JE, Brelsfoard CL. The effects of nano- and microplastic ingestion on the survivorship and reproduction of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Environ Entomol*. 2024;53(4):594–603. <https://doi.org/10.1093/ee/nvae038>.
43. de Nadai BL, Moura L, Castro GB, Silva KJS, Maletzke AG, Corbi JJ, et al. Can microplastic contamination affect the wing morphology and wingbeat frequency of *Aedes aegypti* (Diptera: Culicidae) mosquitoes? *Environ Sci Pollut Res*. 2024;31(49):59289–301. <https://doi.org/10.1007/s11356-024-35161-1>.
44. Li Jhang, Liu X, hui, Liang G, rui, Gao H ting, Guo S han, Zhou X, yu et al. Microplastics affect mosquito from aquatic to terrestrial lifestyles and are transferred to mammals through mosquito bites. *Science of The Total Environment*. 2024;917:170547. <https://doi.org/10.1016/j.scitotenv.2024.170547>.
45. Thormeyer M, Tseng M. No Effect of Realistic Microplastic Exposure on Growth and Development of Wild-caught *Culex* (Diptera: Culicidae) Mosquitoes. *J Med Entomol*. 2023;60(3):604–7.
46. Edwards CC, McConnel G, Ramos D, Gurrola-Mares Y, Dhondiram Arole K, Green MJ, et al. Microplastic ingestion perturbs the microbiome of *Aedes albopictus* (Diptera: Culicidae) and *Aedes aegypti*. *J Med Entomol*. 2023;60(5):884–98. <https://doi.org/10.1093/jme/tjad097>.
47. Jones CM, Hughes GL, Coleman S, Fellows R, Quilliam RS. A perspective on the impacts of microplastics on mosquito biology and their vectorial capacity. *Med Vet Entomol*. 2024;38(2):138–47. <https://doi.org/10.1111/mve.12710>.
48. Maquart PO, Froehlich Y, Boyer S. Plastic pollution and infectious diseases. *Lancet Planet Health*. 2022;6(10):e842–5. [https://doi.org/10.1016/S2542-5196\(22\)00198-X](https://doi.org/10.1016/S2542-5196(22)00198-X) PubMed PMID: 36208647.
49. Zettler ER, Mincer TJ, Amaral-Zettler LA. Life in the plastisphere: microbial communities on plastic marine debris. *Environ Sci Technol*. 2013;47(13):7137–46.
50. Rasool FN, Saavedra MA, Pamba S, Perold V, Mmochi AJ, Maalim M, et al. Isolation and characterization of human pathogenic multidrug resistant bacteria associated with plastic litter collected in Zanzibar. *J Hazard Mater*. 2021;405:124591. <https://doi.org/10.1016/j.jhazmat.2020.124591>. PubMed PMID: 33301968.
51. Metcalf R, White HL, Moresco V, Ormsby MJ, Oliver DM, Quilliam RS. Sewage-associated plastic waste washed up on beaches can act as a reservoir for faecal bacteria, potential human pathogens, and genes for antimicrobial resistance. *Mar Pollut Bull*. 2022;180:113766.
52. Mphasa M, Ormsby MJ, Mwapasa T, Nambala P, Chidziwisano K, Morse T, et al. Urban waste piles are reservoirs for human pathogenic bacteria with high levels of multidrug resistance against last resort antibiotics: A comprehensive temporal and geographic field analysis. *J Hazard Mater*. 2025;484:136639. <https://doi.org/10.1016/j.jhazmat.2024.136639>.
53. Ormsby MJ, White HL, Metcalf R, Oliver DM, Quilliam RS. Clinically important *E. coli* strains can persist, and retain their pathogenicity, on environmental plastic and fabric waste. *Environ Pollut*. 2023;326:121466.
54. Yang Y, Li T, Liu P, Li H, Hu F. The formation of specific bacterial communities contributes to the enrichment of antibiotic resistance genes in the soil plastisphere. *J Hazard Mater*. 2022;436:129247.

55. Burmeister AR. Horizontal Gene Transfer. *Evol Med Public Health*. 2015;2015(1):193–4. <https://doi.org/10.1093/emph/eov.018> PubMed PMID: 26224621; PubMed Central PMCID: PMC4536854.
56. Black RE, Perin J, Yeung D, Rajeev T, Miller J, Elwood SE, et al. Estimated global and regional causes of deaths from diarrhoea in children younger than 5 years during 2000–21: a systematic review and Bayesian multinomial analysis. *Lancet Global Health*. 2024;12(6):e919–28. [https://doi.org/10.1016/S2214-109X\(24\)00078-0](https://doi.org/10.1016/S2214-109X(24)00078-0).
57. Piovani D, Figlioli G, Nikolopoulos GK, Bonovas S The global burden of enteric fever, 2017–2021: a systematic analysis from the global burden of disease study 2021. *eClinicalMedicine*. 2024;77. <https://doi.org/10.1016/j.eclinm.2024.102883>.
58. Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Aguilar GR, Gray A, et al. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*. 2022;399(10325):629–55. [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0) PubMed PMID: 35065702.
59. Naghavi M, Vollset SE, Ikuta KS, Swetschinski LR, Gray AP, Wool EE, et al. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *Lancet*. 2024;404(10459):1199–226. [https://doi.org/10.1016/S0140-6736\(24\)01867-1](https://doi.org/10.1016/S0140-6736(24)01867-1).
60. Yang LY, Huang XR, Neilson R, Li ZL, Yang XR, Su XX. Characterization of microbial community, ecological functions and antibiotic resistance in estuarine plastisphere. *Sci Total Environ*. 2023;161322.
61. Ganesan S, Ruendee T, Kimura SY, Chawengkijwanich C, Janjaroen D. Effect of biofilm formation on different types of plastic shopping bags: Structural and physicochemical properties. *Environ Res*. 2022;206:112542. <https://doi.org/10.1016/j.envres.2021.112542>. PubMed PMID: 34929185.
62. Boni W, Parrish K, Patil S, Fahrenfeld NL. Total coliform and *Escherichia coli* in microplastic biofilms grown in wastewater and inactivation by peracetic acid. *Water Environ Res*. 2021;93(3):334–42.
63. Lear L, Padfield D, Dowsett T, Jones M, Kay S, Hayward A, et al. Bacterial colonisation dynamics of household plastics in a coastal environment. *Sci Total Environ*. 2022;838:156199.
64. Lavery AL, Primpke S, Lorenz C, Gerds G, Dobbs FC. Bacterial biofilms colonizing plastics in estuarine waters, with an emphasis on *Vibrio* spp. and their antibacterial resistance. *PLoS ONE*. 2020;15(8):e0237704.
65. Saygin H, Baysal A. Biofilm formation of clinically important bacteria on bio-based and conventional micro/submicron-sized plastics. *Bull Environ Contam Toxicol*. 2020;105:18–25.
66. Juncheed K, Tanunchai B, Wahdan SFM, Thongsuk K, Schädler M, Noll M, et al. Dark side of a bio-based and biodegradable plastic? Assessment of pathogenic microbes associated with poly(butylene succinate-co-adipate) under ambient and future climates using next-generation sequencing. *Front Plant Sci*. 2022;13. <https://doi.org/10.3389/fpls.2022.966363>.
67. Forero-López AD, Brugnoli LI, Abasto B, Rimondino GN, Lassalle VL, Arduoso MG, et al. Plastisphere on microplastics: *In situ* assays in an estuarine environment. *J Hazard Mater*. 2022;440:129737. <https://doi.org/10.1016/j.jhazmat.2022.129737>.
68. Silva MM, Maldonado GC, Castro RO, de Sá Felizardo J, Cardoso RP, Dos Anjos RM, et al. Dispersal of potentially pathogenic bacteria by plastic debris in Guanabara Bay, RJ, Brazil. *Mar Pollut Bull*. 2019;141:561–8.
69. Rodrigues A, Oliver DM, McCarron A, Quilliam RS. Colonisation of plastic pellets (nurdles) by *E. coli* at public bathing beaches. *Mar Pollut Bull*. 2019;139:376–80. <https://doi.org/10.1016/j.marpolbul.2019.01.011>.
70. Tavşanoğlu ÜN, Başaran Kankılıç G, Akca G, Çirak T, Erdoğan Ş. Microplastics in a dam lake in Turkey: type, mesh size effect, and bacterial biofilm communities. *Environ Sci Pollut Res*. 2020;27(36):45688–98. <https://doi.org/10.1007/s11356-020-10424-9>.
71. Liang H, de Haan WP, Cerda-Domenech M, Méndez J, Lucena F, García-Aljaro C, et al. Detection of faecal bacteria and antibiotic resistance genes in biofilms attached to plastics from human-impacted coastal areas. *Environ Pollut*. 2023;319:120983.
72. Kirstein IV, Kirmizi S, Wichels A, Garin-Fernandez A, Erler R, Löder M, et al. Dangerous hitchhikers? Evidence for potentially pathogenic *Vibrio* spp. on microplastic particles. *Mar Environ Res*. 2016;120:1–8. <https://doi.org/10.1016/j.marenvres.2016.07.004>.
73. Quilliam RS, Jamieson J, Oliver DM. Seaweeds and plastic debris can influence the survival of faecal indicator organisms in beach environments. *Mar Pollut Bull*. 2014;84(1):201–7. <https://doi.org/10.1016/j.marpolbul.2014.05.011>.
74. Li H, Luo Q, Zhao S, Zhao P, Yang X, Huang Q, et al. Watershed urbanization enhances the enrichment of pathogenic bacteria and antibiotic resistance genes on microplastics in the water environment. *Environ Pollut*. 2022;313:120185. <https://doi.org/10.1016/j.envpol.2022.120185>.
75. Naghavi M, Abajobir AA, Abbafati C, Abbas KM, Abd-Allah F, Abera SF, et al. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390(10100):1151–210. [https://doi.org/10.1016/S0140-6736\(17\)32152-9](https://doi.org/10.1016/S0140-6736(17)32152-9).
76. Feasey NA, Gaskell K, Wong V, Msefula C, Selemani G, Kumwenda S, et al. Rapid emergence of multidrug resistant, H58-lineage salmonella Typhi in Blantyre, Malawi. *PLoS Negl Trop Dis*. 2015;9(4):e0003748. <https://doi.org/10.1371/journal.pntd.0003748>.
77. Beloe CJ, Browne MA, Johnston EL. Plastic debris as a vector for bacterial disease: an interdisciplinary systematic review. *Environ Sci Technol*. 2022;56(5):2950–8.
78. WHO. Ethics in public health surveillance [Internet]. [cited 2025 Nov 11]. Available from: <https://www.who.int/news-room/questions-and-answers/item/q-a-ethics-in-public-health-surveillance>
79. Bettencourt LMA, Marchio N. Infrastructure deficits and informal settlements in sub-Saharan Africa. *Nature*. 2025;645(8080):399–406. <https://doi.org/10.1038/s41586-025-09465-2>.
80. Baluwa C, Kalonde PK, Mseka UL, Tsoka O, Kajanga K, Chimphondah F, et al. Updated fine-scale location data for Blantyre district collected using a low-cost approach. *Sci Data*. 2025;12(1):1275. <https://doi.org/10.1038/s41597-025-05605-5>.
81. Mastala PI, Kalonde PK, Chikhola E, Bailey JB, Nkhoma J, Lohr F, et al. Challenges and potential opportunities for improving One Health surveillance in low-resource settings: Insights from rabies surveillance in Malawi. *One Health Outlook*. 2026;8(1):5. <https://doi.org/10.1186/s42522-025-00193-1>.
82. MacPherson P, Choko AT, Webb EL, Thindwa D, Squire SB, Sambakunsi R, et al. Development and Validation of a Global Positioning System–based Map Book System for Categorizing Cluster Residency Status of Community Members Living in High-Density Urban Slums in Blantyre, Malawi. *Am J Epidemiol*. 2013;177(10):1143–7. <https://doi.org/10.1093/aje/kws376>.
83. Baker S, Ali M, Deerin JF, Eltayeb MA, Cruz Espinoza LM, Gasmelseed N, et al. The Typhoid Fever Surveillance in Africa Program: Geospatial Sampling Frames for Household-based Studies: Lessons Learned From a Multicountry Surveillance Network in Senegal, South Africa, and Sudan. *Clin Infect Dis*. 2019;69(Supplement6):S474–82. <https://doi.org/10.1093/cid/ciz755>.
84. Khaki JJ, Meiring JE, Thindwa D, Henrion MYR, Jere TM, Msuku H, et al. Modelling Salmonella Typhi in high-density urban Blantyre neighbourhood, Malawi, using point pattern methods. *Sci Rep*. 2024;14(1):17164. <https://doi.org/10.1038/s41598-024-66436-9>.
85. Thindwa D, Chipeta MG, Henrion MYR, Gordon MA. Distinct climate influences on the risk of typhoid compared to invasive non-typhoid Salmonella disease in Blantyre, Malawi. *Sci Rep*. 2019;9(1):20310. <https://doi.org/10.1038/s41598-019-56688-1>.
86. Osei FB, Duker AA. Spatial dependency of *V. cholera* prevalence on open space refuse dumps in Kumasi, Ghana: a spatial statistical modelling. *Int J Health Geogr*. 2008;7:1–17.
87. Kalonde PK, Tsoka O, Chiepa B, Baluwa C, Nkolokosa C, Mategula D, et al. Mapping and quantifying travel time to define health facility catchment areas in Blantyre city in Malawi. *Commun Med*. 2025;5(1):227. <https://doi.org/10.1038/s43856-025-00845-3>.
88. Gauld JS, Olgemöller F, Heinz E, Nkhata R, Bilima S, Wailan AM, et al. Spatial and Genomic Data to Characterize Endemic Typhoid Transmission. *Clin Infect Dis*. 2022;74(11):1993–2000. <https://doi.org/10.1093/cid/ciab745>.
89. Shen Y, Liu Y, Krafft T, Wang Q. Progress and challenges in infectious disease surveillance and early warning. *Med Plus*. 2025;2(1):100071. <https://doi.org/10.1016/j.medp.2025.100071>.
90. Gibbons CL, Mangan MJ, Plass D, Havelaar AH, Brooke RJ, Kramarz P, et al. Measuring underreporting and under-ascertainment in infectious disease datasets: a comparison of methods. *BMC Public Health*. 2014;14(1):147. <https://doi.org/10.1186/1471-2458-14-147>.
91. Quilliam RS, Cross P, Williams AP, Edwards-Jones G, Salmon RL, Rigby D, et al. Subclinical infection and asymptomatic carriage of gastrointestinal zoonoses: occupational exposure, environmental pathways, and the anonymous spread of disease. *Epidemiol Infect*. 2013;141(10):2011–21. <https://doi.org/10.1017/S0950268813001131>.
92. Crump JA, Youssef FG, Luby SP, Wasfy MO, Rangel JM, Taalat M et al. Estimating the Incidence of Typhoid Fever and Other Febrile Illnesses in Developing Countries - 9, Number 5—May 2003 - Emerging Infectious Diseases journal - CDC [Internet]. 2003 May. <https://doi.org/10.1371/journal.pwat.0000067>

93. Wiens KE, Miller MH, Costello DJ, Solomon AP, Hilbert SM, Shipper AG et al. Care-seeking for diarrheal illness: a systematic review and meta-analysis. *Kulandaipalayam Natarajan, S, editor. PLOS Glob Public Health*. 2025;5(4):e0004483. <https://doi.org/10.1371/journal.pgph.0004483>.
94. Miller MH, Hilbert SM, Rosser EN, Sinko L, Lee EC, Wiens KE. Understanding influences of care-seeking behaviours for diarrhoeal illnesses: a qualitative meta-synthesis. *BMJ Open*. 2026;16(3):e109903. <https://doi.org/10.1136/bmjopen-2025-109903>.
95. Aiemjoy K, Seidman JC, Saha S, Munira SJ, Sajib MSI, Sium SMA, et al. Estimating typhoid incidence from community-based serosurveys: a multicohort study. *Lancet Microbe*. 2022;3(8):e578–87. [https://doi.org/10.1016/S2666-5247\(22\)00114-8](https://doi.org/10.1016/S2666-5247(22)00114-8) PubMed PMID: 35750069.
96. Bi Q, Azman AS, Satter SM, Khan AI, Ahmed D, Rijaz AA, et al. Micro-scale Spatial Clustering of Cholera Risk Factors in Urban Bangladesh. *PLoS Negl Trop Dis*. 2016;10(2):e0004400. <https://doi.org/10.1371/journal.pntd.0004400>.
97. Hellard ME, Sinclair MI, Hogg GG, Fairley CK. Prevalence of enteric pathogens among community based asymptomatic individuals. *J Gastroenterol Hepatol*. 2000;15(3):290–3. <https://doi.org/10.1046/j.1440-1746.2000.02089.x>.
98. Carcelen AC, Kong AC, Takahashi S, Hegde S, Jaenisch T, Chu M, et al. Challenges and Approaches to Establishing Multi-Pathogen Serosurveillance: Findings from the 2023 Serosurveillance Summit. *Am J Trop Med Hyg*. 2024;111(5):1145–52. <https://doi.org/10.4269/ajtmh.24-0296>. PubMed PMID: 39226906; PubMed Central PMCID: PMC11542533.
99. Nakagama Y, Komase Y, Candray K, Nakagama S, Sano F, Tsuchida T, et al. Serological Testing Reveals the Hidden COVID-19 Burden among Health Care Workers Experiencing a SARS-CoV-2 Nosocomial Outbreak. *Microbiol Spectr*. 2021;9(2):e01082–21. <https://doi.org/10.1128/Spectrum.01082-21>.
100. Theu JA, Kabaghe AN, Bello G, Chitsa-Banda E, Kagoli M, Auld A et al. SARS-CoV-2 Prevalence in Malawi Based on Data from Survey of Communities and Health Workers in 5 High-Burden Districts, October 2020 – 28, Supplement—December 2022 - Emerging Infectious Diseases journal - CDC [Internet]. 2022 Dec. <https://doi.org/10.3201/eid0905.020428>
101. Wallmann J. Monitoring of antimicrobial resistance in pathogenic bacteria from livestock animals. *Int J Med Microbiol*. 2006;296:81–6. <https://doi.org/10.1016/j.jimm.2006.01.064>. Risk Management for the Limitation of Antibiotic Resistance.
102. Wilson CN, Pulford CV, Akoko J, Sepulveda BP, Predeus AV, Bevington J, et al. Salmonella identified in pigs in Kenya and Malawi reveals the potential for zoonotic transmission in emerging pork markets. *PLoS Negl Trop Dis*. 2020;14(11):e0008796. <https://doi.org/10.1371/journal.pntd.0008796>.
103. Zhu D, Zhang Y, Zhu YG. Human pathogens in the soil ecosystem: Occurrence, dispersal, and study method. *Curr Opin Environ Sci Health*. 2023;33:100471. <https://doi.org/10.1016/j.coesh.2023.100471>.
104. Hashizume H, Taga S, Sakata MK, Hussein M, Siddig EE, Minamoto T, et al. Environmental detection of eumycetoma pathogens using multiplex real-time PCR for soil DNA in Sennar State, Sudan. *Trop Med Health*. 2023;51(1):71. <https://doi.org/10.1186/s41182-023-00563-3>.
105. Ottesen A, Kocurek B, Ramachandran P, Reed E, Commichaux S, Engelbach G et al. Advancing antimicrobial resistance monitoring in surface waters with metagenomic and quasimetagenomic methods. Charles KJ, editor. *PLOS Water*. 2022;1(12):e0000067. <https://doi.org/10.1371/journal.pwat.0000067>.
106. Barnes KG, Levy JJ, Gauld J, Rigby J, Kanjerwa O, Uzzell CB, et al. Utilizing river and wastewater as a SARS-CoV-2 surveillance tool in settings with limited formal sewage systems. *Nat Commun*. 2023;14(1):7883. <https://doi.org/10.1038/s41467-023-43047-y>.
107. Rigby J, Elmerhebi E, Diness Y, Mkwanda C, Tonthola K, Galloway H, et al. Optimized methods for detecting Salmonella Typhi in the environment using validated field sampling, culture and confirmatory molecular approaches. *J Appl Microbiol*. 2022;132(2):1503–17. <https://doi.org/10.1111/jam.15237>.
108. Cocker D, Mwapasa T, Grabic R, Grabicová K, Staňová AV, Chidziwisano K, et al. Environmental hazards from pollution of antibiotics and resistance-driving chemicals in an urban river network from Malawi. *npj Antimicrob Resist*. 2025;3(1):85. <https://doi.org/10.1038/s44259-025-00149-5>.
109. McConnell BR, Kraft AL, Durso LM, Ibekwe AM, Frye JG, Wells JE, et al. An analysis of culture-based methods used for the detection and isolation of Salmonella spp., Escherichia coli, and Enterococcus spp. from surface water: A systematic review. *Sci Total Environ*. 2024;927:172190. <https://doi.org/10.1016/j.scitotenv.2024.172190>.
110. Ngwira LG, Sharma B, Shrestha KB, Dahal S, Tuladhar R, Manthalu G, et al. Cost of wastewater-based environmental surveillance for SARS-CoV-2: Evidence from pilot sites in Blantyre, Malawi and Kathmandu, Nepal. *PLOS Glob Public Health*. 2022;2(12):e0001377. <https://doi.org/10.1371/journal.pgph.0001377>. PubMed PMID: 36962924; PubMed Central PMCID: PMC10021894.
111. Huijbers PMC, Flach CF, Larsson DGJ. A conceptual framework for the environmental surveillance of antibiotics and antibiotic resistance. *Environ Int*. 2019;130:104880. <https://doi.org/10.1016/j.envint.2019.05.074>.
112. Matrajt G, Lillis L, Meschke JS. Review of Methods Suitable for Environmental Surveillance of Salmonella Typhi and Paratyphi. *Clin Infect Dis*. 2020;71(Supplement2):S79–83. <https://doi.org/10.1093/cid/ciaa487>.
113. Gitter A, Ruvalcaba V, Clark K, Tran Carapucci T, Wu F, Hanson BM, et al. Turning analysis into action: opportunities and challenges in implementing wastewater science for public health decision-making. *Front Public Health*. 2025;13. <https://doi.org/10.3389/fpubh.2025.1562659>.
114. Holm RH, Osborne Jelks N, Schneider R, Smith T, Beyond. *Health Equity*. 2023;7(1):377–9. <https://doi.org/10.1089/heq.2022.0055>. PubMed PMID: 37351532; PubMed Central PMCID: PMC10282970. COVID-19: Designing Inclusive Public Health Surveillance by Including Wastewater Monitoring.
115. Baker KS, Jauneikaite E, Nunn JG, Midega JT, Atun R, Holt KE, et al. Evidence review and recommendations for the implementation of genomics for antimicrobial resistance surveillance: reports from an international expert group. *Lancet Microbe*. 2023;4(12):e1035–9. [https://doi.org/10.1016/S2666-5247\(23\)00281-1](https://doi.org/10.1016/S2666-5247(23)00281-1) PubMed PMID: 37977164.
116. Kilari P, Hill D, Anderson K, Collins MB, Green H, Kmush BL, et al. Wastewater Surveillance for infectious disease: a systematic review. *Am J Epidemiol*. 2023;192(2):305–22. <https://doi.org/10.1093/aje/kwac175>.
117. Duintjer Tebbens RJ, Zimmermann M, Pallansch MA, Thompson KM. Insights from a Systematic Search for Information on Designs, Costs, and Effectiveness of Poliovirus Environmental Surveillance Systems. *Food Environ Virol*. 2017;9(4):361–82. <https://doi.org/10.1007/s12560-017-9314-4>.
118. Berchenko Y, Manor Y, Freedman LS, Kaliner E, Grotto I, Mendelson E, et al. Estimation of polio infection prevalence from environmental surveillance data. *Sci Transl Med*. 2017;9(383):eaaf6786. <https://doi.org/10.1126/scitranslmed.aaf6786>.
119. Asghar H, Diop OM, Weldegebriel G, Malik F, Shetty S, El Bassioni L, et al. Environmental Surveillance for Polioviruses in the Global Polio Eradication Initiative. *J Infect Dis*. 2014;210(suppl1):S294–303. <https://doi.org/10.1093/infdis/jiu384>.
120. Mills C, Chadeau-Hyam M, Elliott P, Donnelly CA. The utility of wastewater surveillance for monitoring SARS-CoV-2 prevalence. *PNAS Nexus*. 2024;3(10):pgae438. <https://doi.org/10.1093/pnasnexus/pgae438>.
121. Morvan M, Jacomo AL, Souque C, Wade MJ, Hoffmann T, Pouwels K, et al. An analysis of 45 large-scale wastewater sites in England to estimate SARS-CoV-2 community prevalence. *Nat Commun*. 2022;13(1):4313. <https://doi.org/10.1038/s41467-022-31753-y>.
122. Schenk H, Rauch W, Zulli A, Boehm AB. SARS-CoV-2 surveillance in US wastewater: Leading indicators and data variability analysis in 2023–2024. *PLoS ONE*. 2024;19(11):e0313927. <https://doi.org/10.1371/journal.pone.0313927>.
123. Parkins MD, Lee BE, Acosta N, Bautista M, Hubert CRJ, Hruday SE, et al. Wastewater-based surveillance as a tool for public health action: SARS-CoV-2 and beyond. Forrest GN, editor. *Clin Microbiol Rev*. 2024;37(1):e00103–22. <https://doi.org/10.1128/cmr.00103-22>.
124. Adhikari S, Halden RU. Opportunities and limits of wastewater-based epidemiology for tracking global health and attainment of UN sustainable development goals. *Environ Int*. 2022;163:107217. <https://doi.org/10.1016/j.envint.2022.107217>.
125. Kumwenda S, Msefula C, Kadewa W, Ngwira B, Morse T. Estimating the Health Risk Associated with the Use of Ecological Sanitation Toilets in Malawi. *J Environ Public Health*. 2017;2017:1–13. <https://doi.org/10.1155/2017/3931802>.
126. Peal A, Evans B, Blackett I, Hawkins P, Heymans C. Fecal sludge management: a comparative analysis of 12 cities. *J Water Sanitation Hygiene Dev*. 2014;4(4):563–75. <https://doi.org/10.2166/wasdev.2014.026>.
127. Kalina M, Kwangulero J, Ali F, Abera YG, Tilley E. Where does it go? Perceptions and problems of riverine and marine litter amongst South Africa and Malawi's urban poor. *PLOS Water*. 2022;1(3):e0000013. <https://doi.org/10.1371/journal.pwat.0000013>.
128. Kalonde PK, Austin AC, Mandevu T, Banda PJ, Banda A, Stanton MC, et al. Determinants of household waste disposal practices and implications for practical community interventions: lessons from Lilongwe. *Environ Res: Infrastruct Sustain*. 2023;3(1):011003. <https://doi.org/10.1088/2634-4505/acbcc>.
129. Boni W, Parrish K, Patil S, Fahrenfeld NL. Total coliform and Escherichia coli in microplastic biofilms grown in wastewater and inactivation by peracetic acid. *Water Environ Res*. 2021;93(3):334–42. <https://doi.org/10.1002/wer.1434>.

130. Ormsby MJ, White HL, Metcalf R, Oliver DM, Feasey NA, Quilliam RS. Enduring pathogenicity of African strains of *Salmonella* on plastics and glass in simulated peri-urban environmental waste piles. *J Hazard Mater*. 2024;461:132439. <https://doi.org/10.1016/j.jhazmat.2023.132439>.
131. Ormsby MJ, Quilliam RS. The aquatic plastisphere: ecology, pathogen dissemination and antimicrobial resistance. *Nat Rev Microbiol*. 2026;24(7):446–62. <https://doi.org/10.1038/s41579-026-01301-2>.
132. Morrison CN, Mair CF, Bates L, Duncan DT, Branas CC, Bushover BR, et al. Defining Spatial Epidemiology: A Systematic Review and Re-orientation. *Epidemiology*. 2024;35(4):542–55. <https://doi.org/10.1097/EDE.0000000000001738>. PubMed PMID: 38534176; PubMed Central PMCID: PMC11196201.
133. Kalonde PK, Mwapa T, Mthawani R, Chidziwisano K, Morse T, Torguson JS, et al. Mapping waste piles in an urban environment using ground surveys, manual digitization of drone imagery, and object based image classification approach. *Environ Monit Assess*. 2025;197(4):374. <https://doi.org/10.1007/s10661-025-13675-6>.
134. Bak S, Hwang D, Kim H, Yoon H. Detection and monitoring of beach litter using Uav image and deep neural network. *international archives of the photogrammetry, remote sensing & spatial information sciences* [Internet]. 2019. Available from: <https://isprs-archives.copernicus.org/articles/XLII-3-W8/55/2019/>.
135. Gonçalves G, Andriolo U, Pinto L, Bessa F. Mapping marine litter using UAS on a beach-dune system: a multidisciplinary approach. *Sci Total Environ*. 2020;706:135742.
136. Chaminé HI, Pereira AJSC, Teodoro AC, Teixeira J. Remote sensing and GIS applications in earth and environmental systems sciences. *SN Appl Sci*. 2021;3(12):870. <https://doi.org/10.1007/s42452-021-04855-3>.
137. Gonçalves G, Andriolo U, Pinto L, Duarte D. Mapping marine litter with Unmanned Aerial Systems: A showcase comparison among manual image screening and machine learning techniques. *Mar Pollut Bull*. 2020;155:111158. <https://doi.org/10.1016/j.marpolbul.2020.111158>.
138. Andriolo U, Garcia-Garin O, Vighi M, Borrell A, Gonçalves G. Beached and floating litter surveys by unmanned aerial vehicles: Operational Analogies and differences. *Remote Sens*. 2022;14(6):1336.
139. Garaba SP, Aitken J, Slat B, Dierssen HM, Lebreton L, Zielinski O, et al. Sensing ocean plastics with an airborne hyperspectral shortwave infrared imager. *Environ Sci Technol*. 2018;52(20):11699–707. <https://doi.org/10.1021/acs.est.8b02855>.
140. Garcia-Garin O, Monleón-Getino T, López-Brosa P, Borrell A, Aguilar A, Borja-Robalino R, et al. Automatic detection and quantification of floating marine macro-litter in aerial images: Introducing a novel deep learning approach connected to a web application in R. *Environ Pollut*. 2021;273:116490. <https://doi.org/10.1016/j.envpol.2021.116490>.
141. Moy K, Neilson B, Chung A, Meadows A, Castrence M, Ambagis S, et al. Mapping coastal marine debris using aerial imagery and spatial analysis. *Mar Pollut Bull*. 2018;SI: Japanese Tsunami Debris132:52–9. <https://doi.org/10.1016/j.marpolbul.2017.11.045>.
142. Biermann L, Clewley D, Martinez-Vicente V, Topouzelis K. Finding Plastic Patches in Coastal Waters using Optical Satellite Data. *Sci Rep*. 2020;10(1):5364. <https://doi.org/10.1038/s41598-020-62298-z>.
143. Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, et al. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science*. 2013;342(6160):850–3. <https://doi.org/10.1126/science.1244693>.
144. Seto KC, Güneralp B, Hutyra LR. Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proc Natl Acad Sci USA*. 2012;109(40):16083–8. <https://doi.org/10.1073/pnas.1211658109>.
145. Viehman S, Vander Pluym JL, Schellinger J. Characterization of marine debris in North Carolina salt marshes. *Mar Pollut Bull*. 2011;62(12):2771–9.
146. Topouzelis K, Papakonstantinou A, Garaba SP. Detection of floating plastics from satellite and unmanned aerial systems (Plastic Litter Project 2018). *Int J Appl Earth Obs Geoinf*. 2019;79:175–83. <https://doi.org/10.1016/j.jag.2019.03.011>.
147. Cocking J, Narayanaswamy BE, Waluda CM, Williamson BJ. Aerial detection of beached marine plastic using a novel, hyperspectral short-wave infrared (SWIR) camera. *ICES J Mar Sci*. 2022;79(3):648–60.
148. Goddijn-Murphy L, Williamson B. On Thermal Infrared Remote Sensing of Plastic Pollution in Natural Waters. *Remote Sens*. 2019;11(18):18. <https://doi.org/10.3390/rs11182159>.
149. Page R, Lavender S, Thomas D, Berry K, Stevens S, Haq M, et al. Identification of Tyre and Plastic Waste from Combined Copernicus Sentinel-1 and -2 Data. *Remote Sens*. 2020;12(17):17. <https://doi.org/10.3390/rs12172824>.
150. Kremezi M, Kristollari V, Karathanassi V, Topouzelis K, Kolokoussis P, Taggio N, et al. Increasing the Sentinel-2 potential for marine plastic litter monitoring through image fusion techniques. *Mar Pollut Bull*. 2022;182:113974.
151. Kikaki K, Kakogeorgiou I, Mikeli P, Raitos DE, Karantzas K. MARIDA: A benchmark for Marine Debris detection from Sentinel-2 remote sensing data. *PLoS ONE*. 2022;17(1):e0262247.
152. Andriolo U, Topouzelis K, van Emmerik THM, Papakonstantinou A, Monteiro JG, Isobe A, et al. Drones for litter monitoring on coasts and rivers: suitable flight altitude and image resolution. *Mar Pollut Bull*. 2023;195:115521. <https://doi.org/10.1016/j.marpolbul.2023.115521>.
153. USGS. Landsat Satellite Missions [Internet]. 2020 [cited 2025 Nov 24]. Available from: <https://www.usgs.gov/landsat-missions/landsat-satellite-missions>
154. ESA. Sentinel-2 [Internet]. 2025 [cited 2025 Nov 24]. Available from: https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-2
155. Planet Labs. PlanetScope [Internet]. 2025 [cited 2025 Nov 24]. Available from: <https://docs.planet.com/data/imagery/planetscope/>
156. ESA. Pleiades [Internet]. 2025 [cited 2025 Nov 24]. Available from: <https://earth.esa.int/eogateway/missions/pleiades>.
157. Vantor. WorldView 2D [Internet]. 2025 [cited 2025 Nov 24]. Available from: <https://vantor.com/product/worldview/2d/>
158. Martin C, Parkes S, Zhang Q, Zhang X, McCabe MF, Duarte CM. Use of unmanned aerial vehicles for efficient beach litter monitoring. *Mar Pollut Bull*. 2018;131:662–73. <https://doi.org/10.1016/j.marpolbul.2018.04.045>.
159. Fallati L, Polidori A, Salvatore C, Saponari L, Savini A, Galli P. Anthropogenic Marine Debris assessment with Unmanned Aerial Vehicle imagery and deep learning: A case study along the beaches of the Republic of Maldives. *Sci Total Environ*. 2019;693:133581. <https://doi.org/10.1016/j.scitotenv.2019.133581>.
160. Wolf M, Berg K, van den, Garaba SP, Gnan N, Sattler K, Stahl F, et al. Machine learning for aquatic plastic litter detection, classification and quantification (APLATIC-Q). *Environ Res Lett*. 2020;15(11):114042. <https://doi.org/10.1088/1748-9326/abb0d1>.
161. Jakovljevic G, Govedarica M, Alvarez-Taboada F. A Deep Learning Model for Automatic Plastic Mapping Using Unmanned Aerial Vehicle (UAV) Data. *Remote Sens*. 2020;12(9):9. <https://doi.org/10.3390/rs12091515>.
162. Papakonstantinou A, Batsaris M, Spondylidis S, Topouzelis K. A Citizen Science Unmanned Aerial System Data Acquisition Protocol and Deep Learning Techniques for the Automatic Detection and Mapping of Marine Litter Concentrations in the Coastal Zone. *Drones*. 2021;5(1):1. <https://doi.org/10.3390/drones5010006>.
163. Pinto L, Andriolo U, Gonçalves G. Detecting stranded macro-litter categories on drone orthophoto by a multi-class Neural Network. *Mar Pollut Bull*. 2021;169:112594. <https://doi.org/10.1016/j.marpolbul.2021.112594>.
164. Merlino S, Paterni M, Berton A, Massetti L. Unmanned aerial vehicles for debris survey in coastal areas: Long-term monitoring programme to study spatial and temporal accumulation of the dynamics of beached marine litter. *Remote Sens*. 2020;12(8):1260.
165. Gonçalves G, Andriolo U, Gonçalves L, Sobral P, Bessa F. Quantifying Marine Macro Litter Abundance on a Sandy Beach Using Unmanned Aerial Systems and Object-Oriented Machine Learning Methods. *Remote Sens*. 2020;12(16):16. <https://doi.org/10.3390/rs12162599>.
166. Fraternali P, Morandini L, Herrera González SL. Solid waste detection, monitoring and mapping in remote sensing images: A survey. *Waste Manag*. 2024;189:88–102. <https://doi.org/10.1016/j.wasman.2024.08.003>.
167. Bao Z, Sha J, Li X, Hanchiso T, Shifaw E. Monitoring of beach litter by automatic interpretation of unmanned aerial vehicle images using the segmentation threshold method. *Mar Pollut Bull*. 2018;137:388–98.
168. Andriolo U, Gonçalves G, Bessa F, Sobral P. Mapping marine litter on coastal dunes with unmanned aerial systems: A showcase on the Atlantic Coast. *Sci Total Environ*. 2020;736:139632. <https://doi.org/10.1016/j.scitotenv.2020.139632>.
169. Mwapa T, Robertson T, Kazembe D, Mnkhwamba A, Kalonde PK, Feasey N, et al. Mapping and quantifying the spatial and temporal composition of waste piles in informal settlements of urban Malawi. *Environ Sci Pollut Res*. 2026 Feb;15. <https://doi.org/10.1007/s11356-026-37534-0>.
170. Kerr GHG, Fischer C, Reulke R. Reliability assessment for remote sensing data: Beyond Cohen's kappa. In: 2015 IEEE International Geoscience and Remote Sensing Symposium (IGARSS) [Internet]. Milan, Italy: IEEE; 2015 [cited 2026 Jul 9]. pp. 4995–8. Available from: <https://doi.org/10.1109/IGARSS.2015.7326954>.
171. Kremezi M, Kristollari V, Karathanassi V, Topouzelis K, Kolokoussis P, Taggio N, et al. Pansharpening PRISMA data for marine plastic litter detection using plastic indexes. *IEEE Access*. 2021;9:61955–71.

172. Sannigrahi S, Basu B, Basu AS, Pilla F. Development of automated marine floating plastic detection system using Sentinel-2 imagery and machine learning models. *Mar Pollut Bull*. 2022;178:113527.
173. Belgiu M, Drăguț L. Random forest in remote sensing: A review of applications and future directions. *ISPRS J Photogrammetry Remote Sens*. 2016;114:24–31. <https://doi.org/10.1016/j.isprsjprs.2016.01.011>.
174. Zhang L, Zhang L, Du B. Deep Learning for Remote Sensing Data: A Technical Tutorial on the State of the Art. *IEEE Geosci Remote Sens Mag*. 2016;4(2):22–40. <https://doi.org/10.1109/MGRS.2016.2540798>.
175. Feasey NA, Masesa C, Jassi C, Faragher EB, Mallewa J, Mallewa M, et al. Three Epidemics of Invasive Multidrug-Resistant Salmonella Bloodstream Infection in Blantyre, Malawi, 1998–2014. *Clin Infect Dis*. 2015;61(suppl4):S363–71. <https://doi.org/10.1093/cid/civ691>.
176. Naghavi M, Vollset SE, Ikuta KS, Swetschinski LR, Gray AP, Wool EE, et al. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *Lancet*. 2024;404(10459):1199–226. [https://doi.org/10.1016/S0140-6736\(24\)01867-1](https://doi.org/10.1016/S0140-6736(24)01867-1) PubMed PMID: 39299261.
177. Krystosik A, Njoroge G, Odhiambo L, Forsyth JE, Mutuku F, LaBeaud AD. Solid wastes provide breeding Sites, Burrows, and food for biological disease vectors, and Urban Zoonotic Reservoirs: a call to action for solutions-based research. *Front Public Health*. 2020;7. <https://doi.org/10.3389/fpubh.2019.00405>.
178. Hamond C, Browne AS, de Wilde LH, Hornsby RL, LeCount K, Anderson T, et al. Assessing rodents as carriers of pathogenic *Leptospira* species in the U.S. Virgin Islands and their risk to animal and public health. *Sci Rep*. 2022;12(1):1132. <https://doi.org/10.1038/s41598-022-04846-3>.
179. Buyukyavuz A, Northcutt JK, Dawson PL. Incidence of bacterial pathogens in flying insects collected near poultry farms. *J Appl Poult Res*. 2024;33(4):100462. <https://doi.org/10.1016/j.japr.2024.100462>.
180. Runcharoen C, Moradigaravand D, Blane B, Paksanont S, Thammachote J, Anun S, et al. Whole genome sequencing reveals high-resolution epidemiological links between clinical and environmental *Klebsiella pneumoniae*. *Genome Med*. 2017;9(1):6. <https://doi.org/10.1186/s13073-017-0397-1>.
181. Rocha J, Henriques I, Gomila M, Manaia CM. Common and distinctive genomic features of *Klebsiella pneumoniae* thriving in the natural environment or in clinical settings. *Sci Rep*. 2022;12(1):10441. <https://doi.org/10.1038/s41598-022-14547-6>.
182. Thorpe HA, Booton R, Kallonen T, Gibbon MJ, Couto N, Passet V, et al. A large-scale genomic snapshot of *Klebsiella* spp. isolates in Northern Italy reveals limited transmission between clinical and non-clinical settings. *Nat Microbiol*. 2022;7(12):2054–67. <https://doi.org/10.1038/s41564-022-01263-0>.
183. Dereeper Alexis G, Gaëlle P, Matthieu C, David B, Elodie B, Sylvaine, et al. Limited Transmission of *Klebsiella pneumoniae* among Humans, Animals, and the Environment in a Caribbean Island, Guadeloupe (French West Indies). *Microbiol Spectr*. 2022;10(5):e01242–22. <https://doi.org/10.1128/spectrum.01242-22>.
184. Musicha P, Beale MA, Cocker D, Oruru FA, Zuza A, Salifu C, et al. Transmission of extended spectrum β -lactamase-producing *Escherichia coli* and antimicrobial resistance gene flow across One Health compartments in eastern Africa: a whole-genome sequence analysis from a prospective cohort study. *Lancet Microbe*. 2026;7(1). <https://doi.org/10.1016/j.lanmic.2025.101224>. PubMed PMID: 41314227.
185. Koolman L, Prakash R, Diness Y, Msefula C, Nyirenda TS, Olgemoeller F, et al. Case-control investigation of invasive *Salmonella* disease in Malawi reveals no evidence of environmental or animal transmission of invasive strains, and supports human to human transmission. *PLoS Negl Trop Dis*. 2022;16(12):e0010982.
186. He X, Huang K. Chapter 7 - Assessment technologies for hazards/risks of wastewater. In: Ren H, Zhang X, editors. *High-Risk Pollutants in Wastewater* [Internet]. Elsevier; 2020. pp. 141–67. Available from: <https://doi.org/10.1016/B978-0-12-816448-8.00007-1>.
187. Katukiza AY, Rontetap M, van der Steen P, Foppen JWA, Lens PNL. Quantification of microbial risks to human health caused by waterborne viruses and bacteria in an urban slum. *J Appl Microbiol*. 2014;116(2):447–63. <https://doi.org/10.1111/jam.12368>. PubMed PMID: 24127653.
188. Nelder JA, Wedderburn RW. Generalized linear models. *J Royal Stat Soc Ser A: Stat Soc*. 1972;135(3):370–84.
189. Laird NM, Ware JH. Random-effects models for longitudinal data. *Biometrics*. 1982;38:307–25.
190. Ester M, Kriegel HP, Sander J, Xu X. A Density-based algorithm for discovering clusters in Large spatial databases with Noise. In: 1996. pp. 226–331. Available from: <https://ui.adsabs.harvard.edu/abs/1996kddm.conf.226E>.
191. Parzen E. On Estimation of a Probability Density Function and Mode. *Ann Math Statist*. 1962;33(3):1065–76. <https://doi.org/10.1214/aoms/1177704472>.
192. Rosenblatt M. Remarks on Some Nonparametric Estimates of a Density Function. *Ann Math Statist*. 1956;27(3):832–7. <https://doi.org/10.1214/aoms/1177728190>.
193. Kulldorff M. A spatial scan statistic. *Communications in Statistics - Theory Methods*. 1997;26(6):1481–96. <https://doi.org/10.1080/03610929708831995>.
194. Anselin L. Local indicators of spatial association—LISA. *Geographical Anal*. 1995;27(2):93–115.
195. Getis A, Ord JK. The analysis of spatial association by use of distance statistics. *Geographical Anal*. 1992;24(3):189–206.
196. Ord JK, Getis A. Local spatial autocorrelation statistics: distributional issues and an application. *Geographical Anal*. 1995;27(4):286–306.
197. Baddeley AJ, Møller J, Waagepetersen R. Non- and semi-parametric estimation of interaction in inhomogeneous point patterns. *Stat Neerl*. 2000;54(3):329–50. <https://doi.org/10.1111/1467-9574.00144>.
198. Ripley BD. Modelling Spatial Patterns. *Journal of the Royal Statistical Society Series. Stat Methodol*. 1977;39(2):172–92. <https://doi.org/10.1111/j.2517-6161.1977.tb01615.x>.
199. Diggle PJ, Tawn JA, Moyeed RA. Model-based geostatistics. *J Royal Stat Soc Ser C: Appl Stat*. 1998;47(3):299–350.
200. Diggle PJ, Giorgi E. Model-based geostatistics for global public health: methods and applications. Chapman and Hall/CRC; 2019.
201. Akullian A, Ng'eno E, Matheson AI, Cosmas L, Macharia D, Fields B, et al. Environmental Transmission of Typhoid Fever in an Urban Slum. *PLoS Negl Trop Dis*. 2015;9(12):e0004212. <https://doi.org/10.1371/journal.pntd.0004212>.
202. Jerie S. Occupational risks associated with solid waste management in the informal sector of Gweru, Zimbabwe. *J Environ Public Health*. 2016;2016(1):9024160.
203. Ormsby MJ, Woodford L, White HL, Fellows R, Oliver DM, Quilliam RS. Toxigenic *Vibrio cholerae* can cycle between environmental plastic waste and floodwater: Implications for environmental management of cholera. *J Hazard Mater*. 2024;461:132492. <https://doi.org/10.1016/j.jhazmat.2023.132492>.
204. Hove J, D'Ambruso L, Mabatha D, van der Merwe M, Byass P, Kahn K, et al. Water is life: developing community participation for clean water in rural South Africa. *BMJ Global Health*. 2019;4(3):e001377. <https://doi.org/10.1136/bmjgh-2018-001377>.
205. Gauld JS, Olgemoeller F, Nkhata R, Li C, Chirambo A, Morse T, et al. Domestic River Water Use and Risk of Typhoid Fever: Results From a Case-control Study in Blantyre, Malawi. *Clin Infect Dis*. 2020;70(7):1278–84. <https://doi.org/10.1093/cid/ciz405>.
206. Ozoh A, Longe B, Akpe V, Cock I. Indiscriminate solid waste disposal and problems with water-polluted urban cities in Africa. *J Coastal Zone Manage*. 2021;24(55):1000005.
207. Ferronato N, Torretta V. Waste mismanagement in developing countries: A review of global issues. *Int J Environ Res Public Health*. 2019;16(6):1060.
208. Silva MM, Maldonado GC, Castro RO, de Sá Felizardo J, Cardoso RP, Anjos dos RM, et al. Dispersal of potentially pathogenic bacteria by plastic debris in Guanabara Bay, RJ, Brazil. *Mar Pollut Bull*. 2019;141:561–8. <https://doi.org/10.1016/j.marpolbul.2019.02.064>.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.